Twenty Years of Cancer Incidence 1964-1983 The Ontario Cancer Registry

E.A. CLARKE
L.D. MARRETT
N. KREIGER

November 1987

THE ONTARIO CANCER TREATMENT AND RESEARCH FOUNDATION

ISBN 0-921196-00-8

This monograph is dedicated to Drs. A.H. Sellers and E.M. MacKay. They developed the concept of a population-based cancer registry "constructed out of documents previously unused or discarded after their primary purpose had been served". A debt of gratitude is owed to them and to the small group of individuals who ensured that their vision became reality.

TABLE OF CONTENTS

Page

Acknowledgements vii
Foreword ix
Index to Tables xi

CANCER REGISTRATION IN ONTARIO
- Introduction 1
- Background 1
- Establishment of the Ontario Cancer Registry 1
- Sources of Data 2
  - Hospital Discharges 2
  - Pathology Reports 2
  - Deaths 2
  - Treatment Centres and Registries 2
  - Other Sources 2
- Coding, Data Entry and Preprocessing of Data 3
- Creation of Cancer Incidence Data 4
  - Linkage of Data Sources 4
  - Allocation of Site, Histology and Other Information 5
- Variables in the Ontario Cancer Registry 7
- Advantages and Limitations of the Ontario Cancer Registry 7
- Conclusion 8

CANCER INCIDENCE IN ONTARIO
- Indices of Quality 11
- Interpretation of Data 11
- Explanatory Notes Regarding the Tables 11
- Calculation of Rates 12

Obtaining More Detailed Information from the Ontario Cancer Registry 14
References 15

TABLES
- Table 1.- ICD9 Code, abbreviated title, and full title for individual cancer sites 16
- Table 2.- Standard Populations 17
- Table 3.- Population of Ontario (in hundreds) 18
- Table 4.- Summary of Cancer Incidence 20
- Table 5.- Indices of Data Quality 22
- Site-Specific Tables 24

Index

Acknowledgements

The generous support of the Ontario Cancer Registry by the Government of the Province of Ontario through the Ontario Cancer Treatment and Research Foundation, together with the encouragement of the members of the Board is gratefully acknowledged. The special grant awarded by the Canadian Cancer Society Ontario Division was of considerable assistance in overcoming the initial difficulties of establishing an ongoing Registry.

The contents of this monograph are based on records received from the Ontario Ministry of Health, the Office of the Registrar General of the Ontario Ministry of Consumer and Commercial Relations, pathology laboratories in the many hospitals of Ontario, the Regional Cancer Centres of the Ontario Cancer Treatment and Research Foundation, and the Ontario Cancer Institute, incorporating the Princess Margaret Hospital. Without the support of these institutions and their staff, the Ontario Cancer Registry would not exist.

The authors wish to acknowledge the contribution of all the staff of the Ontario Cancer Treatment and Research Foundation associated with the Ontario Cancer Registry.

It is impossible to name all the individuals within the Division of Epidemiology and Statistics who have participated in the generation of incidence data for Ontario. However, three must be mentioned for their special contributions: Darlene Dale, Supervisor of Cancer Registry Records, whose attention to detail and knowledge of hospital practice have been invaluable; Mary Nemes, Manager of Health Records, whose knowledge of the history of the Registry was essential in order that the data were consistent; and Hannah Weir, Epidemiologist, whose perspective and meticulous care ensured the appropriate allocation of site and histology.

Without the staff of the Division of Information Systems, the development of the computer techniques in cancer registration would not have been possible. In particular, thanks are owed to Tom Broz, Project Manager, and his staff, including Judy Hum, Jill Chin and Jenny Gangadeen, under the management of June Murphy.

The assistance of Peter Toogood in the preparation of this monograph and the patience of Sheila Wing in typing and proof reading the many revisions of the text are acknowledged with special thanks.

Foreword

The mandate of the Ontario Cancer Treatment and Research Foundation includes "the adequate reporting of cases and the recording and compilation of data".

Initially confined to patients with cancer referred to the Ontario Cancer Treatment and Research Foundation, reporting in 1964 was expanded to include all cancer patients in the province. I am pleased to introduce this volume documenting 20 years of cancer incidence in Ontario. It is intended that this will be the first of a series of monographs of Cancer in Ontario. The Ontario Cancer Registry is a unique provincial resource for researchers, workers in health care, oncologists and others. I trust this volume will help them in their struggle against cancer.

J.W. Meakin, M.D., F.R.C.P. (C)
Executive Director

INDEX TO TABLES OF SITE SPECIFIC INCIDENCE FREQUENCIES AND RATES BY SEX

ICD9 CODE	TITLE	PAGE
140-208 (excl. 173)	ALL SITES, EXCLUDING NON-MELANOMA OF THE SKIN	24
140	LIP	28
141	TONGUE	32
142	SALIVARY GLAND	36
143	GUM	40
144	FLOOR OF MOUTH	44
145	MOUTH UNSPECIFIED	48
146	OROPHARYNX	52
147	NASOPHARYNX	56
148	HYPOPHARYNX	60
149	PHARYNX UNSPECIFIED	64
150	OESOPHAGUS	68
151	STOMACH	72
152	SMALL INTESTINE	76
153	COLON	80
154	RECTUM	84
155	LIVER	88
156	GALLBLADDER	92
157	PANCREAS	96
158	PERITONEUM	100
159	ILL-DEFINED DIGESTIVE	104
160	NOSE, SINUSES	108
161	LARYNX	112
162	LUNG	116
163	PLEURA	120
164	OTHER THORACIC ORGANS	124
165	ILL-DEFINED RESPIRATORY	128
170	BONE	132
171	CONNECTIVE TISSUE	136
172	MELANOMA OF SKIN	140
174/5	BREAST	144
179	UTERUS UNSPECIFIED	148
180	CERVIX UTERI	150
181	PLACENTA	152
182	BODY OF UTERUS	154

ICD9 CODE	TITLE	PAGE
183	OVARY	156
184	OTHER FEMALE GENITAL	158
185	PROSTATE	160
186	TESTIS	162
187	OTHER MALE GENITAL	164
188	BLADDER	166
189	KIDNEY, OTHER URINARY	170
190	EYE	174
191	BRAIN	178
192	NERVOUS SYSTEM UNSPECIFIED	182
193	THYROID	186
194	OTHER ENDOCRINE	190
200	LYMPHOSARCOMA	194
201	HODGKIN'S DISEASE	198
202	OTHER LYMPHOMAS	202
203	MULTIPLE MYELOMA	206
204	LYMPHOID LEUKAEMIA	210
205	MYELOID LEUKAEMIA	214
206	MONOCYTIC LEUKAEMIA	218
207	OTHER LEUKAEMIA	222
208	LEUKAEMIA, CELL TYPE UNSPECIFIED	226
195	ILL-DEFINED SITES	230
196	LYMPH NODES SECONDARY	234
197	RESPIRATORY/DIGESTIVE SECONDARY	238
198	OTHER SECONDARY	242
199	UNSPECIFIED SITE	246

Cancer Registration in Ontario

Introduction

This monograph is the first of a proposed series on cancer incidence, survival and mortality in Ontario. The occurrence of cancer is presented by year, site, sex and age for twenty years in Ontario. This information has not been published in a consolidated form elsewhere and supercedes data previously published for selected time periods.

The monograph is intended as a reference for researchers, health care workers and others. A description of the unique method of cancer registration employed in Ontario precedes the presentation of the data. An understanding of the techniques employed is necessary for interpretation of the data and, where necessary, the reader is cautioned as to possible limitations and artifacts.

Background

Ontario is the most populous province in Canada, with 9.1 million people in 1986[1] and an area of over one million square kilometers between Quebec to the east, Manitoba to the west, Hudson's Bay to the north and the United States to the south. Located between latitudes 41° - 57° north and longitudes 74° - 95° west, the altitude varies from sea level to 690 metres. Although 82% of the population now inhabit urban areas, mostly in the southern part of the province, previously there have been sizeable farming and mining communities. Manufacture of both durable and nondurable goods, trade and communications are the major industries. Eighty percent of Ontario residents were born in Canada but nevertheless represent a wide variety of ethnic groups; 60% are of British origin, 8% of French origin, 6% of Italian origin and 5% of German origin. In addition, there are about 84,000 native people in the province. The rate of emigration from Ontario is low. Of all persons residing in Ontario in 1976, only 4.0% had left the province by 1981. The rate of emigration for those more than 45 years of age, who are at greater risk of cancer, is even lower (1.4%)[2].

The Ontario Cancer Registry (OCR) is a population-based registry covering the entire province of Ontario. It is operated by the Ontario Cancer Treatment and Research Foundation, which was incorporated in 1943 by an Act of the Legislature of the Province of Ontario (The Cancer Act) "to establish a programme of cancer diagnosis, treatment and research" in the province. This act followed a recommendation by a provincial commission (the Cody Commission), that radiotherapy, then the most effective method of cancer treatment other than surgery, be centralized. Regional cancer centres (RCC) were therefore established to provide radiotherapy to outpatients and also included research activities. RCC are now located in Hamilton, Kingston, London, Ottawa, Sudbury, Thunder Bay, Windsor and Toronto. In addition, the Ontario Cancer Institute, incorporating the Princess Margaret Hospital (PMH), was established by statute in 1952 and in 1958 opened in Toronto with two research divisions. The PMH is the only hospital in Ontario dedicated exclusively to the care of cancer patients. Together, the RCC and the PMH provide all the radiation therapy for cancer patients in the province, as well as chemotherapy and consultative services for approximately 50% of cancer patients in Ontario.

The Ontario Cancer Treatment and Research Foundation, including the OCR, is supported primarily by the Ontario Ministry of Health (MOH). Patient care is publicly financed; in Ontario about 95% of Ontario residents are covered by a comprehensive government health insurance plan. While some residents of Ontario seek medical care outside the province, the proportion of claims for in-patient care originating from outside Ontario is less than 1%. The majority of such claims are made by residents of Ontario who live close to its borders.

The Cancer Act of 1943 included provision for "the adequate reporting of cancer cases and the recording and compilation of data". Cancer is not a legally reportable disease in Ontario, but amendments to the Cancer Act since 1943 have provided legal protection for organizations or individuals in the health care system who report information on cases of cancer to the Ontario Cancer Treatment and Research Foundation. These amendments enable information in the OCR to be used for epidemiological and medical research. In addition, each hospital in the province is required to forward diagnostic information on every discharged patient to the MOH. The MOH uses this information for administrative purposes and provides the OCR with copies of data on cancer patients; thus, a degree of compulsory reporting is in effect for hospitalized patients.

Establishment of the Ontario Cancer Registry

In 1936 Dr. A.H. Sellers, at that time with the Department of Health for Ontario - forerunner of Ministry of Health of Ontario - recognized that accurate data on the incidence and prevalence of cancer in Ontario were desirable and necessary. As in the United States, periodic surveys of cancer incidence were then undertaken. The first such survey took place in 1939. Physicians and hospitals in one county of Ontario registered each case of cancer seen in 1939, by completing a form giving residence, age, site and histology[3]. This survey was repeated in the same region in 1953[4].

In 1963 the Ontario Cancer Treatment and Research Foundation considered various methods of obtaining information necessary for the establishment and operation of a population-based cancer registry for Ontario. It was decided to use existing and available information contained in reports routinely prepared for other purposes, without requiring legislation that cancer be made a reportable disease. Initially, a manual system was established in which information from the following sources was recorded:

- hospital in-patient separation forms in which cancer was given as a diagnosis.

- abstracts of death registrations in which cancer was the underlying cause of death, as well as those in which cancer was mentioned.
- abstract cards used by the RCC and the PMH to record information on their cancer patients.
- reports from cancer registries maintained by the RCC.
- reports from hospital-based cancer registries.
- reports on biopsies performed on outpatients with cancer.
- drug prescriptions for needy cancer patients.
- out-patient radiotherapy reports.

After abstraction, coding and transcription of the information from the above sources to cards at the OCR, a manual linkage based on surname and given name was performed to bring together all records for an individual. The primary site of cancer was determined by a physician reviewing all records for each individual. Incidence data for the entire province were published for 1964 in 1968[5], for 1965 in 1970[6] and for 1966 in 1973[7].

The lack of timely incidence data indicated the necessity to streamline the process by implementing computer techniques. It was anticipated that this would permit a small cohesive group to handle a large number of records in a consistent manner and thus generate incidence data that were timely and of high quality.

Sources of Data

There have been major changes over time in the sources and quality of the data used for input to the Registry. The numbers of reports from the various sources used by the OCR are shown in Table A. Although many different sources were employed in the early years, the OCR has relied on only four major sources since 1972. They are:

- hospital separations with cancer as a diagnosis.
- pathology reports with a mention of cancer.
- death certificates in which cancer was the underlying cause of death.
- reports on patients referred to the RCC and PMH.

Hospital Discharges

Hospital inpatient separation data with mention of cancer are forwarded to the OCR by the MOH. These were submitted as documents until 1975, after which time the data were provided on magnetic tape. In 1978 the MOH instituted a requirement that each hospital submit an abstract for each discharge to an independent organization, the Hospital Medical Records Institute (HMRI). The HMRI abstract form provides for the recording of sixteen possible discharge diagnoses (as opposed to the single diagnosis previously permitted on hospital separation forms) but these abstracts do not contain surnames or given names. After processing (which includes some editing), HMRI forwards the resulting file to the MOH where name and Ontario Health Insurance Plan (OHIP) number are added. A subset of this integrated file is created consisting of records in which cancer is one of the discharge diagnoses and this file is forwarded annually to the OCR. The number of reports received in 1964 included patients whose admission to hospital occurred in 1963 or earlier. From 1965 onwards the number of cancer records submitted by hospitals has increased each year, with the exception of 1975 when the MOH changed from a manual to a computer system. The large increase in the number of records in 1978 is attributed to the capture of any record from the HMRI file in which any one of the sixteen diagnoses was cancer. In 1983, the increase is attributed to the inclusion of records in which a history of cancer is given.

Pathology Reports

In 1973 pathology laboratories across the province were asked to submit copies of reports in which cancer was mentioned. By 1980 all were complying. Paper records are provided to the OCR by participating laboratories and are coded by OCR staff.

Deaths

The OCR has data in machine-readable form on all deaths in Ontario residents since 1964. For the years 1964-1980, these data were received from Statistics Canada, by special arrangement with the Office of the Registrar General of Ontario. Since 1981, the Office of the Registrar General of Ontario has annually provided a computer tape directly to the OCR. Underlying cause of death is coded by trained nosologists in the Office of the Registrar General. Deaths with cancer considered to be the underlying cause have been included in the OCR since 1964. In addition, abstracts of death certificates with mention of cancer, provided by the Office of the Registrar General, have been included for the years 1964-1971.

Treatment Centres and Registries

Initially, abstract cards recording minimal information on their cancer patients were completed at each RCC and the PMH. Those from the RCC were forwarded to the OCR for further data abstraction and coding. Between 1964 and 1981, these cards were gradually discontinued at the PMH and the RCC and appropriate data were subsequently forwarded to the OCR in machine-readable form. Abstract cards were also created for tumour registries maintained at the RCC for cases diagnosed in their regions but not referred to the centres. These cards were forwarded to the OCR for abstracting and coding until the registries were discontinued by the RCC.

Other Sources

Pathology reports from hospital laboratories across the province duplicated the information contained in the biopsy reports on outpatients and were more comprehensive. The number of hospital-based registries in the province diminished due to financial constraints and use of these two sources by the OCR was discontinued in 1972. When incidence data from 1969 to 1971 were first generated and reviewed, it was realized that no cases were identified by radiotherapy out-patient reports per se and only 0.3% solely by the Foundation's Drug Service Programme. The identifying information on these sources was poor and it was decided to discontinue their use in 1972.

TABLE A

Reports Used in Registering Cases by Year and Source

Year	Hospital Discharge	Pathology	Death	RCC/PMH	RCC Registry	Hospital Registry	Biopsy Service	Drug Service	Outpatient Radiotherapy	Total
1964	23226	10	8831	6439	481	333	429	162	3	39914
1965	19033	11	9154	6534	984	864	366	206	2	37154
1966	21710	196	9380	6643	1143	836	541	191	18	40658
1967	25918	29	9849	7002	1285	3961	114	334	22	48514
1968	34974	11	11459	7127	1164	3901	13	605	6082	65336
1969	43411	158	11802	7736	2133	3565	1215	506	4456	74982
1970	40916	150	12191	8011	2291	31	449	2373	5858	72270
1971	42768	454	12203	8924	2113	24 [1]	259 [1]	481 [1]	6552 [1]	73778
1972	49699	908	11577	9610	2181	-	-	-	-	73975
1973	56153	11941	11794	9862	1703	-	-	-	-	91453
1974	60005	12519	12088	10720	2289	-	-	-	-	97621
1975	57350	13345	12224	11194	1974	-	-	-	-	96087
1976	61358	15032	12533	11997	1090 [2]	-	-	-	-	102010
1977	63430	15885	13100	12169	-	-	-	-	-	104584
1978	76870	20717	13428	14402	-	-	-	-	-	125417
1979	80257	26495	13961	13851	-	-	-	-	-	134564
1980	84056	28192	14217	14632	-	-	-	-	-	141097
1981	86792	34147	14854	15303	-	-	-	-	-	151096
1982	87098	35955	15306	16139	-	-	-	-	-	154498
1983	99991	43740	15869	17597	-	-	-	-	-	177197
Total	1115015	259895	245820	215892	20831	13515	3386	4858	22993	1902205

[1] Use of these records was discontinued after 1971.
[2] Use of these records was discontinued after 1976.

Coding, Data Entry and Preprocessing of Data

All cancer records submitted to the OCR in the early years (1964-1975), except death records in which cancer was reported as the underlying cause, were coded and entered into the computer centrally by the OCR. Between 1975 and 1977, hospital discharge information was coded at the MOH and from 1978 it has been coded in the medical records departments of hospitals in Ontario. These data have been sent to the OCR on magnetic tape since 1975. Given the fact that a passive system of cancer registration is employed, it is not possible, for the most part, to institute formal methods of quality control in regard to coding.

Pathology reports have always been coded and the data entered by clerks at the OCR. These are subjected to routine assessment of quality, as were other records previously coded at the OCR. Difficult reports are circulated among coding staff and are discussed at regular meetings with the medical staff.

Data from the RCC and PMH have been collected uniformly since their establishment. With computerization of records at these centres, coding has devolved to their medical record staffs. The Managers of Health Records at each RCC and the PMH meet twice a year to discuss coding and other quality control issues. The RCC also send copies of pathology reports and a clinical description of the cancer to the OCR. These reports are recoded, and any discrepancies are corrected after discussion between the RCC and the OCR.

Routine quality control of the data entry phase is carried out on all records of the OCR. Samples of reports entered on-line are verified by routine recoding and key entry. The data entry system requires that certain variables (e.g., surname of patient, date of diagnosis, site of disease) always be entered. As data are entered, they are edited for validity, consistency and plausibility. Data received on magnetic

tapes are also subjected to the same edits; however, these are carried out by batch programmes. Validity edits reject data which are inherently incorrect (e.g., the 13th month, the 32nd day). Consistency edits compare two or more data fields and report contradictions (e.g., a male patient with ovarian cancer, a treatment date preceding date of birth). Edits for plausibility report unlikely but possible situations which are potential errors (e.g., a 110 year old patient, a five year old male with prostatic cancer). These plausibility edits are checked manually and corrected if necessary. Coded data (e.g., residence, hospital, birthplace) are compared with tables constructed by the OCR specifically for validation purposes. Finally, numerical data are validated with check digits.

Site of cancer in abstracts of death certificates for the years 1964-1968 was originally coded to the Seventh Revision of the International Classification of Diseases (ICD)[8], and for 1969-1978 to the Eighth Revision[9]. From 1964 until 1979, site of cancer on all other records in the OCR was coded to the Eighth Revision. Site on all records since 1979 has been coded to the Ninth Revision of the ICD (ICD9)[10]. In addition to the computer edits described above, all ICD codes are converted during processing to the ICD9. Prior to 1979, morphology was coded to the Manual of Tumor Nomenclature and Coding (MOTNAC)[11] and, since 1979, to the International Classification of Diseases for Oncology (ICDO-M)[12]. MOTNAC codes are also converted to ICDO-M codes. All cases in the OCR are reported in ICD9 and ICDO-M.

Creation of Cancer Incidence Data

The methods of generating the cancer incidence data reported in this monograph are essentially the same over the twenty year time period and result from a collaboration between two departments of the Ontario Cancer Treatment and Research Foundation, the Division of Epidemiology and Statistics and Information Systems.

Once the source files have been preprocessed, cancer incidence data are created in two major phases. First, sequential computer linkage of the source files is carried out to bring all records pertaining to an individual together. Then, a set of computerized rules known as the "Case Resolution System" is applied to the linked records, to allocate the appropriate site of disease, histology, date and method of diagnosis, residence, and other information for each case of cancer.

Linkage of Data Sources

As described earlier, all records from 1964-1966 were linked manually. These linked records, together with the source data for 1967-1971, were then keypunched and stored in machine-readable form. The manual linkage of source records was replaced by computer programmes written by OCR staff to identify records having a high probability of belonging to the same individual, using record linkage principles developed by Newcombe, Kennedy and Smith[13,14,15]. The entire output from this record linkage of 1964-1971 data was manually edited to match records that were not linked by the computer and to separate those records that were linked incorrectly. A master record was created for each case and then inspected to determine whether the site, histology and date of diagnosis had been correctly assigned; these were then modified, if indicated. This simple linkage system and detailed manual editing were very labour intensive, again resulting in a delay in the production of incidence data. Cancer incidence data for 1969 to 1971 were not published until 1978[16].

Fortunately, by 1978 more efficient computers and sophisticated computerized record linkage systems were available which facilitated the task of linking large numbers of records. In order to link together the large volume of data, OCR staff have developed a sophisticated computer record linkage system based on the Generalized Iterative Record Linkage System (GIRLS)[17] designed by Statistics Canada in conjunction with the Epidemiology Unit of the National Cancer Institute of Canada.

Since Ontario does not have a unique number in the health or political system which identifies an individual throughout life, linkage is based on a number of identifying variables including name, date of birth, OHIP number, hospital where diagnosed and hospital chart number. It should be noted that an OHIP number is allocated to a family, and does not distinguish between individual members of that family. When a child reaches the age of 18 (or 21, if attending university), he or she is assigned a new OHIP number. Change of employment, or divorce, may also result in the allocation of new OHIP numbers to individuals.

The present computer linkage is completed in several stages. First, a NYSIIS (New York State Intelligence Identification System) code is created which is a phonetic version of the surname. Only records which have the same NYSIIS code are compared for possible linkage; therefore, records with names having similar spellings but different NYSIIS codes do not have an opportunity to link. Records with the same NYSIIS code constitute a "pocket" within which records are compared. A numeric score or "weight" is assigned to each variable when two records are compared. The greater the sum of the weights of the variables compared, the greater the probability that two records linked by the system belong to the same individual. The word "iterative" in the acronym GIRLS indicates that this process of allocating weights is repeated more than once. The system uses previous observations to assign more precise weights.

Each link (i.e., each pair of records brought together) is classified into one of three categories: definite, possible or rejected, based on the magnitude of the total weight. The distribution of the total weights in the linked file is usually bimodal, clustering around a high weight (definite, i.e., likely to be true links) and a low weight (rejected, i.e., unlikely to be true links). The middle range of weights contains possible links, i.e., those in which it is uncertain that paired records relate to the same individual. Linked records in this range (the grey area) are reviewed by health record staff of the OCR who have access to additional data that were not used in the linkage. An example would be information contained in the complete pathology report which might confirm the suspicion by the staff that an earlier biopsy had

been performed. Decisions are made to accept or to reject each link in the grey area and the result is then entered into the linked files. This manual resolution reduces the number of false links accepted and missed true links, but both still occur. The size of the grey area varies according to the files being linked; between 2 and 12% of potential links are manually resolved.

Linkages of source files are performed in sequence (Figure 1). Each year, hospital reports are linked internally to bring together multiple admissions for the same patient. Pathology reports are then linked to these aggregated hospital records, since most pathology reports will be related to a hospital stay. This combined hospital-pathology file is subsequently linked with previous years' incidence to identify incident (as opposed to prevalent) cases, producing provisional incidence data. Every second or third year, deaths due to cancer and records from the RCC and the PMH are linked to these provisional data. These final linkages add few new cases of cancer, although RCC and PMH records improve data quality, particularly the specificity of site and histology. This sequence of linkages has been applied to data from 1972 to the present. All RCC, PMH and death records from 1964 to 1983 have been included in the data presented in this monograph.



Fig. 1 Sequence of linkages of source files for generation of cancer incidence data.

An internal linkage has been performed on the entire 1964-1983 file using pockets other than those created by NYSIIS codes. This linkage allowed groups of records with different NYSIIS codes to be compared so that records which may have pertained to the same patient had the opportunity to link. There were three distinct phases to this linkage. Within pockets created in each phase, comparisons of all possible pairs of records were carried out and weights were assigned, as in other linkages. In the first phase, pockets were assigned using OHIP number and sex. In the second phase, pockets were formed using birth year and the first three letters of the given name. The third phase utilized the first three characters of the surname. Records linking at a high weight in one phase were not included in subsequent phases. The grey area resulting from this three-phase linkage was resolved as in other routine linkages. As a result of this linkage, 2.5% of patients were identified as being recorded twice in the OCR and records for these patients were amalgamated. This reduced the effect on the OCR due to errors in spelling or in transcription of surnames.

Nearly all cancer patients have multiple source records due to this method of cancer registration. A set of computer programmes has been developed by OCR staff to create a composite identification record containing the "best" identifying information as determined from all source records on a patient (e.g., surname, given names, birthdate, sex). This is then carried forward into the next linkage. These programmes also find conflicts between individual source records which may be the result of false links which had not been identified earlier. These conflicts are reported and reviewed by OCR health record staff, who make corrections as indicated.

Results of the early manual linkages have been set aside, original 1964-66 source records have been computerized and the same linkage techniques and rules have been applied over the twenty years of data reported in this monograph. As a result, data reported previously for earlier years[5,6,7] are not the same as those reported here.

Allocation of Site, Histology and Other Information

Groups of linked source records for individual patients are processed by a second major system, "Case Resolution". This consists of a series of computer modules developed by OCR staff, and applies medical logic to the source records for a patient to determine the appropriate site of disease, histology and date of diagnosis, since these may vary between source records.

In the Case Resolution system, cancer sites on all source records belonging to one individual are examined to determine the most specific site in a rubric or group of rubrics of the ICD. Only the most specific site codes are retained for further processing. Thus, if one source record indicated "malignant neoplasm of digestive tract" (ICD9 159.9), another indicated "malignant neoplasm of stomach not otherwise specified" (ICD9 151.9), another "malignant neoplasm of pylorus" (ICD9 151.1) and another "malignant neoplasm of plyoric antrum" (ICD9 151.2), only codes ICD9

151.1 and ICD9 151.2 would be retained because they are the most specific.

If at this stage only one site code remains, it is deemed to represent the primary site. If more than one remains but the only difference occurs in the 4th digit of the ICD9 (e.g., 151.1 and 151.2), then the site is selected from the most reliable source. For this purpose, RCC and PMH records are considered to be the most reliable source, followed by pathology records, then hospital discharge records and, finally, death certificates.

If more than one 3-digit site code remains, histology codes on each record for a patient are compared. Histology codes considered to be the same are organized into groups, according to a modification of the classification prepared by Berg[18] and are presented in Table B. Records with a blank histology field, or in which the histology is either "neoplasms not otherwise specified" (ICDO-M 800) or "no microscopic confirmation of tumour" (ICDO-M 999), are included in all histology groups. In addition, records in which the histology given is "carcinoma not otherwise specified" (ICDO-M 801) and "carcinoma undifferentiated type not otherwise specified" (ICDO-M 802) are included with all histology groups except sarcoma, lymphoma and leukaemia. The ICDO-M codes not given in the table are considered to each have a different histology from any other, for example, "mucoepidermoid neoplasms" (ICDO-M code 843).

The OCR considers a second site of cancer in the same individual to be metastatic unless clearly shown to be

TABLE B

Groupings of histology codes considered the "same" for allocation of site in the OCR

Name ICDO-M	Code ICDO-M
Squamous cell carcinomas	807-808
Transitional cell carcinomas	812-813
Adenocarcinomas	814,816,818-823, 825-838,857
Adnexal carcinomas	839-842
Cystic, mucinous and serous carcinomas	844-848
Ductal carcinomas	850-854
Specialized gonadal carcinomas	859,860,862-867
Paragangliomas and glomus tumours	868-871
Melanomas	872-874,876-878
Sarcomas and other soft tissue tumours	880,881,883-886, 889-892,899
Teratomatous carcinomas	908,909
Blood vessel and lymphatic vessel carcinomas	912-915,917
Osteosarcomas, chondrosarcomas and odontogenic tumours	918,919,922-927,929-933
Other tumours (pinealoma, chordoma and granular cell myoblastoma)	936,937
Gliomas	938-948
Neuroepitheliomatous tumours	949,950
Nerve sheath tumours	954,956
Lymphomas and Hodgkin's disease	959-966,969-972,975
Leukaemias	980-994

otherwise. Thus the rules for reporting second primary cancers are conservative. For two different primary sites to be reported in the same individual, the sites must be different at the 3-digit ICD9 level and the histologies of the two sites must be in different groups, as given in Table B. The only exception to this rule is breast cancer. Other sites rarely metastasize to the breast; if breast cancer is given as the site on a source record, then it will always be reported as a primary site, even if the histology is in the same group as that of other primary sites in the linked records. The case resolved from the other primary sites is also reported.

If different 3-digit site records have histologies in the same group according to Table B, one or more sites are considered to be metastatic and the site reported by the most reliable source, as defined earlier, will be allocated as the primary site. If the sources are equally reliable, a broad code which encompasses all the more specific site codes from the most reliable sources is selected as representing the primary site. For example, adenocarcinomas of the transverse colon (ICD9 153.1) and of the rectum (ICD9 154.1), reported by equally reliable sources, would be allocated to "large intestine, not otherwise specified" (ICD9 153.9).

Although the OCR does not report non-melanotic skin cancer (ICD9 173), records of this site are used, if appropriate, to override the site and histology given by less reliable sources. Thus, a pathology record with a diagnosis of cancer of skin of lip (ICD9 173.0) will result in a non-reported case, even although the hospital record indicated a diagnosis of cancer of the lip (ICD9 140.9) as the site of cancer.

The primary site is resolved to "malignant neoplasm without specification" (ICD9 199) when more specific allocation cannot be achieved. This can happen in three ways: first, if the only site recorded is 199; second, if there are two possible primaries in different organ groups with the same histology and equally reliable sources; and third, if more than one secondary site (ICD9 196-198) is specified in the absence of a primary site.

If more than one primary site is identified, each source record is examined and associated with the most appropriate of these diagnoses, thereby aggregating all data for each site. Finally, a composite case record for each case diagnosed is created comprising the best set of diagnostic information, as determined by the computer, using all source records. Checks are made to ensure consistency of composite information, (e.g., that site and histology are not in conflict, that date of death does not precede date last known alive).

The Case Resolution system is being continually improved. Cases which cannot be resolved by the rules, or which include inconsistencies, are reviewed by OCR staff and may result in subsequent modification of the computer rules. Rule changes can be encoded into the system and the entire registry file reprocessed according to the new rules.

Variables in the Ontario Cancer Registry

Three kinds of records exist in the registry: source records, composite identification records and composite case records. Source records are obtained from the external sources previously described, and represent cancer-related events. Composite identification records are created by record linkage, and represent the best identifying and demographic information on each cancer patient. Finally, composite case records are created by the Case Resolution system, and each composite case record describes an individual primary case of cancer. Most analyses use the variables in the composite records as listed in Table C.

Advantages and Limitations of the Ontario Cancer Registry

The advantages of the unique system of registration in Ontario are several. The multiple sources of data combine to provide incidence data of good quality and completeness. A recent study of completeness of cancer registration in 1982, using capture-recapture methodology, estimated completeness for all sites combined as more than 95%, with a low of 91% for cutaneous malignant melanoma to a high of over 98% for deep-seated digestive organs[19]. When data from several sources are present for a given diagnosis, the OCR system takes advantage of the known strengths of each particular source to select the best information for each variable.

The multiple sources generate a "patient profile", in that data from all hospital discharges related to the cancer diagnosis are stored in the OCR, so that length of stay and other data which are valuable to health planners are available. The patient profile also readily permits identification of multiple cancers in the same patient.

TABLE C

Variables Retained in the Composite OCR Record

Composite Identification Record

- registry identification number.
- names (including alternates).
- sex, date and place of birth.
- hospital and residence codes.
- last known date.
- vital status as of last known date.
- cause of death from the death certificate, if deceased.

Composite Case Record

- cancer site and histology codes.
- date of diagnosis.
- method of confirmation.
- residence at time of diagnosis.
- earliest known treatment date.
- hospital and RCC/PMH chart numbers.
- hospital of diagnosis.

This unique method of registration is relatively inexpensive, an important feature in a jurisdiction the size of Ontario.

Finally, the use of computerized linkage and Case Resolution systems ensures that records are processed in a consistent fashion. If the rules for allocating primary site are enhanced, the quality of the incidence data for the entire period of the OCR can be improved, as the complete registry data base (more than 6 million records on more than 500,000 cases) can be processed by the improved system. In addition, the impact of different rules for multiple primaries on incidence rates could be assessed by processing the entire registry file through two separate case resolution programs and comparing the results.

There are, however, some limitations of the system as well. These are primarily related to reliance on, and therefore limited control over, the type, quality and flow of input data. For some data sources, coding of site, histology and residence is decentralized to hospital medical record departments (hospital separation reports) and the Office of the Registrar General (death certificates). Thus, the OCR has no control over the quality of data from these sources, both in coding and in the number of records received. Nevertheless, approximately 50% of cancer cases in the province are eventually referred to the RCC or the PMH, where data quality and uniformity of coding are ensured.

Changes in any of the input data sources can affect registration, so that time trends in cancer incidence may be subject to artifacts related to changes in or problems with sources. For example, in 1978, changes in the administrative arrangements concerning provision of hospital discharge data to the MOH resulted in increased numbers of hospital discharge reports being received by the OCR, thereby producing a sudden increase in incidence rates. Such effects are likely to become less frequent as the OCR matures. Another problem with dependence on outside data sources is the resultant delay in generation of incidence data. HMRI processes all hospital discharge data for a fiscal year at the end of the year; only afterwards are data passed to the OCR. Since hospital discharges comprise the major registry source, linkages cannot begin until these data are received. Nine months of processing are required at the OCR to add one year of data. Therefore incidence data for a particular year are not available until 18-24 months after the close of that year.

Throughout the 1970's the annual number of hospital pathology reports voluntarily submitted to the OCR increased dramatically. Thus, during these years, the OCR would be expected to include increasing numbers of patients reported solely by pathology departments and never admitted to hospital or referred to the PMH or the RCC. For some sites, particularly those in which admission to hospital as an in-patient is not common, artifactual increases may thus be evident throughout this period.

Further problems, occurring from the use of data generated for purposes other than cancer registration, are lack of complete demographic/geographic/tumour-specific information. For example, data are not available on clinical stage at diagnosis except for a proportion of patients seen at the RCC or PMH. Municipality of residence is not available historically for many patients or, currently, for those for whom pathology reports only are received. Age and exact date of birth, an important linkage variable, are sometimes missing, particularly from pathology reports. This contributes to the relatively high proportion (0.06%) of cases reported with unknown age.

In addition, it is likely that there is some over-reporting in the OCR. Although a large proportion of duplicate records (where records for the same individual have not been brought together by the linkage system) were eliminated by the internal linkage process using pockets other than NYSIIS, it is estimated that 0.2% duplicates remain. This is an insoluble problem in a province where individuals do not have unique identifiers. However, the magnitude of the problem of over-reporting, due to failure to correctly link all records, is much less than it would have been if a completely manual linkage were performed. Comparison of data using the present system, published herein, with results of the original 1965-1966 manually generated data[6,7] demonstrates an 11% reduction in the number of cases. Similarly, there is a 5% reduction in the number of cases for 1969-1971 when incidence was generated using simplified computer linkage techniques[16]. When hospital records were coded at the OCR, control of quality and consistency of the coding could be conducted. Currently, as these source records are submitted to the OCR in machine-readable form, control is not possible. This has led to some over-reporting in that hospitals do, on occasion, report cancer cases, as cancer was the admitting diagnosis, whereas the tumours are found upon further investigation in hospital to be non-malignant. The OCR is constantly reviewing sites and age groups where this may occur and taking corrective measures.

Conclusion

Although data for 1964-1982 were generated in large segments, (those for 1972-1976 becoming available in September 1983 and those for 1977-1982 in late 1984), incidence rates since that time have been generated annually. Currently, incidence data are not available for a particular year until 18-24 months after the close of that year. It is expected that this interval will shorten with increasing computerization of hospital discharge and pathology reports at source.

Additional sources, such as cytopathology and haematology reports, may be added to the routine data sources; private pathology laboratories began submitting reports in 1984. This should improve both completeness and data quality. In addition, the OCR is continually investigating new sources of data which could be used to improve the quality of the OCR. For example, a linkage with a government data source is in progress now to improve residence information.

The linkage and Case Resolution systems are constantly being improved and streamlined, and the quality and timeliness of OCR incidence data will improve as they do.

Although the OCR contains twenty years of data, it is a new registry. Only through critical use of the data and comments from researchers using the database will the OCR reach its potential. Already, numerous research projects using the

OCR have resulted in improvement. The innovative method of cancer registration used in Ontario has resulted in a cancer registry of good quality, particularly in later years where the proportion of cases identified by death certificates is low and the proportion histologically verified is more than 85%. As more computers are introduced into different aspects of the health care system, this unique approach may prove to be the optimal technique for cancer registration in the future.

Cancer Incidence in Ontario

Indices of Quality

The three indices of quality traditionally used by cancer registries are percentage histologically verified (HV%), percentage death certificate only (DCO%) and mortality to incidence ratio expressed as a percentage (M/I%). Since the more precise definition of these indices varies from registry to registry, a description of the way they are defined in the OCR follows.

Histologically verified cases are those where at least one source record for a case indicates microscopic confirmation. The years 1967-1968 and 1972-1977 must be regarded as transition years for the OCR, when development of the unique system of registration did not permit allocation of resources to monitor the quality of source records. For these transition years the proportion histologically verified is low and the data should be interpreted accordingly.

A death certificate only case is one in which the only supporting source record for that case is a death certificate. No routine follow-up of DCO cases is conducted; thus, the proportion of cases not ascertained through one of the other sources is truly represented by the percentage DCO.

The mortality/incidence ratio is the number of deaths in a given time period divided by the number of incident cases for that time period expressed as a percentage. If the number of deaths in a period exceeds the number of incident cases, cancer registration is considered incomplete. However, the Mortality/Incidence ratio expressed as a percentage may be in excess of 100 for reasons other than under-reporting. Thus, for cancers coded to other and ill-defined sites within a system such as "malignant neoplasm of other and ill defined sites within the oral cavity and pharynx" (ICD9 149), the OCR could have more specific information than was given on the death certificate. This results in a reduced number of incident cases, compared to deceased cases, in that specific rubric and, hence, in an artifactually high index.

Interpretation of Data

As in all registries, rates are high for 1964, the initial year of cancer registration, which inevitably includes prevalent cases. In addition, there are a few aspects of the OCR which are unique and result in caveats which should be kept in mind when interpreting the data.

The OCR contains a high proportion of relatively non-specific sites, such as "other and ill defined sites within the digestive organs" (ICD9 code 159). This results from the rules used for allocating primary site in Case Resolution. This will, of necessity, result in somewhat reduced rates for specific sites within the same organ system.

The OCR contains a high proportion of cancers of "unspecified primary site" (ICD9 199), again resulting from the rules applied in Case Resolution where this rubric is used as a default if the system cannot identify the primary site.

Reliance on data from other sources results in artifacts which must also be remembered in interpreting the data. As mentioned earlier, there was a reduction in the number of hospital reports received in 1975, which is reflected in a fall in incidence rates for that year. In 1978 a marked increase in the number of hospital reports received resulted in a sudden increase in incidence rates. The increasing number of pathology reports received during the 1970's might contribute to part of the increasing incidence rates in those sites treated predominantly as out-patients, such as malignant melanoma (ICD9 172). The years 1967-1968 and 1972-1977 must be regarded as transition years when resources were not available to monitor the quality of source records.

Over-reporting in the OCR has been found in two specific areas, namely childhood cancer and cancer of the cervix. Very high rates were observed for childhood cancers with the inclusion of some unusual sites and a low proportion of histologic verification, suggestive of over-reporting. Since 1975, pathology reports have been requested for all cases of childhood cancer reported by hospital records only. Approximately 50% were found not to be invasive cancer, thereby reducing rates substantially. Thus, rates of childhood cancer prior to 1975 are believed to be inflated.

Similarly, apparently high rates in 1971 of cervical cancer led to a special study[20]. Only 25% of cases registered by hospital source records alone were confirmed, on receipt of further information from the hospital, as being invasive cancer. Difficulties by staff of medical records departments in distinguishing other conditions of the cervix (e.g., carcinoma in situ, chronic cervicitis) from invasive cancer led to the over-reporting. Additional information has been routinely requested on all cases of cancer of the cervix, originally registered by hospital source records only since 1969, and has been included in the data reported here. Rates of cervical cancer prior to 1969 should be interpreted with care.

It is unfortunate that it has not been possible to secure additional information for the early years of the OCR when over-reporting in such specific areas has been identified, but with the passage of time, hospital records become more difficult, if not impossible, to retrieve. The above examples of over-reporting confirm the salient comment by Waterhouse et al.[21] that "Cancer registration is not a matter of setting up a registry and then leaving it to run. Rather, it involves constant vigilance and monitoring at all levels so as to identify changes and take appropriate action as quickly as possible."

Explanatory Notes Regarding the Tables

Table 1 shows a list of 3-digit ICD9 site titles. ICD9 codes and abbreviated titles are used in the tables. The full descriptive title is taken from the "International Classification of Diseases, Ninth Revision"[10].

Table 2 presents the two standard populations used for the

calculation of age-adjusted rates. The first part of Table 2 includes the sex and age-specific population of Canada in 1971[22]. These age groupings were chosen to provide comparable rates to those already published for other jurisdictions in Canada[16]. The second part of Table 2 gives the "world" standard population by age group. The same world standard is used for both males and females, and yields rates which are comparable to those published in Cancer Incidence in Five Continents [21]

Table 3 displays the population of Ontario by calendar year and age group, for males and females separately. These populations may be used, together with cancer frequencies, to calculate age-specific or age-adjusted rates.

Table 4 provides a summary of the occurrence of cancer in Ontario. It includes the frequency of incident cancers (N), the percentage distribution of cancer by site of disease (% dist), and the incidence rate age-adjusted to the world standard population (AAR World) by site and sex, averaged over 5-year calendar periods.

Table 5 provides a summary of indices of the quality of data by site and sex, averaged over 5-year calendar periods.

Detailed site-specific tables follow and are arranged in numerical order by 3-digit ICD9 rubric, with the exception that sites 195-199 are placed at the end of the tables. For each site and sex, there are two tables, the first containing incidence frequencies and the second age-specific and summary rates. Methods used to calculate the statistics presented are detailed below.

Calculation of Rates

An age-specific rate (ASR) is calculated as the frequency in a given age and sex subgroup divided by the population for that same subgroup, and expressed per 100,000 population.

$$ASR_i = (F_i / P_i) \times 100{,}000$$

F_i: Number of new cancers occurring in the i^{th} age group.

P_i: Population of the i^{th} age group in the Ontario population.

Example:

The ASR_{0-4} of cancer incidence (ICD9 140-208, excl. 173) is calculated for Ontario males under 5 years of age in 1964:

$F_{0-4} = 90$

$P_{0-4} = 391{,}100$

$ASR_{0-4} = (90/391{,}100) \times 100{,}000 = 23.0$

Age adjusted rates (AAR) are standardized to the Canadian population (AAR Can) or to the world population (AAR World). AAR Can uses as its standard the population of Canada in 1971. AAR World uses an arbitrary population (called the "world" population) as its standard for age adjustment. Both types of age-adjusted rates were calculated using the entire standard population. The interested reader may calculate any particular truncated age-adjusted rates that are desired (truncation uses only a portion of the standard population to calculate rates). AAR is expressed per 100,000 population and is calculated using the formula:

$$AAR = \left\{ \sum_{i=1}^{n} (F_i \times P_{i,std} \times 100{,}000)/P_i \right\} / \left(\sum_{i=1}^{n} P_{i,std} \right).$$

n: Number of age subgroups included in AAR.

F_i: Number of new cancers occurring in the i^{th} age group.

P_i: Population of i^{th} age group in the Ontario population.

$P_{i,std}$: Population of i^{th} age group in standard population.

As an example, the $AAR_{all\ ages}$ for all cancer sites (ICD9 140-208, excl. 173) is calculated for Ontario males in 1964 adjusted to the world standard (Table D).

Cumulative Risk per cent for ages 0-74 is an estimate of an individual's risk of developing cancer of a particular type, up to the age of 74 years:

$$\text{CUM RISK \%}_{0-74} = \left\{1-e^{-y}\right\} \times 100, \text{ where } y = \sum_{i=1}^{n} (F_i \times T_i / P_i)$$

n: Number of age groups which CUM RISK includes.

F_i: Number of new cancers occurring in the i^{th} age group.

P_i: Population of i^{th} age group in the Ontario population.

T_i: Number of years in i^{th} age group.

As an example, the CUM RISK $\%_{0-74}$ for all types of cancer, excluding skin, is calculated for Ontario males in 1964 (Table E).

Table D

Sample Calculation of an AAR

i	Age Group	F_i	$P_{i,std}$	P_i	AAR
1	0-4	90	12000	391100	2.76
2	5-9	55	10000	371500	1.48
3	10-14	42	9000	331100	1.14
4	15-19	57	9000	273700	1.87
5	20-24	58	8000	212000	2.19
6	25-29	58	8000	204100	2.27
7	30-34	107	6000	225500	2.85
8	35-39	183	6000	236200	4.65
9	40-44	273	6000	217900	7.52
10	45-49	411	6000	188600	13.08
11	50-54	659	5000	170000	19.38
12	55-59	994	4000	139700	28.46
13	60-64	1300	4000	113900	45.65
14	65-69	1404	3000	90000	46.80
15	70-74	1382	2000	68300	40.47
16	75-79	1179	1000	45400	25.97
17	80-84	796	500	25400	15.67
18	85+	447	500	13600	16.43
TOTAL					278.64

$AAR_{\text{all ages}} = 278.6$

Table E

Sample Calculation of CUM RISK $\%_{0\text{-}74}$

i	Age Group	F_i	T_i	P_i	$(F_i x T_i / P_i)(x10^{-4})$
1	0-4	90	5	391100	11.51
2	5-9	55	5	371500	7.40
3	10-14	42	5	331100	6.34
4	15-19	57	5	273700	10.41
5	20-24	58	5	212000	13.68
6	25-29	58	5	204100	14.21
7	30-34	107	5	225500	23.73
8	35-39	183	5	236200	38.74
9	40-44	273	5	217900	62.64
10	45-49	411	5	188600	108.96
11	50-54	659	5	170000	193.82
12	55-59	994	5	139700	355.76
13	60-64	1300	5	113900	570.68
14	65-69	1404	5	90000	780.00
15	70-74	1382	5	68300	1011.71
TOTAL					3209.59

$CUM\ RISK_{0\text{-}74} = 1 - e^{-0.3210}$

$= 1 - 0.7255$

$= 0.2745$

$CUM\ RISK\ \%_{0\text{-}74} = 27.45\%$

Obtaining More Detailed Information From the Ontario Cancer Registry

The data in this monograph are available on magnetic tape and can be obtained on request. In addition to the material presented, the OCR routinely provides similar data for smaller regions in the province (e.g., county) and at the 4-digit ICD9 level. Such data can normally be provided in four to eight weeks from receipt of a request. Requests are generally free of charge and can be made in writing to:

DR. E.A. CLARKE, Director
Ontario Cancer Registry
The Ontario Cancer Treatment
and Research Foundation
7 Overlea Blvd.
Toronto, Ontario, Canada
M4H 1A8

Tel: (416) 423-4240
Fax: (416) 423-2017

References

1. 1986 Census of Canada: Final Population and Dwelling Counts. Ottawa: Statistics Canada; 1987.

2. 1981 Census of Canada: Population Mobility Status. Cat. No. 92-907, Vol. 1: National Series. Ottawa: Statistics Canada; 1983.

3. Sellers AH, Marr WB, Kelly AD: A survey of cancer in Middlesex County, Ontario. *Can J Public Health* 1950; **41**: 314.

4. MacKay EN, Sellers AH: A repeat survey of cancer in Middlesex County, Ontario. The Ontario Cancer Treatment and Research Foundation, 1953.

5. MacKay EN, Sellers AH: A study of cancer incidence and prevalence in Ontario, 1964. *Can Med Assoc J* 1968; **99**: 313.

6. MacKay EN, Sellers AH: The Ontario cancer incidence survey, 1965: A progress report. *Can Med Assoc J* 1970; **103**: 51.

7. MacKay EN, Sellers AH: The Ontario cancer incidence survey, 1964-1966: a new approach to cancer data acquisition. *Can Med Assoc J* 1973; **109**: 489.

8. International Classification of Diseases: Manual of the International Statistical Classification of Diseases, Injuries, and Causes of Death. 1955 Revision; vol. 1. Geneva: World Health Organization; 1957.

9. Eighth Revision: International Classification of Diseases Public Health Service Publication No. 1693. Washington DC: U.S. Department of Health, Education, and Welfare.

10. International Classification of Diseases: Manual of the International Statistical Classification of Diseases, Injuries, and Causes of Death. 1975 Revision, vol. 1. Geneva: World Health Organization; 1977.

11. American Cancer Society Inc.: Manual of Tumor Nomenclature and Coding, 1968.

12. World Health Organization Manual of the International Classification of Diseases for Oncology, 1st edition. Geneva: World Health Organization; 1976.

13. Newcombe HB, Kennedy JM, Axford SJ, et al: Automatic linkage of vital records. *Science* 1959; **130**: 954.

14. Kennedy JM, Newcombe HB, Okazaki EA, et al: Computer methods for family linkage of vital and health records. Chalk River: Atomic Energy of Canada Limited, 1965, Document AECL-2222.

15. Smith ME, Newcombe HB:Methods for computer linkage of hospital admission - separation records into cumulative health histories. *Methods Inf Med* 1975; **14**: 118.

16. New Primary Sites of Malignant Neoplasms in Canada. Cat. No. 82-207. Ottawa: Statistics Canada; 1978.

17. Howe GR, Lindsay J: A generalized iterative record linkage computer system for use in medical follow-up studies. *Comput Biomed Res* 1981; **14**: 327.

18. Berg JW: Morphological Classification of Human Cancer. *In*: Schottenfeld D and Fraumeni JF, eds. Cancer Epidemiology and Prevention. Philadelphia: WB Saunders, 1982: 74.

19. Robles SC, Marrett LD, Clarke EA, Risch HA: An application of capture-recapture methods to the estimation of completeness of cancer registration. 1987: *J Chron Dis* (in press).

20. Clarke EA, Hilditch S: Problems in determining the incidence of cervical cancer. *Can Med Assoc J* 1983; **129**: 1271.

21. Waterhouse, et al. (eds.): Cancer Incidence in Five Continents, Volume IV, World Health Organization, Lyon: 1982.

22. Cancer in Canada 1981. Cat. No. 82-207. Ottawa: Statistics Canada; 1986.

TABLE 1

ICD9 CODE, ABBREVIATED TITLE, AND FULL DESCRIPTIVE TITLE FOR INDIVIDUAL CANCER SITES

ICD9 CODE	ABBREVIATED TITLE	FULL DESCRIPTIVE TITLE
140-208,E173	ALL SITES	ALL MALIGNANT NEOPLASMS (EXCLUDING SKIN)
140	LIP	MALIGNANT NEOPLASM OF LIP
141	TONGUE	MALIGNANT NEOPLASM OF TONGUE
142	SALIVARY GLAND	MALIGNANT NEOPLASM OF MAJOR SALIVARY GLANDS
143	GUM	MALIGNANT NEOPLASM OF GUM
144	FLOOR OF MOUTH	MALIGNANT NEOPLASM OF FLOOR OF MOUTH
145	MOUTH UNSPECIFIED	MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF MOUTH
146	OROPHARYNX	MALIGNANT NEOPLASM OF OROPHARYNX
147	NASOPHARYNX	MALIGNANT NEOPLASM OF NASOPHARYNX
148	HYPOPHARYNX	MALIGNANT NEOPLASM OF HYPOPHARYNX
149	PHARYNX UNSPECIFIED	MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE LIP, ORAL CAVITY, AND PHARYNX
150	OESOPHAGUS	MALIGNANT NEOPLASM OF OESOPHAGUS
151	STOMACH	MALIGNANT NEOPLASM OF STOMACH
152	SMALL INTESTINE	MALIGNANT NEOPLASM OF SMALL INTESTINE, INCLUDING DUODENUM
153	COLON	MALIGNANT NEOPLASM OF COLON
154	RECTUM	MALIGNANT NEOPLASM OF RECTUM, RECTOSIGMOID JUNCTION AND ANUS
155	LIVER	MALIGNANT NEOPLASM OF LIVER AND INTRAHEPATIC BILE DUCTS
156	GALLBLADDER	MALIGNANT NEOPLASM OF GALLBLADDER AND EXTRAHEPATIC BILE DUCTS
157	PANCREAS	MALIGNANT NEOPLASM OF PANCREAS
158	PERITONEUM	MALIGNANT NEOPLASM OF RETROPERITONEUM AND PERITONEUM
159	ILL-DEFINED DIGESTIVE	MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE DIGESTIVE ORGANS AND PERITONEUM.
160	NOSE, SINUSES	MALIGNANT NEOPLASM OF NASAL CAVITIES, MIDDLE EAR AND ACCESSORY SINUSES
161	LARYNX	MALIGNANT NEOPLASM OF LARYNX
162	LUNG	MALIGNANT NEOPLASM OF TRACHEA, BRONCHUS AND LUNG
163	PLEURA	MALIGNANT NEOPLASM OF PLEURA
164	OTHER THORACIC ORGANS	MALIGNANT NEOPLASM OF THYMUS, HEART AND MEDIASTINUM
165	ILL-DEFINED RESPIRATORY	MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE RESPIRATORY SYSTEM AND INTRATHORACIC ORGANS
170	BONE	MALIGNANT NEOPLASM OF BONE AND ARTICULAR CARTILAGE
171	CONNECTIVE TISSUE	MALIGNANT NEOPLASM OF CONNECTIVE AND OTHER SOFT TISSUE
172	MELANOMA OF SKIN	MALIGNANT MELANOMA OF SKIN
174/5	BREAST	MALIGNANT NEOPLASM OF FEMALE/MALE BREAST
179	UTERUS UNSPECIFIED	MALIGNANT NEOPLASM OF UTERUS, PART UNSPECIFIED
180	CERVIX UTERI	MALIGNANT NEOPLASM OF CERVIX UTERI
181	PLACENTA	MALIGNANT NEOPLASM OF PLACENTA
182	BODY OF UTERUS	MALIGNANT NEOPLASM OF BODY OF UTERUS
183	OVARY	MALIGNANT NEOPLASM OF OVARY AND OTHER UTERINE ADNEXA
184	OTHER FEMALE GENITAL	MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED FEMALE GENITAL ORGANS
185	PROSTATE	MALIGNANT NEOPLASM OF PROSTATE
186	TESTIS	MALIGNANT NEOPLASM OF TESTIS
187	MALE GENITAL	MALIGNANT NEOPLASM OF PENIS AND OTHER MALE GENITAL ORGANS
188	BLADDER	MALIGNANT NEOPLASM OF BLADDER
189	KIDNEY, OTHER URINARY	MALIGNANT NEOPLASM OF KIDNEY AND OTHER UNSPECIFIED URINARY ORGANS
190	EYE	MALIGNANT NEOPLASM OF EYE
191	BRAIN	MALIGNANT NEOPLASM OF BRAIN
192	NERV. SYST. UNSPECIFIED	MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF NERVOUS SYSTEM
193	THYROID	MALIGNANT NEOPLASM OF THYROID GLAND
194	OTHER ENDOCRINE	MALIGNANT NEOPLASM OF OTHER ENDOCRINE GLANDS AND RELATED STRUCTURES
200	LYMPHOSARCOMA	LYMPHOSARCOMA AND RETICULOSARCOMA
201	HODGKIN'S DISEASE	HODGKIN'S DISEASE
202	OTHER LYMPHOMAS	OTHER MALIGNANT NEOPLASM OF LYMPHOID AND HISTIOCYTIC TISSUE
203	MULTIPLE MYELOMA	MULTIPLE MYELOMA AND IMMUNOPROLIFERATIVE NEOPLASMS
204	LYMPHOID LEUKAEMIA	LYMPHOID LEUKAEMIA
205	MYELOID LEUKAEMIA	MYELOID LEUKAEMIA
206	MONOCYTIC LEUKAEMIA	MONOCYTIC LEUKAEMIA
207	OTHER LEUKAEMIA	OTHER SPECIFIED LEUKAEMIA
208	LEUKAEMIA, CELL UNSP.	LEUKAEMIA OF UNSPECIFIED CELL TYPE
195	ILL-DEFINED SITES	MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES
196	LYMPH NODES SECONDARY	SECONDARY AND UNSPECIFIED NEOPLASM OF LYMPH NODES
197	RESP/DIGEST SECONDARY	SECONDARY MALIGNANT NEOPLASM OF RESPIRATORY AND DIGESTIVE SYSTEMS
198	OTHER SECONDARY	SECONDARY MALIGNANT NEOPLASM OF OTHER SPECIFIED SITES
199	UNSPECIFIED SITE	MALIGNANT NEOPLASM WITHOUT SPECIFICATION OF SITE

TABLE 2

STANDARD POPULATIONS

STANDARD POPULATION, CANADA 1971

	AGE (YEARS)								
SEX	0-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	ALL AGES
FEMALE	5,104,960	1,427,985	1,240,575	1,159,260	877,635	575,730	304,000	82,805	10,772,950
MALE	5,279,700	1,461,590	1,285,830	1,132,315	854,085	501,625	225,675	54,535	10,795,355

STANDARD POPULATION, WORLD

AGE (YEARS)																	
0-4	5-9	10-14	15-19	20-24	25-29	30-34	34-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	ALL AGES
12000	10000	9000	9000	8000	8000	6000	6000	6300	6000	5000	4000	4000	3000	2000	1000	500	500	100000

SOURCES:

1) STATISTICS CANADA: CANCER IN CANADA 1981, CAT. NO. 82-207, OTTAWA: 1986: 175
2) WATERHOUSE, ET AL. (EDS.): CANCER INCIDENCE IN FIVE CONTINENTS, VOLUME IV, WORLD HEALTH ORGANIZATION, LYON: 1982: 673

TABLE 3

POPULATION OF ONTARIO: MALES (IN HUNDREDS)

YEAR	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	ALL AGES
1964	3911	3715	3311	2737	2120	2041	2255	2362	2179	1886	1700	1397	1139	900	683	454	254	136	33180
1965	3893	3823	3404	2901	2245	2073	2252	2376	2259	1909	1748	1432	1167	915	694	459	259	139	33948
1966	3825	3935	3519	3047	2407	2154	2266	2396	2330	1955	1790	1471	1196	931	704	466	260	140	34792
1967	3729	3988	3634	3188	2613	2265	2285	2404	2384	2014	1819	1520	1229	950	709	475	264	145	35615
1968	3600	4026	3742	3300	2803	2380	2294	2394	2408	2089	1828	1572	1259	972	711	478	268	150	36274
1964-68	18958	19487	17610	15173	12188	10913	11352	11932	11560	9853	8885	7392	5990	4668	3501	2332	1305	710	173809
1969	3454	4038	3850	3396	2977	2511	2304	2381	2414	2175	1831	1624	1289	1000	718	481	272	157	36872
1970	3353	4035	3954	3515	3177	2688	2354	2387	2417	2261	1849	1671	1324	1030	729	485	278	164	37671
1971	3263	4012	4037	3622	3348	2867	2420	2393	2420	2325	1888	1704	1361	1061	744	489	282	173	38409
1972	3239	3887	4090	3723	3347	3048	2507	2376	2432	2334	1987	1708	1410	1080	767	494	287	179	38895
1973	3235	3700	4148	3815	3384	3195	2604	2367	2436	2352	2089	1696	1455	1101	796	498	291	182	39344
1969-73	16544	19672	20079	18071	16233	14309	12189	11904	12119	11447	9644	8403	6839	5272	3754	2447	1410	855	191191
1974	3235	3549	4188	3925	3502	3350	2748	2402	2444	2364	2189	1686	1503	1129	823	506	292	185	40020
1975	3204	3459	4185	4008	3608	3475	2859	2437	2426	2380	2235	1720	1552	1160	844	524	292	190	40558
1976	3117	3420	4093	4127	3683	3564	2957	2473	2408	2390	2271	1790	1575	1205	864	544	297	194	40972
1977	3098	3379	3985	4154	3759	3526	3161	2543	2395	2404	2276	1888	1578	1259	883	565	302	197	41352
1978	3088	3359	3823	4187	3822	3528	3300	2652	2386	2418	2290	1990	1575	1301	907	589	308	200	41723
1974-78	15742	17166	20274	20401	18374	17443	15025	12507	12059	11956	11261	9074	7783	6054	4321	2728	1491	966	204625
1979	3078	3305	3650	4208	3861	3542	3398	2743	2407	2410	2302	2085	1567	1345	937	612	318	205	41973
1980	3053	3252	3533	4209	3890	3584	3493	2825	2434	2367	2329	2128	1612	1388	970	629	331	207	42254
1981	3037	3173	3467	4134	3927	3588	3571	2915	2472	2365	2355	2158	1688	1411	1007	647	341	212	42468
1982	3063	3126	3437	4024	4000	3656	3525	3129	2540	2354	2369	2182	1778	1427	1046	665	357	216	42894
1983	3087	3092	3409	3861	4112	3739	3522	3279	2641	2355	2371	2210	1877	1423	1091	685	378	223	43355
1979-83	15318	15948	17496	20436	19790	18109	17509	14891	12494	11671	11726	10763	8522	6994	5051	3238	1725	1063	212944

TABLE 3

POPULATION OF ONTARIO: FEMALES (IN HUNDREDS)

YEAR	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	ALL AGES
1964	3732	3549	3163	2661	2159	2071	2185	2340	2216	1851	1645	1380	1180	1016	830	586	346	220	33130
1965	3710	3655	3251	2812	2278	2107	2186	2327	2300	1893	1697	1420	1211	1036	856	604	359	230	33932
1966	3632	3766	3364	2945	2444	2185	2206	2330	2364	1955	1744	1462	1245	1061	886	619	372	237	34817
1967	3542	3811	3481	3098	2623	2276	2224	2311	2401	2029	1787	1514	1284	1085	901	647	390	251	35655
1968	3408	3835	3585	3223	2802	2374	2236	2290	2400	2121	1813	1571	1321	1113	912	670	407	265	36346
1964-68	18024	18616	16844	14739	12306	11013	11037	11598	11681	9849	8686	7347	6241	5311	4385	3126	1874	1203	173880
1969	3273	3838	3678	3314	2977	2489	2254	2277	2372	2220	1834	1628	1357	1146	926	688	424	283	36978
1970	3192	3840	3770	3421	3199	2645	2303	2285	2344	2311	1869	1686	1396	1181	947	705	442	303	37839
1971	3110	3823	3840	3512	3393	2807	2361	2292	2321	2373	1923	1732	1439	1217	971	721	462	325	38622
1972	3081	3693	3891	3614	3373	3047	2439	2292	2321	2370	2039	1747	1490	1252	991	743	475	343	39201
1973	3071	3524	3940	3701	3417	3192	2553	2297	2321	2369	2159	1750	1540	1286	1018	760	490	359	39747
1969-73	15727	18718	19119	17562	16359	14180	11910	11443	11679	11643	9824	8543	7222	6082	4853	3617	2293	1613	192387
1974	3071	3381	3983	3800	3544	3354	2685	2344	2337	2356	2277	1762	1595	1324	1049	779	501	379	40521
1975	3038	3295	3984	3864	3670	3488	2806	2383	2335	2346	2338	1814	1660	1352	1081	800	516	394	41164
1976	2955	3259	3895	3953	3761	3595	2908	2427	2338	2330	2379	1904	1688	1404	1109	827	533	414	41679
1977	2943	3206	3788	3987	3835	3589	3128	2504	2333	2329	2364	2022	1712	1456	1148	847	554	433	42178
1978	2938	3178	3632	4030	3866	3628	3288	2617	2331	2334	2357	2145	1723	1502	1187	874	572	449	42671
1974-78	14945	16319	19282	19634	18696	17654	14815	12275	11674	11695	11715	9647	8378	7038	5574	4127	2676	2069	208213
1979	2929	3127	3469	4053	3905	3670	3409	2716	2362	2333	2337	2251	1733	1549	1229	902	596	471	43041
1980	2909	3074	3354	4047	3925	3739	3535	2801	2399	2321	2337	2302	1788	1611	1262	933	618	491	43446
1981	2893	3000	3297	3955	3970	3760	3644	2897	2447	2315	2334	2331	1877	1644	1303	963	639	515	43784
1982	2922	2967	3257	3829	4017	3828	3623	3121	2522	2314	2338	2335	1983	1672	1344	1001	654	540	44267
1983	2948	2942	3225	3675	4089	3894	3649	3287	2628	2323	2336	2347	2103	1670	1396	1037	681	569	44799
1979-83	14601	15110	16602	19559	19906	18891	17860	14822	12358	11606	11682	11566	9484	8146	6534	4836	3188	2586	219337

TABLE 4

SUMMARY OF CANCER INCIDENCE

MALES

SITE	1964-68			1969-73			1974-78			1979-83		
	N	XDIST	AAR WORLD	N	XDIST	AAR WORLD	N	XDIST	AAR WORLD	N	XDIST	AAR WORLD
ALL SITES	44194	100.0	249.4	50952	100.0	259.8	63683	100.0	289.8	74787	100.0	301.4
LIP	1136	2.6	6.4	1138	2.2	5.8	1105	1.7	5.0	1117	1.5	4.5
TONGUE	288	0.7	1.6	300	0.6	1.5	390	0.6	1.8	474	0.6	2.0
SALIVARY GLANDS	155	0.4	0.9	135	0.3	0.7	187	0.3	0.9	223	0.3	0.9
GUM	70	0.2	0.4	86	0.2	0.4	90	0.1	0.4	105	0.1	0.4
FLOOR OF MOUTH	160	0.4	0.9	206	0.4	1.1	278	0.4	1.3	313	0.4	1.3
MOUTH UNSPECIFIED	193	0.4	1.1	193	0.4	1.0	252	0.4	1.2	275	0.4	1.1
OROPHARYNX	192	0.4	1.1	310	0.6	1.6	359	0.6	1.7	311	0.4	1.3
NASOPHARYNX	92	0.2	0.5	139	0.3	0.7	160	0.3	0.8	164	0.2	0.7
HYPOPHARYNX	141	0.3	0.8	179	0.4	0.9	249	0.4	1.2	298	0.4	1.2
PHARYNX UNSPECIFIED	172	0.4	1.0	78	0.2	0.4	186	0.2	0.5	128	0.2	0.5
OESOPHAGUS	785	1.8	4.4	826	1.6	4.2	945	1.5	4.3	971	1.3	3.9
STOMACH	3130	7.1	17.5	3034	6.0	15.3	2966	4.7	13.4	2888	3.9	11.5
SMALL INTESTINE	88	0.2	0.5	109	0.2	0.5	158	0.2	0.7	202	0.3	0.8
COLON	4048	9.2	22.6	4631	9.1	23.5	5917	9.3	26.7	7268	9.7	28.8
RECTUM	2563	5.8	14.4	2882	5.7	14.6	3203	5.0	14.5	3528	4.7	14.1
LIVER	347	0.8	2.0	441	0.9	2.3	470	0.7	2.2	524	0.7	2.2
GALLBLADDER	322	0.7	1.8	352	0.7	1.8	409	0.6	1.9	504	0.7	2.0
PANCREAS	1606	3.6	9.1	1969	3.9	10.0	2030	3.2	9.3	2155	2.9	8.6
PERITONEUM	56	0.1	0.3	44	0.1	0.2	62	0.1	0.3	67	0.1	0.3
ILL-DEFINED DIGESTIVE	250	0.6	1.4	268	0.5	1.4	423	0.7	1.9	543	0.7	2.2
NOSE, SINUSES	148	0.3	0.8	192	0.4	1.0	302	0.5	1.4	203	0.3	0.8
LARYNX	820	1.9	4.7	1039	2.0	5.4	1293	2.0	6.0	1579	2.1	6.5
LUNG	7834	17.7	45.3	10327	20.3	53.5	13418	21.1	61.5	15620	20.9	63.0
PLEURA	21	0.0	0.1	34	0.1	0.2	55	0.1	0.2	128	0.2	0.5
OTHER THORACIC ORGANS	52	0.1	0.3	70	0.1	0.4	89	0.1	0.4	106	0.1	0.5
ILL-DEFINED RESPIRATORY	50	0.1	0.3	54	0.1	0.3	77	0.1	0.4	81	0.1	0.3
BONE	198	0.4	1.1	257	0.5	1.3	391	0.6	1.8	312	0.4	1.4
CONNECTIVE TISSUE	367	0.8	2.1	476	0.9	2.5	628	1.0	3.0	613	0.8	2.6
MELANOMA OF SKIN	534	1.2	3.0	733	1.4	3.7	1244	2.0	5.7	1822	2.4	7.5
BREAST	107	0.2	0.6	127	0.2	0.7	169	0.3	0.8	146	0.2	0.6
UTERUS UNSPECIFIED	-	-	-	-	-	-	-	-	-	-	-	-
CERVIX UTERI	-	-	-	-	-	-	-	-	-	-	-	-
PLACENTA	-	-	-	-	-	-	-	-	-	-	-	-
BODY OF UTERUS	-	-	-	-	-	-	-	-	-	-	-	-
OVARY	-	-	-	-	-	-	-	-	-	-	-	-
OTHER FEMALE GENITAL	-	-	-	-	-	-	-	-	-	-	-	-
PROSTATE	6343	14.4	34.3	6692	13.1	33.0	8851	13.9	39.0	11832	14.8	42.4
TESTIS	448	1.0	2.6	556	1.1	2.9	713	1.1	3.3	833	1.1	3.5
OTHER MALE GENITAL	102	0.4	1.0	167	0.3	0.8	209	0.3	0.9	225	0.3	0.9
BLADDER	3088	7.0	17.4	3483	6.8	17.6	4549	7.1	20.5	5605	7.5	22.4
KIDNEY, OTHER URINARY	1057	2.4	6.0	1245	2.4	6.5	1456	2.3	6.8	1876	2.5	7.8
EYE	154	0.3	0.9	185	0.4	1.0	199	0.3	1.0	282	0.3	0.9
BRAIN	931	2.1	5.4	1089	2.1	5.7	1298	2.0	6.2	1620	2.2	7.0
NERV. SYST. UNSPECIFIED	98	0.2	0.6	96	0.2	0.5	193	0.3	0.9	155	0.2	0.7
THYROID	188	0.4	1.1	189	0.4	1.0	288	0.5	1.3	361	0.5	1.5
OTHER ENDOCRINE	62	0.1	0.4	67	0.1	0.4	109	0.2	0.5	153	0.2	0.8
LYMPHOSARCOMA	973	2.2	5.6	1146	2.2	5.9	1441	2.3	6.7	1483	2.0	6.1
HODGKIN'S DISEASE	613	1.4	3.6	665	1.3	3.4	656	1.0	3.0	808	1.1	3.5
OTHER LYMPHOMAS	321	0.7	1.8	396	0.8	2.0	594	0.9	2.7	1096	1.5	4.6
MULTIPLE MYELOMA	472	1.1	2.7	597	1.2	3.0	777	1.2	3.5	979	1.3	3.9
LYMPHOID LEUKAEMIA	642	1.5	3.6	787	1.5	4.2	989	1.6	4.9	1338	1.8	6.0
MYELOID LEUKAEMIA	362	0.8	2.0	574	1.1	2.9	777	1.2	3.6	1016	1.4	4.1
MONOCYTIC LEUKAEMIA	41	0.1	0.2	48	0.1	0.2	42	0.1	0.2	50	0.1	0.2
OTHER LEUKAEMIA	8	0.0	0.0	36	0.1	0.2	39	0.1	0.2	44	0.1	0.2
LEUKAEMIA UNSPECIFIED	778	1.8	4.4	513	1.0	2.7	429	0.7	2.0	303	0.4	1.2
ILL-DEFINED SITES	200	0.5	1.1	372	0.7	1.8	569	0.9	2.5	516	0.7	2.1
LYMPH NODES SECONDARY	68	0.2	0.4	99	0.2	0.5	125	0.2	0.6	175	0.2	0.7
RESP/DIGEST SECONDARY	156	0.4	0.9	235	0.5	1.2	278	0.4	1.2	429	0.6	1.7
OTHER SECONDARY	128	0.3	0.7	196	0.4	1.0	227	0.4	1.0	343	0.5	1.4
UNSPECIFIED SITE	956	2.2	5.4	898	1.8	4.6	1458	2.3	6.5	1477	2.0	5.8

SEE TEXT SECTION 'INTERPRETATION OF DATA'

TABLE 4

SUMMARY OF CANCER INCIDENCE

FEMALES

	1964-68			1969-73			1974-78			1979-83		ICD9
N	%DIST	AAR WORLD	N	%DIST	AAR WORLD	N	%DIST	AAR WORLD	N	%DIST	AAR WORLD	
44187	100.0	225.5	51221	100.0	228.4	63209	100.0	247.1	71953	100.0	246.4	140-208,E173
49	0.1	0.2	91	0.2	0.4	140	0.2	0.5	205	0.3	0.6	140
138	0.3	0.7	152	0.3	0.7	222	0.4	0.9	238	0.3	0.8	141
131	0.3	0.7	127	0.2	0.6	214	0.3	0.9	175	0.2	0.6	142
38	0.1	0.2	69	0.1	0.3	56	0.1	0.2	74	0.1	0.2	143
40	0.1	0.2	59	0.1	0.3	107	0.2	0.4	114	0.2	0.4	144
84	0.2	0.4	112	0.2	0.5	163	0.3	0.6	185	0.3	0.6	145
65	0.1	0.3	124	0.2	0.6	133	0.2	0.6	156	0.2	0.6	146
52	0.1	0.3	63	0.1	0.3	67	0.1	0.3	92	0.1	0.4	147
48	0.1	0.2	77	0.2	0.3	68	0.1	0.3	95	0.1	0.3	148
41	0.1	0.3	39	0.1	0.2	47	0.1	0.2	48	0.1	0.2	149
336	0.8	1.6	408	0.8	1.7	474	0.7	1.6	499	0.7	1.5	150
1832	4.1	8.3	1818	3.5	7.1	1845	2.9	6.2	1896	2.6	5.5	151
108	0.2	0.5	119	0.2	0.5	149	0.2	0.6	192	0.3	0.6	152
5010	11.3	23.9	5841	11.4	23.9	7091	11.2	25.1	8212	11.4	24.8	153
2034	4.6	10.0	2197	4.3	9.3	2548	4.0	9.1	2841	3.9	8.9	154
257	0.6	1.2	267	0.5	1.1	340	0.5	1.2	306	0.4	1.0	155
603	1.4	2.8	638	1.2	2.6	657	1.0	2.3	800	1.1	2.3	156
1194	2.7	5.5	1471	2.9	5.9	1620	2.6	5.6	1928	2.7	5.8	157
67	0.2	0.3	56	0.1	0.2	55	0.1	0.2	62	0.1	0.2	158
245	0.6	1.1	277	0.5	1.0	471	0.7	1.5	403	0.6	1.1	159
106	0.2	0.5	121	0.2	0.5	285	0.5	1.0	139	0.2	0.5	160
91	0.2	0.5	148	0.3	0.7	230	0.4	0.9	244	0.3	0.9	161
1344	3.0	6.9	2234	4.4	10.1	3635	5.8	14.5	5581	7.8	19.6	162
12	0.0	0.1	21	0.0	0.1	29	0.0	0.1	48	0.1	0.2	163
43	0.1	0.2	42	0.1	0.2	56	0.1	0.2	93	0.1	0.4	164
10	0.0	0.1	8	0.0	0.0	17	0.0	0.1	19	0.0	0.1	165
122	0.3	0.6	213	0.4	1.0	295	0.5	1.2	242	0.3	0.9	170
310	0.7	1.7	406	0.8	2.0	585	0.9	2.5	530	0.7	1.9	171
738	1.7	4.0	958	1.9	4.6	1492	2.4	6.2	2046	2.8	7.8	172
12024	27.2	62.7	14082	27.5	64.7	16870	26.7	68.9	18357	25.5	65.9	174/5
282	0.6	1.5	295	0.6	1.4	291	0.5	1.2	179	0.2	0.6	179
3987	9.0	21.7	5377	6.6	16.6	2956	4.7	12.6	2781	3.9	10.7	180
23	0.1	0.1	24	0.0	0.1	74	0.1	0.3	33	0.0	0.1	181
2463	5.6	13.1	3159	6.2	14.7	4432	7.0	18.2	4882	6.8	17.5	182
2289	5.2	12.2	2573	5.0	12.0	3044	4.8	12.5	3261	4.5	11.9	183
460	1.0	2.2	562	1.1	2.3	708	1.1	2.6	671	0.9	2.1	184
-	-	-	-	-	-	-	-	-	-	-	-	185
-	-	-	-	-	-	-	-	-	-	-	-	186
-	-	-	-	-	-	-	-	-	-	-	-	187
1148	2.6	5.3	1302	2.5	5.2	1681	2.7	5.9	1970	2.7	6.1	188
662	1.5	3.4	728	1.4	3.3	850	1.3	3.4	1095	1.5	3.9	189
127	0.3	0.7	142	0.3	0.7	171	0.3	0.8	210	0.3	0.8	190
610	1.4	3.4	757	1.5	3.8	1010	1.6	4.5	1257	1.7	5.0	191
88	0.2	0.5	113	0.2	0.6	158	0.2	0.7	167	0.2	0.6	192
499	1.1	2.7	620	1.2	3.0	805	1.3	3.4	1069	1.5	4.1	193
45	0.1	0.3	38	0.1	0.2	85	0.1	0.4	159	0.2	0.7	194
750	1.7	3.9	967	1.9	4.3	1315	2.1	5.1	1259	1.7	4.2	200
431	1.0	2.4	489	1.0	2.4	501	0.8	2.2	563	0.8	2.2	201
248	0.6	1.3	311	0.6	1.4	574	0.9	2.2	997	1.4	3.5	202
367	0.8	1.8	553	1.1	2.3	754	1.2	2.7	900	1.3	2.7	203
387	0.9	2.0	511	1.0	2.4	654	1.0	2.8	910	1.3	3.4	204
361	0.8	1.9	513	1.0	2.4	608	1.0	2.5	907	1.3	3.1	205
48	0.1	0.2	27	0.1	0.1	38	0.1	0.1	32	0.0	0.1	206
6	0.0	0.0	19	0.0	0.1	11	0.0	0.0	22	0.0	0.1	207
588	1.3	3.1	398	0.8	1.7	336	0.5	1.2	252	0.4	0.7	208
156	0.4	0.8	321	0.6	1.3	442	0.7	1.6	496	0.7	1.5	195
53	0.1	0.3	88	0.2	0.4	105	0.2	0.4	150	0.2	0.5	196
146	0.3	0.7	289	0.6	1.2	355	0.6	1.2	524	0.7	1.5	197
101	0.2	0.5	155	0.3	0.6	231	0.4	0.8	358	0.5	1.1	198
670	1.5	3.3	652	1.3	2.7	1061	1.7	3.8	1029	1.4	3.0	199

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

TABLE 5

INDICES OF DATA QUALITY

MALES

SITE	1964-68			1969-73			1974-78			1979-83		
	MVX	DCOX	M/IX	MVX	DCOX	M/IX	MVX	DCOX	M/IX	MVX	DCOX	M/IX
ALL SITES	79.0	8.5	58.6	82.2	4.7	60.0	74.6	2.9	54.5	85.8	1.3	54.0
LIP	98.2	0.4	2.8	97.9	0.0	4.4	86.3	0.0	4.3	96.3	0.2	3.1
TONGUE	95.5	1.0	64.9	96.3	0.0	59.7	92.6	1.5	55.1	96.4	0.2	55.3
SALIVARY GLANDS	94.8	1.9	27.7	91.9	2.2	46.7	83.4	0.5	28.9	97.8	0.9	27.4
GUM	97.1	0.0	0.0	96.5	2.3	26.7	93.3	0.0	17.8	98.1	1.0	18.1
FLOOR OF MOUTH	96.9	1.3	38.1	99.0	0.0	25.7	97.1	0.4	22.3	97.1	0.0	22.7
MOUTH UNSPECIFIED	91.2	3.1	67.9	97.4	1.0	60.1	91.3	1.2	52.8	97.1	0.4	48.4
OROPHARYNX	97.4	1.0	33.9	98.7	0.6	33.5	94.4	0.8	35.9	97.4	0.3	40.8
NASOPHARYNX	92.4	3.3	60.9	97.1	0.0	54.8	95.0	0.0	58.1	97.6	0.0	56.7
HYPOPHARYNX	97.9	0.7	29.1	95.5	1.1	35.4	97.2	0.4	37.3	97.7	0.0	30.5
PHARYNX UNSPECIFIED	93.0	4.1	66.3	91.0	5.1	167.9	89.6	2.8	134.9	95.3	1.6	151.6
OESOPHAGUS	80.9	8.5	85.9	84.3	4.6	89.8	78.2	3.3	93.1	90.1	2.1	106.7
STOMACH	64.5	14.5	78.7	74.7	7.2	82.6	70.9	3.7	79.4	87.9	2.1	76.4
SMALL INTESTINE	78.4	8.8	67.8	78.9	6.4	70.6	72.8	3.8	48.7	94.1	0.8	51.5
COLON	76.2	10.7	58.2	82.2	5.4	59.8	73.8	2.4	50.4	91.1	0.8	48.1
RECTUM	83.0	8.2	54.3	89.6	3.8	50.1	81.1	1.7	44.3	92.8	0.9	42.2
LIVER	53.8	28.8	108.1	59.8	19.8	88.9	46.8	10.8	83.2	67.2	4.6	94.3
GALLBLADDER	69.9	17.4	83.9	79.3	9.4	84.1	69.7	5.4	79.7	83.5	1.6	68.8
PANCREAS	57.8	18.4	91.8	65.4	9.6	91.6	46.9	6.7	90.1	57.3	3.3	96.7
PERITONEUM	66.1	26.8	132.1	77.3	11.4	195.5	64.5	6.5	143.5	68.7	1.5	119.4
ILL-DEFINED DIGESTIVE	52.8	23.2	98.8	51.9	19.4	123.1	51.5	8.5	112.5	73.1	5.7	144.8
NOSE, SINUSES	91.2	6.1	36.5	98.1	0.5	42.7	59.9	0.7	25.8	98.1	0.5	35.8
LARYNX	93.4	2.1	39.9	95.5	1.3	38.4	91.6	0.6	36.8	96.8	0.1	33.6
LUNG	78.1	9.3	81.2	79.6	5.8	82.4	73.5	4.1	79.6	81.8	2.1	83.8
PLEURA	81.8	19.8	66.7	76.5	20.6	55.9	74.5	3.6	52.7	95.3	1.6	43.8
OTHER THORACIC ORGANS	86.5	7.7	67.3	88.6	8.6	38.6	88.8	4.5	39.3	81.1	0.9	31.1
ILL-DEFINED RESPIRATORY	96.8	0.8	0.8	96.3	1.9	7.4	81.8	5.2	16.9	87.7	0.8	1.2
BONE	87.4	8.6	80.8	88.3	3.5	62.6	71.1	1.8	45.3	81.7	0.6	36.2
CONNECTIVE TISSUE	95.4	3.8	35.7	96.8	1.1	37.2	75.6	0.6	22.9	92.4	0.5	25.3
MELANOMA OF SKIN	94.6	3.7	36.7	98.1	0.5	32.9	85.3	0.3	28.1	97.2	0.1	24.8
BREAST	91.6	4.7	30.8	88.2	1.6	28.3	81.7	1.2	27.8	93.2	0.8	28.8
UTERUS UNSPECIFIED	-	-	-	-	-	-	-	-	-	-	-	-
CERVIX UTERI	-	-	-	-	-	-	-	-	-	-	-	-
PLACENTA	-	-	-	-	-	-	-	-	-	-	-	-
BODY OF UTERUS	-	-	-	-	-	-	-	-	-	-	-	-
OVARY	-	-	-	-	-	-	-	-	-	-	-	-
OTHER FEMALE GENITAL	-	-	-	-	-	-	-	-	-	-	-	-
PROSTATE	76.8	6.8	38.8	84.3	2.9	41.8	77.8	1.7	35.8	90.7	0.6	33.3
TESTIS	97.3	1.1	31.8	97.8	0.7	28.4	93.7	0.1	23.8	98.2	0.1	18.9
OTHER MALE GENITAL	88.5	4.4	31.9	91.8	0.6	28.7	80.4	1.8	24.4	90.7	0.9	20.8
BLADDER	90.1	3.3	35.9	94.8	1.5	33.9	86.5	0.9	27.1	96.5	0.2	24.5
KIDNEY, OTHER URINARY	85.1	7.2	61.1	88.3	4.3	56.8	78.3	2.6	53.3	84.4	0.9	49.1
EYE	94.8	1.9	27.9	94.6	2.2	27.6	69.8	0.5	22.1	91.6	1.5	25.7
BRAIN	82.8	12.1	75.6	85.6	9.6	79.2	75.8	6.3	70.6	80.8	2.2	70.4
NERV. SYST. UNSPECIFIED	87.8	7.1	39.8	85.4	6.3	50.8	60.6	2.6	26.9	73.5	1.9	16.8
THYROID	91.5	1.6	36.7	96.8	1.6	40.7	82.6	0.7	22.6	94.2	0.8	16.1
OTHER ENDOCRINE	77.4	9.7	87.1	82.1	4.5	46.3	54.1	8.3	39.4	64.1	0.7	52.9
LYMPHOSARCOMA	89.2	5.5	57.8	91.4	1.7	49.6	92.6	0.8	33.7	95.9	0.0	26.9
HODGKIN'S DISEASE	93.3	2.8	53.2	92.2	1.8	48.1	93.0	0.6	38.4	95.7	0.4	28.6
OTHER LYMPHOMAS	72.6	12.5	57.3	65.4	8.3	63.1	53.0	4.5	74.9	85.6	1.1	71.6
MULTIPLE MYELOMA	66.9	11.8	60.4	58.8	6.9	68.5	57.0	4.2	62.8	73.7	1.2	64.5
LYMPHOID LEUKAEMIA	84.4	5.3	37.4	69.6	4.2	61.1	65.7	1.7	47.7	69.2	1.8	39.8
MYELOID LEUKAEMIA	82.8	5.5	37.8	54.4	6.8	74.4	50.5	3.6	76.2	76.8	0.9	61.6
MONOCYTIC LEUKAEMIA	46.3	26.8	95.1	50.8	12.5	70.8	40.5	2.4	66.7	74.8	0.8	44.8
OTHER LEUKAEMIA	100.0	0.0	0.0	66.7	16.7	44.4	64.1	2.6	56.4	56.8	2.3	77.3
LEUKAEMIA UNSPECIFIED	55.5	16.5	105.8	25.5	9.9	82.7	17.8	8.2	94.2	44.6	5.3	142.9
ILL-DEFINED SITES	85.5	0.0	0.0	79.6	3.8	18.3	50.8	2.6	15.1	58.7	0.8	15.3
LYMPH NODES SECONDARY	94.1	0.8	36.8	94.9	1.0	36.4	92.8	0.8	14.4	94.3	0.0	0.8
RESP/DIGEST SECONDARY	61.5	9.8	44.9	66.8	13.2	94.5	60.4	10.8	93.9	64.3	0.0	0.8
OTHER SECONDARY	85.2	0.8	0.8	70.4	9.7	79.1	63.8	7.9	67.4	63.6	0.0	0.8
UNSPECIFIED SITE	71.2	10.7	53.9	68.2	6.8	48.2	49.1	5.7	45.2	57.5	5.8	93.8

SEE TEXT SECTION 'INTERPRETATION OF DATA'

TABLE 5

INDICES OF DATA QUALITY

FEMALES

1964-68			1969-73			1974-78			1979-83			ICD9
MVX	DCOX	M/IX	MVX	DCOX	M/IX	MVX	DCOX	M/IX	MVX	DCOX	M/IX	
84.4	7.0	49.2	86.6	3.8	49.1	77.8	2.7	45.7	87.4	1.6	47.4	140-208,E173
95.9	0.0	0.0	92.3	0.0	1.1	74.3	0.0	4.3	93.2	0.0	1.5	140
94.2	0.7	45.7	96.1	2.0	59.2	92.8	0.5	50.5	96.2	0.4	40.3	141
89.3	2.3	30.5	92.1	3.1	29.1	79.9	0.5	15.0	96.0	0.0	23.4	142
100.0	0.0	0.0	97.1	0.0	24.6	94.6	0.0	21.4	95.9	0.0	17.6	143
97.5	2.5	25.0	100.0	0.0	22.0	94.4	0.0	20.6	96.5	0.9	23.7	144
95.2	2.4	58.3	95.5	1.8	48.2	90.2	0.6	41.7	98.7	0.5	41.1	145
96.9	1.5	36.9	99.2	0.0	28.2	97.7	0.0	18.3	97.4	0.6	37.2	146
92.3	3.8	48.4	95.2	0.0	42.9	89.6	1.5	40.3	89.1	2.2	56.5	147
100.0	0.0	39.6	96.1	0.0	33.8	95.6	0.0	22.1	97.9	0.0	30.5	148
93.4	3.3	44.3	84.6	7.7	102.6	76.6	2.1	93.6	91.7	2.1	93.8	149
69.3	15.8	79.8	83.1	5.6	81.1	72.2	6.1	83.3	87.8	2.6	89.6	150
59.6	17.1	78.5	70.3	9.1	79.9	60.3	6.4	77.8	79.4	4.1	74.5	151
77.8	13.9	77.8	91.6	4.2	48.7	77.9	2.8	34.2	91.1	3.1	61.5	152
78.6	10.6	55.9	82.8	4.8	56.4	73.0	3.1	50.0	89.2	1.8	47.8	153
84.7	6.4	46.5	89.3	3.6	45.9	80.8	1.7	41.2	89.2	1.5	39.8	154
44.4	29.6	111.7	57.7	16.1	83.5	42.9	16.2	82.4	57.5	5.2	101.3	155
65.0	20.9	88.6	76.5	7.5	87.9	71.4	4.9	87.2	83.0	2.8	76.4	156
59.4	17.6	87.9	62.5	11.4	88.2	43.2	8.4	93.0	54.5	5.0	94.1	157
71.6	25.4	132.8	60.7	12.5	151.8	54.5	7.3	132.7	79.0	0.0	122.6	158
42.4	33.9	122.4	43.3	22.4	140.4	37.2	20.8	126.8	49.4	16.9	207.9	159
92.5	2.8	34.9	84.3	8.8	28.1	49.1	0.7	14.0	86.3	0.0	20.1	160
83.5	4.4	37.4	89.2	4.1	33.1	81.7	0.9	33.0	90.6	3.7	43.4	161
75.5	9.2	72.8	79.9	5.0	73.7	73.7	3.7	72.8	81.3	1.9	76.7	162
50.0	33.3	133.3	85.7	9.5	42.9	79.3	0.0	65.5	79.2	0.0	43.8	163
76.7	14.0	51.2	81.0	11.9	47.6	80.4	3.6	42.9	84.9	2.2	35.5	164
90.0	0.0	10.0	87.5	0.0	25.0	82.4	0.0	17.6	68.4	0.0	0.0	165
83.6	6.6	64.8	91.1	2.3	55.4	66.8	2.0	38.3	75.6	1.7	33.9	170
94.2	4.5	34.5	96.1	2.0	29.8	77.8	1.0	22.6	73.0	0.2	28.9	171
97.3	2.0	23.2	97.7	0.7	23.8	83.6	0.7	18.5	96.6	0.1	16.2	172
92.6	2.9	38.0	95.3	1.2	36.0	88.5	0.9	35.6	95.5	0.6	37.5	174/5
81.9	4.6	26.6	83.4	4.4	34.6	43.3	2.4	37.5	76.8	5.0	76.5	179
95.1	2.0	31.7	96.2	1.2	32.8	91.6	1.1	34.8	96.8	0.6	35.2	180
73.9	21.7	47.8	91.7	4.2	45.8	56.8	0.0	10.8	69.7	0.0	18.2	181
95.2	2.5	24.8	95.9	1.6	19.5	91.4	0.7	15.8	98.0	0.4	15.8	182
85.3	6.9	67.0	89.4	3.5	68.0	83.7	2.1	62.9	91.6	1.4	63.4	183
94.3	3.3	33.0	95.4	1.8	36.5	85.5	1.6	28.1	93.1	1.8	29.2	184
-	-	-	-	-	-	-	-	-	-	-	-	185
-	-	-	-	-	-	-	-	-	-	-	-	186
-	-	-	-	-	-	-	-	-	-	-	-	187
86.6	6.1	41.6	90.2	2.0	35.1	83.1	1.8	33.0	94.1	0.7	29.9	188
85.3	6.2	52.6	88.5	4.8	51.5	78.1	2.5	48.9	82.1	0.8	49.5	189
92.1	4.7	32.3	93.7	2.1	28.2	70.8	1.2	29.8	91.4	1.4	21.9	190
80.8	13.6	74.8	81.4	12.4	73.7	70.8	8.1	69.8	74.8	3.0	69.1	191
94.3	3.4	28.4	82.3	7.1	28.3	53.2	5.1	31.6	70.7	1.2	22.2	192
72.8	3.0	26.7	93.4	1.5	19.5	80.7	0.7	18.1	95.0	0.5	12.8	193
64.4	24.4	102.2	68.4	15.8	65.8	50.6	4.8	31.3	59.1	2.5	44.7	194
90.0	4.4	53.2	91.2	2.6	47.8	91.3	0.8	33.9	94.8	0.3	26.0	200
89.6	5.1	43.9	94.5	0.6	38.2	91.2	0.6	32.1	94.8	0.2	25.9	201
74.6	13.3	54.0	60.5	9.6	74.3	46.7	5.9	73.9	83.1	1.8	70.7	202
64.9	9.8	62.4	61.8	4.9	59.5	54.9	4.0	57.0	71.8	1.6	64.1	203
85.3	4.1	33.3	62.2	6.3	57.5	58.0	4.6	49.5	65.1	1.8	38.4	204
83.4	6.6	43.2	57.3	6.6	75.8	50.5	3.9	77.5	74.6	1.3	63.5	205
56.3	22.9	77.1	63.0	7.4	70.4	42.1	5.3	71.1	65.6	3.1	78.1	206
100.0	0.0	0.0	73.7	10.5	42.1	45.5	9.1	100.0	68.2	0.0	81.8	207
55.4	15.6	104.8	25.1	11.6	85.7	17.9	10.7	84.5	41.7	11.9	131.0	208
76.3	0.8	0.0	67.9	7.5	38.9	35.5	8.4	43.2	45.0	5.0	29.2	195
92.5	3.8	34.0	87.5	2.3	37.5	79.0	0.0	6.7	86.7	0.0	0.0	196
61.0	9.6	40.4	64.7	12.5	92.4	62.5	9.3	67.6	62.0	0.0	0.0	197
79.2	0.0	0.0	65.2	10.3	69.7	59.7	6.5	48.5	62.8	0.0	0.0	198
60.0	19.7	79.3	56.4	10.4	54.4	39.8	9.5	59.6	47.8	10.1	136.9	199

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

ALL MALIGNANT NEOPLASMS (EXCLUDING SKIN) (ICD9 140-208 EXCL. 173)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	90	55	42	57	58	58	107	183	273	411	659	994	1308	1404	1382	1179	796	447	1	9496
1965	85	42	43	54	50	85	107	155	241	392	640	886	1025	1232	1267	1084	765	420	1	8574
1966	92	62	34	46	51	62	85	168	289	382	676	932	1079	1303	1279	1026	698	408	0	8672
1967	77	38	41	46	56	76	68	145	252	420	593	885	1126	1310	1279	1109	694	440	1	8656
1968	86	52	50	53	65	83	119	154	283	441	595	907	1205	1268	1238	1028	724	441	4	8796
1964-68	430	249	210	256	280	364	486	805	1338	2046	3163	4604	5735	6517	6445	5426	3677	2156	7	44194
1969	78	55	52	63	75	91	121	166	300	444	688	981	1246	1454	1356	1111	811	474	6	9572
1970	69	42	44	61	80	102	109	159	279	477	659	1058	1265	1385	1369	1067	742	538	5	9510
1971	73	44	39	59	81	90	106	168	319	528	643	1055	1320	1521	1393	1180	777	495	2	9893
1972	71	47	42	71	74	109	111	158	352	517	792	1100	1435	1568	1490	1221	843	594	4	10599
1973	85	46	52	77	95	111	131	174	336	558	842	1078	1548	1659	1698	1353	894	603	38	11378
1969-73	376	234	229	331	405	503	578	825	1586	2524	3624	5272	6814	7587	7306	5932	4067	2704	55	50952
1974	83	36	67	72	134	134	155	198	336	589	907	1151	1560	1797	1759	1336	891	651	66	11922
1975	81	41	56	82	103	144	147	177	343	541	905	1144	1643	1786	1785	1415	902	603	47	11945
1976	70	46	57	104	117	151	148	170	313	639	1013	1336	1753	2003	1935	1480	1011	645	34	13025
1977	83	54	45	68	111	156	175	183	321	634	1019	1319	1690	1950	1875	1500	932	643	31	12789
1978	47	40	49	105	125	151	156	166	313	558	1081	1417	1885	2206	2077	1710	1119	757	40	14002
1974-78	364	217	274	431	590	736	781	894	1626	2961	4925	6367	8531	9742	9431	7441	4855	3299	218	63683
1979	79	44	61	97	109	133	164	239	299	623	1025	1532	1798	2174	2095	1737	1066	757	31	14063
1980	74	31	36	91	97	162	219	228	328	557	1012	1580	1754	2365	2181	1822	1156	794	35	14522
1981	71	37	42	88	124	146	204	234	304	552	1008	1620	1990	2409	2357	1805	1180	794	43	15008
1982	80	38	39	74	114	141	194	232	328	550	1007	1583	2047	2425	2464	1946	1171	795	50	15278
1983	70	63	40	75	129	149	179	248	326	548	1013	1674	2211	2433	2578	1972	1302	854	52	15916
1979-83	374	213	218	425	573	731	960	1181	1585	2830	5065	7989	9800	11806	11675	9282	5875	3994	211	74787

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

ALL MALIGNANT NEOPLASMS (EXCLUDING SKIN) (ICD9 140-208 EXCL. 173)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	23.0	14.8	12.7	20.8	27.4	28.4	47.5	77.5	125.3	217.9	387.6	711.5	1141.4	1560.0	2023.4	2596.9	3133.9	3286.8	286.9	278.6	27.5
1965	21.8	11.0	12.6	18.6	22.3	41.0	47.5	65.2	106.7	205.3	366.1	618.7	878.3	1346.4	1825.6	2361.7	2953.7	3021.6	254.2	245.7	24.4
1966	24.1	15.8	9.7	15.1	21.2	28.8	37.5	70.1	124.0	195.4	377.7	633.6	902.2	1399.6	1816.8	2201.7	2684.6	2914.3	252.1	245.0	24.7
1967	20.6	9.5	11.3	14.4	21.4	33.6	29.8	60.3	105.7	208.5	326.0	582.2	916.2	1378.9	1803.9	2334.7	2628.8	3034.5	246.7	239.8	24.1
1968	23.9	12.9	13.4	16.1	23.2	34.9	51.9	64.3	117.5	211.1	325.5	577.0	957.1	1304.5	1741.2	2150.6	2701.5	2940.0	245.8	239.5	23.9
1964-68	22.7	12.8	11.9	16.9	23.0	33.4	42.8	67.5	115.7	207.7	356.0	622.8	957.4	1396.1	1840.9	2326.8	2817.6	3036.6	256.8	249.4	24.9
1969	22.6	13.6	13.5	18.6	25.2	36.2	52.5	69.7	124.3	204.1	375.8	604.1	966.6	1454.0	1888.6	2309.8	2981.6	3019.1	262.0	255.0	25.4
1970	20.6	10.4	11.1	17.4	25.2	37.9	46.3	66.6	115.4	211.0	356.4	633.2	955.4	1344.7	1877.9	2200.0	2669.1	3280.5	254.4	248.5	24.9
1971	22.4	11.0	9.7	16.3	24.2	31.4	43.8	70.2	131.8	227.1	340.6	619.1	969.9	1433.6	1872.3	2413.1	2755.3	2861.3	258.8	252.2	25.3
1972	21.9	12.1	10.3	19.1	22.1	35.8	44.3	66.5	144.7	221.5	398.6	644.0	1017.7	1451.9	1942.6	2471.7	2937.3	3318.4	271.9	264.5	26.1
1973	26.3	12.4	12.5	20.2	28.1	34.7	50.3	73.5	137.9	237.2	403.1	635.6	1063.9	1506.8	2133.2	2716.9	3072.2	3313.2	286.2	277.4	27.3
1969-73	22.7	11.9	11.4	18.3	24.9	35.2	47.4	69.3	130.9	220.5	375.8	627.4	996.3	1439.1	1946.2	2424.2	2884.4	3162.6	266.9	259.8	25.8
1974	25.7	10.1	16.0	18.3	38.3	40.0	56.4	82.4	137.5	249.2	414.3	682.7	1037.9	1591.7	2137.3	2640.3	3051.4	3518.9	293.3	284.3	27.9
1975	25.3	11.9	13.4	20.5	28.5	41.4	51.4	72.6	141.4	227.3	404.9	665.1	1058.6	1539.7	2114.9	2700.4	3089.0	3173.7	288.0	278.4	27.4
1976	22.5	13.5	13.9	25.2	31.8	42.4	50.1	68.7	130.0	267.4	446.1	746.4	1113.0	1662.2	2239.6	2720.6	3404.0	3324.7	306.4	296.6	29.1
1977	26.8	16.0	11.3	16.4	29.5	44.2	55.4	72.0	134.0	263.7	447.7	698.6	1071.0	1548.8	2123.4	2654.9	3086.1	3264.0	293.2	285.1	28.0
1978	15.2	11.9	12.8	25.1	32.7	42.8	47.3	62.6	131.2	230.8	472.1	712.1	1196.8	1695.6	2290.0	2903.2	3633.1	3785.0	313.1	303.5	29.5
1974-78	23.1	12.6	13.5	21.1	32.1	42.2	52.0	71.5	134.8	247.7	437.4	701.7	1096.1	1609.2	2182.6	2727.6	3256.2	3415.1	299.0	289.8	28.4
1979	25.7	13.3	16.7	23.1	28.2	37.5	48.3	87.1	124.2	258.5	445.3	734.8	1147.4	1616.4	2235.9	2838.2	3352.2	3692.7	307.0	298.7	29.0
1980	24.2	9.5	10.2	21.6	24.9	45.2	62.7	80.7	134.8	233.3	434.5	742.5	1088.1	1703.9	2248.5	2896.7	3492.4	3835.7	309.0	299.7	29.1
1981	23.4	11.7	12.1	21.3	31.6	40.7	57.1	80.3	123.0	233.4	428.0	750.7	1178.9	1707.3	2340.6	2789.8	3460.4	3745.3	312.0	302.9	29.7
1982	26.1	12.2	11.3	18.4	28.5	38.6	55.0	74.1	129.1	233.6	425.1	725.5	1151.3	1699.4	2355.6	2926.3	3280.1	3680.6	309.9	300.4	29.5
1983	22.7	20.4	11.7	19.4	31.4	39.9	50.8	75.6	123.4	232.7	427.2	757.5	1177.9	1709.8	2363.0	2878.8	3444.4	3829.6	315.3	304.7	29.8
1979-83	24.4	13.4	12.5	20.8	29.0	40.4	54.8	79.3	126.9	238.4	431.9	742.3	1150.0	1688.0	2311.4	2866.6	3405.8	3757.3	310.8	301.4	29.4

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

ALL MALIGNANT NEOPLASMS (EXCLUDING SKIN) (ICD9 140-208 EXCL. 173)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	83	36	33	58	56	105	204	403	667	742	892	926	1008	1039	1081	864	649	398	1	9245
1965	69	46	27	37	62	110	230	375	588	726	824	823	899	895	971	803	572	406	1	8464
1966	60	43	24	49	70	113	218	356	655	726	806	906	917	986	933	856	620	427	0	8765
1967	63	37	18	42	70	124	175	372	638	777	888	894	936	992	979	871	591	412	1	8880
1968	57	33	41	37	69	129	228	376	610	793	856	938	942	948	942	826	569	437	2	8833
1964-68	332	195	143	223	327	581	1055	1882	3158	3764	4266	4487	4702	4860	4906	4220	3001	2080	5	44187
1969	53	44	26	47	87	126	216	362	616	854	949	1018	1045	1065	1060	905	658	523	2	9656
1970	56	27	50	61	81	119	194	319	585	850	868	1056	1078	1003	1025	864	680	502	4	9422
1971	53	35	30	45	95	152	215	361	595	908	947	1139	1084	1086	1061	966	698	536	1	10007
1972	50	43	55	68	94	136	248	325	598	915	988	1124	1211	1178	1113	998	762	627	7	10540
1973	74	37	49	65	124	191	233	370	633	915	1123	1205	1307	1259	1233	1167	854	717	40	11596
1969-73	286	186	210	286	481	724	1106	1737	3027	4442	4875	5542	5725	5591	5492	4900	3652	2905	54	51221
1974	58	37	46	76	111	203	231	379	612	983	1224	1241	1364	1419	1321	1193	890	772	48	12208
1975	58	33	29	82	122	171	254	360	592	935	1222	1294	1398	1361	1307	1127	843	780	42	12010
1976	56	25	51	88	142	241	298	378	601	996	1277	1393	1523	1559	1368	1189	960	772	27	12944
1977	49	35	36	69	134	209	290	367	554	907	1208	1390	1508	1549	1447	1172	919	776	24	12643
1978	56	34	35	74	118	188	317	403	594	928	1201	1530	1632	1601	1452	1282	1016	917	26	13404
1974-78	277	164	197	389	627	1012	1390	1887	2953	4749	6132	6848	7425	7489	6895	5963	4628	4017	167	63209
1979	61	33	29	64	114	227	307	407	575	858	1184	1616	1616	1685	1560	1373	1057	942	22	13730
1980	62	41	33	75	110	219	340	428	590	891	1158	1635	1713	1719	1567	1367	1095	934	30	14007
1981	47	29	32	68	126	209	354	472	615	875	1218	1599	1675	1786	1714	1410	1152	940	26	14347
1982	53	28	43	57	126	223	330	454	564	873	1191	1507	1766	1870	1712	1475	1155	1086	48	14561
1983	54	35	41	71	117	213	336	499	600	874	1205	1646	1842	1869	1964	1498	1217	1160	59	15308
1979-83	277	166	178	335	593	1091	1667	2260	2944	4371	5956	8003	8612	8929	8517	7123	5676	5070	185	71953

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000# FEMALES

ALL MALIGNANT NEOPLASMS (EXCLUDING SKIN) (ICD9 140-208 EXCL. 173)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	22.2	10.1	10.4	21.8	25.9	50.7	93.4	172.2	301.0	400.9	542.2	671.0	854.2	1022.6	1302.4	1474.4	1875.7	1809.1	277.9	248.8	24.0
1965	18.6	12.6	8.3	13.2	27.2	52.2	105.2	161.2	255.7	383.5	485.6	579.6	742.4	863.9	1134.3	1329.5	1593.3	1765.2	248.1	222.0	21.5
1966	16.5	11.4	7.1	16.6	28.6	51.7	98.8	152.8	277.1	371.4	462.2	619.7	736.5	929.3	1053.0	1382.9	1666.7	1801.7	250.0	223.2	21.5
1967	17.8	9.7	5.2	13.6	26.7	54.5	78.7	161.0	265.7	382.9	496.9	590.5	729.0	914.3	1086.6	1346.2	1515.4	1641.4	245.8	220.6	21.5
1968	16.7	8.6	11.4	11.5	24.6	54.3	102.0	164.2	254.2	373.9	472.1	597.1	713.1	851.8	1032.9	1232.8	1398.0	1649.1	238.3	214.7	20.9
1964-68	18.4	10.5	8.5	15.1	26.6	52.8	95.6	162.3	270.4	382.2	491.1	610.7	753.4	915.1	1118.8	1350.0	1601.4	1729.0	251.5	225.5	21.9
1969	16.2	11.5	7.1	14.2	29.2	50.6	95.8	159.0	259.7	384.7	517.4	625.3	770.1	929.3	1144.7	1315.4	1551.9	1848.1	253.7	228.0	22.2
1970	17.5	7.0	13.3	17.8	25.3	45.0	84.2	139.6	249.6	367.8	464.4	626.3	772.2	849.3	1082.4	1225.5	1538.5	1656.8	240.6	216.2	21.2
1971	17.0	9.2	7.8	12.8	28.0	54.2	91.1	157.5	256.4	382.6	492.5	657.6	753.3	892.4	1092.7	1339.8	1510.8	1649.2	248.6	223.5	21.7
1972	16.2	11.6	14.1	18.8	27.9	44.6	101.7	141.8	257.6	386.1	484.6	643.4	812.8	940.9	1123.1	1343.2	1604.2	1828.0	255.3	228.9	22.2
1973	24.1	10.5	12.4	17.6	36.3	59.8	91.3	161.1	272.7	386.2	520.1	688.6	848.7	979.0	1211.2	1535.5	1742.9	1997.2	273.6	244.2	23.4
1969-73	18.2	9.9	11.0	16.3	29.4	51.1	92.9	151.8	259.2	381.5	496.2	648.7	792.7	919.3	1131.7	1354.7	1592.7	1801.0	254.6	228.4	22.2
1974	18.9	10.9	11.5	20.0	31.3	60.5	86.0	161.7	261.9	417.2	537.5	704.3	855.2	1071.8	1259.3	1531.5	1776.4	2036.9	280.4	250.1	24.1
1975	19.1	10.0	7.3	21.2	33.2	49.0	90.5	151.1	253.5	398.6	522.7	713.3	842.2	1006.7	1209.1	1408.7	1633.7	1979.7	269.5	240.9	23.4
1976	19.0	7.7	13.1	22.3	37.8	67.0	102.5	155.7	257.1	427.5	536.8	731.6	902.3	1110.4	1233.5	1437.7	1801.1	1864.7	283.5	254.1	24.5
1977	16.6	10.9	9.5	17.3	34.9	58.2	92.7	146.6	237.5	389.4	511.0	687.4	880.8	1063.9	1260.5	1383.7	1658.8	1792.1	269.5	241.4	23.7
1978	19.1	10.7	9.6	18.4	30.4	51.8	96.4	154.0	254.8	397.6	509.5	713.3	947.2	1065.9	1223.3	1466.8	1776.2	2042.3	278.0	248.7	24.1
1974-78	18.5	10.0	10.2	19.8	33.5	57.3	93.8	153.7	253.0	406.1	523.4	709.9	886.2	1064.1	1237.0	1444.9	1729.4	1941.5	276.2	247.1	24.0
1979	20.8	10.6	8.4	15.8	29.2	61.9	90.1	149.9	243.4	367.8	506.6	717.9	932.5	1087.8	1269.3	1522.2	1773.5	2000.0	277.1	247.5	24.1
1980	21.3	13.3	9.8	18.5	28.0	58.6	96.2	152.8	245.9	383.9	495.5	710.3	958.1	1067.0	1241.7	1465.2	1771.8	1902.2	276.1	247.5	24.0
1981	16.2	9.7	9.7	17.2	31.7	55.6	97.1	162.9	251.3	378.0	521.9	686.0	892.4	1086.4	1315.4	1464.2	1802.8	1825.2	276.3	246.6	24.2
1982	18.1	9.4	13.2	14.9	31.4	58.3	91.1	145.5	223.6	377.3	509.4	645.4	890.6	1118.4	1273.8	1473.5	1766.1	2011.1	272.4	242.6	23.7
1983	18.3	11.9	12.7	19.3	28.6	54.7	92.1	151.8	228.3	376.2	515.8	701.3	875.9	1119.2	1406.9	1444.6	1787.1	2052.7	278.9	248.1	24.5
1979-83	19.0	11.0	10.7	17.1	29.8	57.8	93.3	152.5	238.2	376.6	509.8	691.9	908.1	1096.1	1303.5	1472.9	1780.4	1960.6	276.1	246.4	24.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF LIP (ICD9 140)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	1	2	2	5	5	13	13	21	25	40	32	31	25	16	12	0	243
1965	0	0	0	0	1	1	5	4	10	25	22	28	28	34	39	23	17	13	0	250
1966	0	0	0	0	1	1	3	7	11	14	23	29	28	26	24	25	18	6	0	216
1967	0	0	0	0	1	3	5	2	12	15	25	26	24	30	25	24	10	10	0	212
1968	0	0	0	0	0	1	5	8	10	4	16	27	28	36	24	27	14	15	0	215
1964-68	0	0	0	1	5	8	23	26	56	71	107	135	148	158	143	124	75	56	0	1136
1969	0	0	0	0	0	1	4	2	12	11	23	19	24	33	30	23	24	13	0	219
1970	0	0	0	0	0	0	0	6	4	16	23	28	43	29	36	19	15	12	2	233
1971	0	1	0	0	0	0	5	2	14	21	19	30	36	28	27	34	15	12	0	244
1972	1	1	0	0	1	1	6	5	11	13	22	27	22	34	34	23	19	16	0	236
1973	0	0	0	0	0	1	1	8	10	18	17	21	29	23	19	15	19	17	0	198
1969-73	1	2	0	0	1	3	16	23	51	79	104	125	154	147	146	114	92	70	2	1130
1974	0	1	0	0	0	0	1	11	9	10	22	24	33	37	28	23	19	17	2	237
1975	0	0	0	0	0	3	2	4	17	20	24	19	31	31	31	23	15	14	3	237
1976	0	0	0	0	0	5	4	4	8	14	12	33	37	24	31	18	17	8	1	216
1977	0	0	0	0	1	3	4	3	6	7	22	27	27	25	29	19	16	16	0	205
1978	0	0	0	1	0	4	2	1	1	13	17	21	22	33	19	29	25	17	5	210
1974-78	0	1	0	1	1	15	13	23	41	64	97	124	150	150	138	112	92	72	11	1105
1979	0	0	0	0	1	1	2	4	6	20	10	31	26	35	30	28	17	17	1	229
1980	0	0	0	0	1	4	3	6	7	9	16	26	28	40	31	30	16	17	6	240
1981	0	0	0	1	0	0	6	7	6	10	17	24	22	34	27	16	16	22	3	211
1982	0	0	0	0	0	1	1	7	9	9	18	24	28	30	37	29	14	14	3	224
1983	0	0	0	0	1	1	4	3	4	10	17	23	27	31	38	24	14	12	4	213
1979-83	0	0	0	1	3	7	16	27	32	58	78	128	131	170	163	127	77	82	17	1117

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF LIP (ICD9 140)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.4	0.9	1.0	2.2	2.1	6.0	6.9	12.4	17.9	35.1	35.6	45.4	55.1	63.0	88.2	7.3	7.2	0.8
1965	0.0	0.0	0.0	0.0	0.4	0.5	2.2	1.7	4.4	13.1	12.6	19.6	24.0	37.2	56.2	50.1	65.6	93.5	7.4	7.3	0.9
1966	0.0	0.0	0.0	0.0	0.4	0.5	1.3	2.9	4.7	7.2	12.8	19.7	23.4	27.9	34.1	53.6	69.2	42.9	6.2	6.0	0.7
1967	0.0	0.0	0.0	0.0	0.4	1.3	2.2	0.8	5.0	7.4	13.7	17.1	19.5	31.6	35.3	50.5	37.9	69.0	6.0	5.9	0.7
1968	0.0	0.0	0.0	0.0	0.0	0.4	2.2	3.3	4.2	1.9	8.8	17.2	22.2	37.0	33.8	56.5	52.2	100.0	6.0	5.9	0.7
1964-68	0.0	0.0	0.0	0.1	0.4	0.7	2.0	2.2	4.8	7.2	12.0	18.3	24.7	33.8	40.8	53.2	57.5	78.9	6.6	6.4	0.7
1969	0.0	0.0	0.0	0.0	0.0	0.4	1.7	0.8	5.0	5.1	12.6	11.7	18.6	33.0	41.8	47.8	88.2	82.8	6.0	5.8	0.7
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.5	1.7	7.1	12.4	16.8	32.5	28.2	49.4	39.2	54.0	73.2	6.1	6.1	0.7
1971	0.0	0.2	0.0	0.0	0.0	0.0	2.1	0.8	5.8	9.0	10.1	17.6	26.5	26.4	36.3	69.5	53.2	69.4	6.4	6.2	0.7
1972	0.3	0.3	0.0	0.0	0.3	0.3	2.4	2.1	4.5	5.6	11.1	15.8	15.6	31.5	44.3	46.6	66.2	89.4	6.0	5.9	0.7
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.4	3.4	4.1	7.7	8.1	12.4	19.9	20.9	23.9	30.1	65.3	93.4	5.0	4.9	0.5
1969-73	0.1	0.1	0.0	0.0	0.1	0.2	1.3	1.9	4.2	6.9	10.8	14.9	22.5	27.9	38.9	46.6	65.2	81.9	5.9	5.8	0.6
1974	0.0	0.3	0.0	0.0	0.0	0.0	0.4	4.6	3.7	4.2	10.1	14.2	22.0	32.8	34.0	45.5	65.1	91.9	5.8	5.7	0.6
1975	0.0	0.0	0.0	0.0	0.0	0.9	0.7	1.6	7.0	8.4	10.7	11.0	20.0	26.7	36.7	43.9	51.4	73.7	5.6	5.5	0.6
1976	0.0	0.0	0.0	0.0	0.0	1.4	1.4	1.6	3.3	5.9	5.3	18.4	23.5	19.9	35.9	33.1	57.2	41.2	5.0	4.9	0.6
1977	0.0	0.0	0.0	0.0	0.3	0.9	1.3	1.2	2.5	2.9	9.7	14.3	17.1	19.9	32.8	33.6	53.0	81.2	4.7	4.6	0.5
1978	0.0	0.0	0.0	0.2	0.0	1.1	0.6	0.4	0.4	5.4	7.4	10.6	14.0	25.4	20.9	49.2	81.2	85.0	4.6	4.4	0.4
1974-78	0.0	0.1	0.0	0.0	0.1	0.9	0.9	1.8	3.4	5.4	8.6	13.7	19.3	24.8	31.9	41.1	61.7	74.5	5.1	5.0	0.6
1979	0.0	0.0	0.0	0.0	0.3	0.3	0.6	1.5	2.5	8.3	4.3	14.9	16.6	26.0	32.0	45.8	53.5	82.9	5.0	4.8	0.5
1980	0.0	0.0	0.0	0.0	0.3	1.1	0.9	2.1	2.9	3.8	6.9	12.2	17.4	28.8	32.0	47.7	48.3	82.1	5.0	4.8	0.5
1981	0.0	0.0	0.0	0.2	0.0	0.0	1.7	2.4	2.4	4.2	7.2	11.1	13.0	24.1	26.8	24.7	46.9	103.8	4.3	4.3	0.5
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.3	2.2	3.5	3.8	7.6	11.0	15.7	21.0	35.4	43.6	39.2	64.8	4.5	4.4	0.5
1983	0.0	0.0	0.0	0.0	0.2	0.3	1.1	0.9	1.5	4.2	7.2	10.4	14.4	21.8	34.8	35.0	37.0	53.8	4.1	4.0	0.5
1979-83	0.0	0.0	0.0	0.0	0.2	0.4	0.9	1.8	2.6	4.9	6.7	11.9	15.4	24.3	32.3	39.2	44.6	77.1	4.6	4.5	0.5

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF LIP (ICD9 140)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	1	0	2	1	1	1	3	2	1	3	1	0	16
1965	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	2	1	2	0	6
1966	0	0	0	0	0	0	0	1	0	1	0	1	3	1	3	2	0	0	0	12
1967	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	1	0	2	0	5
1968	0	0	0	0	0	0	0	2	0	3	1	2	2	0	0	0	0	0	0	10
1964-68	0	0	0	0	0	0	0	4	0	8	3	4	6	4	5	6	4	5	0	49
1969	0	0	0	0	0	0	0	0	1	1	2	2	1	1	1	2	2	1	0	14
1970	0	0	0	0	0	0	0	0	0	2	0	1	0	2	4	2	1	0	0	12
1971	0	0	0	1	0	0	0	0	2	1	1	2	6	3	1	2	1	1	0	21
1972	0	0	0	0	1	1	0	0	0	2	0	0	4	2	1	0	2	3	0	16
1973	0	0	0	0	0	0	0	0	1	0	2	2	3	2	6	6	2	3	1	28
1969-73	0	0	0	1	1	1	0	0	4	6	5	7	14	10	13	12	8	8	1	91
1974	0	0	0	0	0	0	0	0	1	2	3	2	3	2	2	2	4	3	0	24
1975	1	0	0	0	0	0	0	1	0	0	2	4	1	2	4	3	1	4	0	23
1976	1	0	0	0	0	0	0	2	0	2	2	1	4	2	6	5	7	1	0	33
1977	0	0	0	0	0	0	0	1	0	2	1	2	3	3	4	4	3	6	0	29
1978	0	0	0	0	0	0	0	0	0	2	3	1	1	6	4	7	3	4	0	31
1974-78	2	0	0	0	0	0	0	4	1	8	11	10	12	15	20	21	18	18	0	140
1979	0	0	0	0	0	0	0	0	1	2	5	4	5	8	3	6	3	3	0	40
1980	0	0	0	0	0	1	1	1	0	0	6	2	6	5	5	7	3	5	1	43
1981	0	0	0	0	1	0	0	0	2	2	1	2	5	6	3	5	6	0	1	34
1982	0	0	0	0	0	0	0	0	1	2	1	3	4	6	7	5	9	5	2	45
1983	0	0	0	0	0	0	0	1	0	2	1	3	3	4	4	9	10	4	2	43
1979-83	0	0	0	0	1	1	1	2	4	8	14	14	23	29	22	32	31	17	6	205

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF LIP (ICD9 140)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.1	0.6	0.7	0.8	3.0	2.4	1.7	8.7	4.5	0.5	0.4	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	3.3	2.8	8.7	0.2	0.1	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	0.0	0.7	2.4	0.9	3.4	3.2	0.0	0.0	0.3	0.3	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.6	0.0	0.0	0.0	0.0	1.5	0.0	8.0	0.1	0.1	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	1.4	0.6	1.3	1.5	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.8	0.3	0.5	1.0	0.8	1.1	1.9	2.1	4.2	0.3	0.2	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	1.1	1.2	0.7	0.9	1.1	2.9	4.7	3.5	0.4	0.3	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.6	0.0	1.7	4.2	2.8	2.3	0.0	0.3	0.3	0.0
1971	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.9	0.4	0.5	1.2	4.2	2.5	1.0	2.8	2.2	3.1	0.5	0.5	0.1
1972	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.0	0.8	0.0	0.0	2.7	1.6	1.0	0.0	4.2	8.7	0.4	0.3	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	1.1	1.9	1.6	5.9	7.9	4.1	8.4	0.6	0.5	0.1
1969-73	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.0	0.3	0.5	0.5	0.8	1.9	1.6	2.7	3.3	3.5	5.0	0.4	0.4	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.3	1.1	1.9	1.5	1.9	2.6	8.0	7.9	0.5	0.5	0.0
1975	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.9	2.2	0.6	1.5	3.7	3.7	1.9	10.2	0.5	0.4	0.0
1976	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.9	0.8	0.5	2.4	1.4	5.4	6.0	13.1	2.4	0.7	0.6	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	0.4	1.0	1.8	2.1	3.5	4.7	5.4	13.9	0.6	0.5	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.3	0.5	0.6	4.0	3.4	8.0	5.2	8.9	0.6	0.5	0.1
1974-78	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.1	0.7	0.9	1.0	1.4	2.1	3.6	5.1	6.7	8.7	0.6	0.5	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	2.1	1.8	2.9	5.2	2.4	6.7	5.0	6.4	0.8	0.7	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.0	0.0	2.6	0.9	3.4	3.1	4.0	7.5	4.9	10.2	0.8	0.7	0.1
1981	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.8	0.9	0.4	0.9	2.7	3.6	2.3	5.2	9.4	0.0	0.6	0.5	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.4	1.3	2.0	3.6	5.2	5.0	13.8	9.3	0.8	0.6	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.9	0.4	1.3	1.4	2.4	2.9	8.7	14.7	7.0	0.7	0.5	0.0
1979-83	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.3	0.7	1.2	1.2	2.4	3.6	3.4	6.6	9.7	6.6	0.7	0.6	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF TONGUE (ICD9 141)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	1	7	3	4	8	7	3	7	4	6	0	50
1965	0	0	0	0	0	0	0	0	5	5	6	7	7	11	8	5	3	1	0	58
1966	0	0	0	0	0	0	1	0	0	4	4	5	8	13	5	9	3	2	0	54
1967	0	0	0	1	0	0	1	1	0	5	7	11	3	9	7	8	4	3	0	60
1968	0	0	0	0	0	0	1	1	4	8	7	7	12	4	5	8	7	2	0	66
1964-68	0	0	0	1	0	0	3	2	10	29	27	34	38	44	28	37	21	14	0	288
1969	0	0	0	0	0	0	0	0	3	0	7	7	5	6	7	8	3	3	0	49
1970	0	0	0	0	0	0	0	1	5	4	3	10	1	8	13	3	6	2	0	56
1971	0	0	0	0	0	0	0	1	1	5	7	13	4	8	9	7	5	3	0	63
1972	0	0	0	0	0	0	2	1	5	7	9	10	8	8	10	5	2	4	0	71
1973	0	1	0	0	0	0	0	0	4	7	8	6	11	10	5	1	6	1	1	61
1969-73	0	1	0	0	0	0	2	3	18	23	34	46	29	40	44	24	22	13	1	300
1974	1	0	0	0	0	1	0	0	1	5	7	4	13	10	3	3	5	1	0	54
1975	0	0	0	0	1	1	0	2	2	3	12	8	13	12	10	6	1	3	0	74
1976	0	0	0	0	1	3	1	4	2	9	9	9	13	13	9	6	6	0	0	85
1977	0	0	0	0	1	0	1	0	4	9	13	12	19	10	10	11	5	1	2	98
1978	0	0	0	0	0	1	1	1	7	7	4	7	18	7	10	13	1	2	0	79
1974-78	1	0	0	0	3	6	3	7	16	33	45	40	76	52	42	39	18	7	2	390
1979	0	0	0	0	0	0	0	1	0	4	14	16	14	14	9	5	3	2	0	82
1980	0	0	0	0	0	0	2	3	4	9	14	22	17	5	10	5	2	4	0	97
1981	0	0	0	0	1	0	2	0	3	6	7	14	16	13	12	4	5	3	1	87
1982	0	0	0	0	0	0	2	2	3	7	10	13	20	14	18	9	3	0	1	102
1983	0	0	0	1	0	0	2	0	6	7	13	22	11	22	9	5	4	3	1	106
1979-83	0	0	0	1	1	0	8	6	16	33	58	87	78	68	58	28	17	12	3	474

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF TONGUE (ICD9 141)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	3.7	1.8	2.9	7.0	7.8	4.4	15.4	15.7	44.1	1.5	1.5	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.2	2.6	3.4	4.9	6.0	12.0	11.5	10.9	11.6	7.2	1.7	1.7	0.2
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	2.0	2.2	3.4	6.7	14.0	7.1	19.3	11.5	14.3	1.6	1.5	0.2
1967	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	0.0	2.5	3.8	7.2	2.4	9.5	9.9	16.8	15.2	20.7	1.7	1.6	0.2
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.7	3.8	3.8	4.5	9.5	4.1	7.0	16.7	26.1	13.3	1.8	1.8	0.2
1964-68	0.0	0.0	0.0	0.1	0.0	0.0	0.3	0.2	0.9	2.9	3.0	4.6	6.3	9.4	8.0	15.9	16.1	19.7	1.7	1.6	0.2
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.0	3.8	4.3	3.9	6.0	9.7	16.6	11.0	19.1	1.3	1.3	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.1	1.8	1.6	6.0	0.8	7.8	17.8	6.2	21.6	12.2	1.5	1.4	0.2
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.2	3.7	7.6	2.9	7.5	12.1	14.3	17.7	17.3	1.6	1.6	0.2
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	2.1	3.0	4.5	5.9	5.7	7.4	13.0	10.1	7.0	22.3	1.8	1.8	0.2
1973	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	1.6	3.0	3.8	3.5	7.6	9.1	6.3	2.0	20.6	5.5	1.5	1.5	0.2
1969-73	0.0	0.1	0.0	0.0	0.0	0.0	0.2	0.3	1.5	2.0	3.5	5.5	4.2	7.6	11.7	9.8	15.6	15.2	1.6	1.5	0.2
1974	0.3	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	2.1	3.2	2.4	8.6	8.9	3.6	5.9	17.1	5.4	1.3	1.3	0.1
1975	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.8	0.8	1.3	5.4	4.7	8.4	10.3	11.8	11.5	3.4	15.8	1.8	1.8	0.2
1976	0.0	0.0	0.0	0.0	0.3	0.8	0.3	1.6	0.8	3.8	4.0	5.0	8.3	10.8	10.4	11.0	20.2	0.0	2.0	2.0	0.2
1977	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.0	1.7	3.7	5.7	6.4	12.0	7.9	11.3	19.5	16.6	5.1	2.2	2.2	0.2
1978	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	2.9	2.9	1.7	3.5	11.4	5.4	11.0	22.1	3.2	10.0	1.8	1.8	0.2
1974-78	0.1	0.0	0.0	0.0	0.2	0.3	0.2	0.6	1.3	2.8	4.0	4.4	9.8	8.6	9.7	14.3	12.1	7.2	1.8	1.8	0.2
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.7	6.1	7.7	8.9	10.4	9.6	8.2	9.4	9.8	1.8	1.8	0.2
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.6	1.1	1.6	3.8	6.0	10.3	10.5	3.6	10.3	7.9	6.0	19.3	2.1	2.1	0.2
1981	0.0	0.0	0.0	0.0	0.3	0.0	0.6	0.0	1.2	2.5	3.0	6.5	9.5	9.2	11.9	6.2	14.7	14.2	1.8	1.8	0.2
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.6	1.2	3.0	4.2	6.0	11.2	9.8	17.2	13.5	8.4	0.0	2.0	2.0	0.3
1983	0.0	0.0	0.0	0.3	0.0	0.0	0.6	0.0	2.3	3.0	5.5	10.0	5.9	15.5	8.2	7.3	10.6	13.5	2.1	2.1	0.3
1979-83	0.0	0.0	0.0	0.0	0.1	0.0	0.5	0.4	1.3	2.8	4.9	8.1	9.2	9.7	11.5	8.6	9.9	11.3	2.0	2.0	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF TONGUE (ICD9 141)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	1	0	2	2	0	0	1	5	5	5	3	2	0	26
1965	0	0	0	0	0	0	0	1	1	5	4	2	1	2	4	1	3	0	0	24
1966	0	0	0	0	0	0	0	1	0	3	2	4	2	4	2	2	2	0	0	22
1967	0	0	0	0	0	0	0	0	0	4	8	10	2	2	2	2	6	4	0	40
1968	0	0	0	0	0	0	0	2	0	1	3	3	6	4	2	3	2	0	0	26
1964-68	0	0	0	0	0	0	1	4	3	15	17	19	12	17	15	13	16	6	0	138
1969	0	0	0	0	0	0	0	0	0	1	3	2	0	4	6	2	4	2	0	24
1970	0	0	0	1	1	0	2	0	3	1	0	2	4	6	2	5	2	2	0	31
1971	0	0	0	0	0	2	0	0	1	1	6	5	8	5	1	4	0	5	0	38
1972	0	0	0	0	0	0	0	2	1	0	3	2	9	1	4	3	3	6	0	34
1973	0	0	0	0	0	0	0	0	0	2	0	1	6	5	1	4	4	2	0	25
1969-73	0	0	0	1	1	2	2	2	5	5	12	12	27	21	14	18	13	17	0	152
1974	0	0	0	0	1	1	1	0	0	2	4	4	6	6	7	3	3	1	1	40
1975	0	0	0	0	0	0	2	1	2	6	2	4	6	4	6	3	4	2	0	42
1976	0	0	0	0	0	0	1	1	0	1	2	8	8	6	4	5	8	1	0	45
1977	0	0	0	2	1	0	1	1	1	3	4	10	8	6	5	4	1	1	0	48
1978	0	0	0	0	1	0	0	0	3	4	6	8	3	8	8	1	2	3	0	47
1974-78	0	0	0	2	3	1	5	3	6	16	18	34	31	30	30	16	18	8	1	222
1979	0	0	0	1	0	3	0	0	1	4	4	7	5	5	5	3	4	3	1	46
1980	0	0	0	0	0	0	0	2	0	3	6	3	6	7	5	4	3	5	0	44
1981	0	0	0	0	0	0	0	0	1	3	5	17	1	7	3	6	2	8	0	53
1982	0	0	0	0	0	2	1	2	3	5	5	1	4	8	8	2	2	4	0	47
1983	0	0	0	1	0	1	0	0	1	3	6	5	7	8	5	6	2	3	0	48
1979-83	0	0	0	2	0	6	1	4	6	18	26	33	23	35	26	21	13	23	1	238

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF TONGUE (ICD9 141)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.9	1.1	0.0	0.0	0.8	4.9	6.0	8.5	8.7	9.1	0.8	0.6	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.6	2.4	1.4	0.8	1.9	4.7	1.7	8.4	0.0	0.7	0.6	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.5	1.1	2.7	1.6	3.8	2.3	3.2	5.4	0.0	0.6	0.6	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.0	4.5	6.6	1.6	1.8	2.2	3.1	15.4	15.9	1.1	1.0	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.5	1.7	1.9	4.5	3.6	2.2	4.5	4.9	0.0	0.7	0.6	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.3	1.5	2.0	2.6	1.9	3.2	3.4	4.2	8.5	5.0	0.8	0.7	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.6	1.2	0.0	3.5	6.5	2.9	9.4	7.1	0.6	0.5	0.1
1970	0.0	0.0	0.0	0.3	0.3	0.0	0.9	0.0	1.3	0.4	0.0	1.2	2.9	5.1	2.1	7.1	4.5	6.6	0.8	0.7	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0	0.4	0.4	3.1	2.9	5.6	4.1	1.0	5.5	0.0	15.4	0.9	0.9	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	0.0	1.5	1.1	6.0	0.8	4.0	4.0	6.3	17.5	0.8	0.7	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.6	3.9	3.9	1.0	5.3	8.2	5.6	0.6	0.5	0.1
1969-73	0.0	0.0	0.0	0.1	0.1	0.1	0.2	0.2	0.4	0.4	1.2	1.4	3.7	3.5	2.9	5.0	5.7	10.5	0.8	0.7	0.1
1974	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.0	0.0	0.8	1.8	2.3	3.8	4.5	6.7	3.9	6.0	2.6	0.9	0.8	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.4	0.9	2.6	0.9	2.2	3.6	3.0	5.6	3.7	7.8	5.1	0.9	0.8	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.0	0.4	0.8	4.2	4.7	4.3	3.6	6.0	15.0	2.4	1.0	0.8	0.1
1977	0.0	0.0	0.0	0.5	0.3	0.0	0.3	0.4	0.4	1.3	1.7	4.9	4.7	4.1	4.4	4.7	1.8	2.3	1.0	1.0	0.1
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	1.3	1.7	2.5	3.7	1.7	5.3	6.7	1.1	3.5	6.7	1.0	0.9	0.1
1974-78	0.0	0.0	0.0	0.1	0.2	0.1	0.3	0.2	0.5	1.4	1.5	3.5	3.7	4.3	5.4	3.9	6.7	3.9	1.0	0.9	0.1
1979	0.0	0.0	0.0	0.2	0.0	0.8	0.0	0.0	0.4	1.7	1.7	3.1	2.9	3.2	4.1	3.3	6.7	6.4	0.9	0.8	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.0	1.3	2.6	1.3	3.4	4.3	4.0	4.3	4.9	10.2	0.9	0.8	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	2.1	7.3	0.5	4.3	2.3	6.2	3.1	15.5	1.0	0.9	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.5	0.3	0.6	1.2	2.2	2.1	0.4	2.0	4.8	6.0	2.0	3.1	7.4	0.9	0.8	0.1
1983	0.0	0.0	0.0	0.3	0.0	0.3	0.0	0.0	0.4	1.3	2.6	2.1	3.3	4.8	3.6	5.8	2.9	5.3	0.9	0.8	0.1
1979-83	0.0	0.0	0.0	0.1	0.0	0.3	0.1	0.3	0.5	1.6	2.2	2.9	2.4	4.3	4.0	4.3	4.1	8.9	0.9	0.8	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF MAJOR SALIVARY GLAND (ICD9 142)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	2	0	1	0	1	2	1	4	3	4	3	0	0	3	0	24
1965	0	0	1	1	0	0	1	1	1	3	1	3	4	1	1	5	3	2	0	28
1966	1	0	0	0	0	0	1	3	0	1	5	0	0	6	3	4	2	3	0	29
1967	0	2	1	0	0	2	0	0	3	4	3	2	5	2	2	4	5	2	0	37
1968	0	0	0	0	0	2	3	6	0	0	2	2	4	3	7	3	2	3	0	37
1964-68	1	2	2	1	2	4	6	10	5	10	12	11	16	16	16	16	12	13	0	155
1969	0	0	0	0	2	1	2	0	0	3	2	3	2	4	1	2	1	3	0	26
1970	0	0	1	0	0	0	0	2	1	3	0	4	4	1	0	4	3	1	0	24
1971	0	0	1	0	0	1	2	0	2	0	0	5	2	3	0	4	2	2	0	24
1972	0	0	0	0	1	2	0	3	0	0	1	2	6	3	4	3	2	6	0	33
1973	0	0	0	1	0	0	2	3	2	2	2	4	4	2	1	4	1	0	0	28
1969-73	0	0	2	1	3	4	6	8	5	8	5	18	18	13	6	17	9	12	0	135
1974	0	1	0	1	2	0	0	1	3	2	2	7	7	6	4	6	0	1	0	43
1975	0	1	1	0	0	2	1	3	2	0	3	5	2	5	3	4	5	1	0	38
1976	0	0	0	0	1	2	2	0	1	2	3	2	5	5	0	1	0	3	0	27
1977	0	0	1	0	1	1	2	2	1	2	4	4	4	4	4	2	4	1	0	37
1978	0	0	2	0	1	0	3	2	1	6	2	2	5	6	3	1	4	3	1	42
1974-78	0	2	4	1	5	5	8	8	8	12	14	20	23	26	14	14	13	9	1	187
1979	0	0	0	0	1	1	1	4	2	3	2	4	3	9	6	7	3	3	0	49
1980	0	0	0	1	0	0	1	0	6	1	3	8	2	3	1	5	4	2	0	37
1981	0	0	0	0	0	2	2	1	0	5	6	2	7	10	6	4	2	1	0	48
1982	0	0	2	0	0	1	1	2	4	1	6	5	6	3	10	2	3	4	0	50
1983	0	0	0	1	1	0	1	2	1	1	2	3	4	7	9	4	2	1	0	39
1979-83	0	0	2	2	2	4	6	9	13	11	19	22	22	32	32	22	14	11	0	223

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF MAJOR SALIVARY GLAND (ICD9 142)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.9	0.0	0.4	0.0	0.5	1.1	0.6	2.9	2.6	4.4	4.4	0.0	0.0	22.1	0.7	0.8	0.1
1965	0.0	0.0	0.3	0.3	0.0	0.0	0.4	0.4	0.4	1.6	0.6	2.1	3.4	1.1	1.4	10.9	11.6	14.4	0.8	0.8	0.1
1966	0.3	0.0	0.0	0.0	0.0	0.0	0.4	1.3	0.0	0.5	2.8	0.0	0.0	6.4	4.3	8.6	7.7	21.4	0.8	0.8	0.1
1967	0.0	0.5	0.3	0.0	0.0	0.9	0.0	0.0	1.3	2.0	1.6	1.3	4.1	2.1	2.8	8.4	18.9	13.8	1.0	1.0	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.8	1.3	2.5	0.0	0.0	1.1	1.3	3.2	3.1	9.8	6.3	7.5	20.0	1.0	1.0	0.1
1964-68	0.1	0.1	0.1	0.1	0.2	0.4	0.5	0.8	0.4	1.0	1.4	1.5	2.7	3.4	4.6	6.9	9.2	18.3	0.9	0.9	0.1
1969	0.0	0.0	0.0	0.0	0.7	0.4	0.9	0.0	0.0	1.4	1.1	1.8	1.6	4.0	1.4	4.2	3.7	19.1	0.7	0.7	0.1
1970	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.8	0.4	1.3	0.0	2.4	3.0	1.0	0.0	8.2	10.8	6.1	0.6	0.6	0.0
1971	0.0	0.0	0.2	0.0	0.0	0.3	0.8	0.0	0.8	0.0	0.0	2.9	1.5	2.8	0.0	8.2	7.1	11.6	0.6	0.6	0.0
1972	0.0	0.0	0.0	0.0	0.3	0.7	0.0	1.3	0.0	0.0	0.5	1.2	4.3	2.8	5.2	6.1	7.0	33.5	0.8	0.8	0.1
1973	0.0	0.0	0.0	0.3	0.0	0.0	0.8	1.3	0.8	0.9	1.0	2.4	2.7	1.8	1.3	8.0	3.4	0.0	0.7	0.7	0.1
1969-73	0.0	0.0	0.1	0.1	0.2	0.3	0.5	0.7	0.4	0.7	0.5	2.1	2.6	2.5	1.6	6.9	6.4	14.0	0.7	0.7	0.1
1974	0.0	0.3	0.0	0.3	0.6	0.0	0.0	0.4	1.2	0.8	0.9	4.2	4.7	5.3	4.9	11.9	0.0	5.4	1.1	1.0	0.1
1975	0.0	0.3	0.2	0.0	0.0	0.6	0.3	1.2	0.8	0.0	1.3	2.9	1.3	4.3	3.6	7.6	17.1	5.3	0.9	0.9	0.1
1976	0.0	0.0	0.0	0.0	0.3	0.6	0.7	0.0	0.4	0.8	1.3	1.1	3.2	4.1	0.0	1.8	0.0	15.5	0.6	0.6	0.1
1977	0.0	0.0	0.3	0.0	0.3	0.3	0.6	0.8	0.4	0.8	1.8	2.1	2.5	3.2	4.5	3.5	13.2	5.1	0.9	0.8	0.1
1978	0.0	0.0	0.5	0.0	0.3	0.0	0.9	0.8	0.4	2.5	0.9	1.0	3.2	4.6	3.3	1.7	13.0	15.0	0.9	0.9	0.1
1974-78	0.0	0.1	0.2	0.0	0.3	0.3	0.5	0.6	0.7	1.0	1.2	2.2	3.0	4.3	3.2	5.1	8.7	9.3	0.9	0.9	0.1
1979	0.0	0.0	0.0	0.0	0.3	0.3	0.3	1.5	0.8	1.2	0.9	1.9	1.9	6.7	6.4	11.4	9.4	14.6	1.1	1.0	0.1
1980	0.0	0.0	0.0	0.2	0.0	0.0	0.3	0.0	2.5	0.4	1.3	3.8	1.2	2.2	1.0	7.9	12.1	9.7	0.8	0.7	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.6	0.6	0.3	0.0	2.1	2.5	0.9	4.1	7.1	6.0	6.2	5.9	4.7	1.0	1.0	0.1
1982	0.0	0.0	0.6	0.0	0.0	0.3	0.3	0.6	1.6	0.4	2.5	2.3	3.4	2.1	9.6	3.0	8.4	18.5	1.0	1.0	0.1
1983	0.0	0.0	0.0	0.3	0.2	0.0	0.3	0.6	0.4	0.4	0.8	1.4	2.1	4.9	8.2	5.8	5.3	4.5	0.8	0.7	0.1
1979-83	0.0	0.0	0.1	0.1	0.1	0.2	0.3	0.6	1.0	0.9	1.6	2.0	2.6	4.6	6.3	6.8	8.1	10.3	0.9	0.9	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF MAJOR SALIVARY GLAND (ICD9 142)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	1	1	0	3	2	2	2	2	2	1	2	3	0	1	3	2	0	27
1965	0	0	0	2	2	1	2	1	3	5	2	2	0	1	4	1	1	3	0	30
1966	0	0	0	0	0	1	3	0	2	2	1	1	3	3	1	3	2	0	0	22
1967	0	0	0	2	1	1	0	2	2	1	2	3	2	3	0	2	0	1	0	22
1968	0	0	1	0	1	1	0	3	1	2	5	3	1	4	4	2	1	1	0	30
1964-68	0	0	2	5	4	7	7	8	10	12	12	10	8	14	9	9	7	7	0	131
1969	0	0	0	0	1	1	1	0	1	1	3	1	0	1	3	4	2	2	0	21
1970	0	0	1	0	1	2	3	1	3	4	2	1	1	2	2	1	1	4	0	29
1971	0	0	0	1	1	2	0	0	3	0	1	1	3	2	6	3	2	3	0	28
1972	0	0	0	0	1	2	0	2	4	0	3	0	3	1	2	3	3	1	0	25
1973	0	0	0	0	0	3	0	3	1	4	1	3	1	2	2	2	1	1	0	24
1969-73	0	0	1	1	4	10	4	6	12	9	10	6	8	8	15	13	9	11	0	127
1974	0	0	0	0	1	3	3	1	6	0	4	1	3	5	5	3	1	2	0	38
1975	0	0	0	0	2	3	4	2	4	7	5	5	1	1	2	1	2	2	0	41
1976	0	0	0	0	1	4	4	3	2	7	4	5	5	7	0	2	1	4	0	49
1977	0	0	0	1	2	1	1	1	4	4	1	3	3	5	1	1	3	2	1	34
1978	0	0	0	3	1	2	3	0	1	3	9	6	2	6	4	4	5	3	0	52
1974-78	0	0	0	4	7	13	15	7	17	21	23	20	14	24	12	11	12	13	1	214
1979	0	0	0	0	2	1	1	2	2	1	2	3	3	6	6	3	3	2	0	37
1980	0	0	0	0	1	1	2	0	1	2	1	4	4	4	4	2	2	3	0	31
1981	0	0	0	0	0	1	0	3	0	1	1	3	4	1	2	4	1	4	0	25
1982	0	0	0	0	0	4	0	1	1	2	6	4	5	2	3	5	0	7	1	41
1983	1	0	0	2	4	3	2	1	1	1	1	3	3	6	6	2	1	4	0	41
1979-83	1	0	0	2	7	10	5	7	5	7	11	17	19	19	21	16	7	20	1	175

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF MAJOR SALIVARY GLAND (ICD9 142)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.3	0.4	0.0	1.4	0.9	0.9	0.9	1.1	1.2	0.7	1.7	3.0	0.0	1.7	8.7	9.1	0.8	0.8	0.1
1965	0.0	0.0	0.0	0.7	0.9	0.5	0.9	0.4	1.3	2.6	1.2	1.4	0.0	1.0	4.7	1.7	2.8	13.0	0.9	0.8	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.5	1.4	0.0	0.8	1.0	0.6	0.7	2.4	2.8	1.1	4.8	5.4	0.0	0.6	0.6	0.1
1967	0.0	0.0	0.0	0.6	0.4	0.4	0.0	0.9	0.8	0.5	1.1	2.0	1.6	2.8	0.0	3.1	0.0	4.0	0.6	0.6	0.1
1968	0.0	0.0	0.3	0.0	0.4	0.4	0.0	1.3	0.4	0.9	2.8	1.9	0.8	3.6	4.4	3.0	2.5	3.8	0.8	0.7	0.1
1964-68	0.0	0.0	0.1	0.3	0.3	0.6	0.6	0.7	0.9	1.2	1.4	1.4	1.3	2.6	2.1	2.9	3.7	5.8	0.7	0.7	0.1
1969	0.0	0.0	0.0	0.0	0.3	0.4	0.4	0.0	0.4	0.5	1.6	0.6	0.0	0.9	3.2	5.8	4.7	7.1	0.6	0.5	0.0
1970	0.0	0.0	0.3	0.0	0.3	0.8	1.3	0.4	1.3	1.7	1.1	0.6	0.7	1.7	2.1	1.4	2.3	13.2	0.7	0.7	0.1
1971	0.0	0.0	0.0	0.3	0.3	0.7	0.0	0.0	1.3	0.0	0.5	0.6	2.1	1.6	6.2	4.2	4.3	9.2	0.7	0.6	0.1
1972	0.0	0.0	0.0	0.0	0.3	0.7	0.0	0.9	1.7	0.0	1.5	0.0	2.0	0.8	2.0	4.0	6.3	2.9	0.6	0.5	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.9	0.0	1.3	0.4	1.7	0.5	1.7	0.6	1.6	2.0	2.6	2.0	2.8	0.6	0.5	0.1
1969-73	0.0	0.0	0.1	0.1	0.2	0.7	0.3	0.5	1.0	0.8	1.0	0.7	1.1	1.3	3.1	3.6	3.9	6.8	0.6	0.6	0.1
1974	0.0	0.0	0.0	0.0	0.3	0.9	1.1	0.4	2.6	0.0	1.8	0.6	1.9	3.8	4.8	3.9	2.0	5.3	0.9	0.8	0.1
1975	0.0	0.0	0.0	0.0	0.5	0.9	1.4	0.8	1.7	3.0	2.1	2.8	0.6	0.7	1.9	1.2	3.9	5.1	0.9	0.9	0.1
1976	0.0	0.0	0.0	0.0	0.3	1.1	1.4	1.2	0.9	3.0	1.7	2.6	3.0	5.0	0.0	2.4	1.9	9.7	1.1	1.0	0.1
1977	0.0	0.0	0.0	0.3	0.5	0.3	0.3	0.4	1.7	1.7	0.4	1.5	1.8	3.4	0.9	1.2	5.4	4.6	0.7	0.7	0.1
1978	0.0	0.0	0.0	0.7	0.3	0.6	0.9	0.0	0.4	1.3	3.8	2.8	1.2	4.0	3.4	4.6	8.7	6.7	1.1	0.9	0.1
1974-78	0.0	0.0	0.0	0.2	0.4	0.7	1.0	0.6	1.5	1.8	2.0	2.1	1.7	3.4	2.2	2.7	4.5	6.3	0.9	0.9	0.1
1979	0.0	0.0	0.0	0.0	0.5	0.3	0.3	0.7	0.8	0.4	0.9	1.3	1.7	3.9	4.9	3.3	5.0	4.2	0.7	0.7	0.1
1980	0.0	0.0	0.0	0.0	0.3	0.3	0.6	0.0	0.4	0.9	0.4	1.7	2.2	2.5	3.2	2.1	3.2	6.1	0.6	0.5	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.0	1.0	0.0	0.4	0.4	1.3	2.1	0.6	1.5	4.2	1.6	7.8	0.5	0.4	0.0
1982	0.0	0.0	0.0	0.0	0.0	1.0	0.0	0.3	0.4	0.9	2.6	1.7	2.5	1.2	2.2	5.0	0.0	13.0	0.7	0.7	0.1
1983	0.3	0.0	0.0	0.5	1.0	0.8	0.5	0.3	0.4	0.4	0.4	1.3	1.4	3.6	4.3	1.9	1.5	7.0	0.8	0.7	0.1
1979-83	0.1	0.0	0.0	0.1	0.4	0.5	0.3	0.5	0.4	0.6	0.9	1.5	2.0	2.3	3.2	3.3	2.2	7.7	0.7	0.6	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF GUM (ICD9 143)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	0	2	0	1	1	2	6	1	1	0	14
1965	0	0	0	0	0	0	0	0	2	0	1	0	1	3	3	1	2	1	0	14
1966	0	0	0	0	0	0	0	0	0	1	3	1	1	0	2	2	1	1	0	12
1967	0	0	0	0	0	0	0	0	0	1	2	1	1	3	1	1	1	0	0	11
1968	0	0	0	0	0	0	1	0	0	1	2	3	2	4	2	2	1	1	0	19
1964-68	0	0	0	0	0	0	1	0	2	3	10	5	6	11	10	12	6	4	0	70
1969	0	0	1	0	0	0	0	1	0	2	4	2	4	0	3	2	4	3	0	26
1970	0	0	0	0	0	0	0	0	0	0	2	1	3	0	4	2	2	0	0	14
1971	0	0	0	0	0	0	0	0	1	0	0	2	1	1	3	5	2	3	0	18
1972	0	0	0	0	0	0	0	0	1	1	1	1	3	3	1	1	3	0	0	15
1973	0	0	0	0	0	0	0	1	0	1	3	1	1	1	2	1	0	2	0	13
1969-73	0	0	1	0	0	0	0	2	2	4	10	7	12	5	13	11	11	8	0	86
1974	0	0	0	0	0	1	0	0	0	1	1	2	3	4	4	3	1	3	0	23
1975	0	0	0	0	0	0	0	0	1	1	1	3	3	4	3	3	1	4	0	24
1976	0	0	0	0	0	0	0	0	0	0	1	1	1	4	2	3	0	2	0	14
1977	0	0	0	0	2	0	0	1	1	0	0	2	1	6	3	3	0	0	0	19
1978	0	0	0	0	0	0	0	0	0	0	1	2	2	0	2	1	1	1	0	10
1974-78	0	0	0	0	2	1	0	1	2	2	4	10	10	18	14	13	3	10	0	90
1979	0	0	0	0	0	0	2	0	0	1	2	1	6	2	2	0	3	3	0	22
1980	1	0	0	0	0	0	0	0	0	0	0	2	3	3	1	3	2	0	0	15
1981	1	0	0	0	1	0	0	0	0	3	0	5	3	0	1	3	1	2	0	20
1982	0	0	0	0	0	0	0	0	1	1	3	3	4	5	2	3	1	1	1	25
1983	0	0	0	0	0	0	0	0	0	1	1	3	3	3	3	4	2	3	0	23
1979-83	2	0	0	0	1	0	2	0	1	6	6	14	19	13	9	13	9	9	1	105

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF GUM (ICD9 143)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.0	0.9	1.1	2.9	13.2	3.9	7.4	0.4	0.4	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.6	0.0	0.9	3.3	4.3	2.2	7.7	7.2	0.4	0.4	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.7	0.7	0.8	0.0	2.8	4.3	3.8	7.1	0.3	0.3	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.1	0.7	0.8	3.2	1.4	2.1	3.8	0.0	0.3	0.3	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	1.1	1.9	1.6	4.1	2.8	4.2	3.7	6.7	0.5	0.5	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.3	1.1	0.7	1.0	2.4	2.9	5.1	4.6	5.6	0.4	0.4	0.0
1969	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.0	0.9	2.2	1.2	3.1	0.0	4.2	4.2	14.7	19.1	0.7	0.7	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.6	2.3	0.0	5.5	4.1	7.2	0.0	0.4	0.4	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.2	0.7	0.9	4.0	10.2	7.1	17.3	0.5	0.4	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.6	2.1	2.8	1.3	2.0	10.5	0.0	0.4	0.4	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	1.4	0.6	0.7	0.9	2.5	2.0	0.0	11.0	0.3	0.3	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.3	1.0	0.8	1.8	0.9	3.5	4.5	7.8	9.4	0.5	0.4	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	0.5	1.2	2.0	3.5	4.9	5.9	3.4	16.2	0.6	0.6	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	1.7	1.9	3.4	3.6	5.7	3.4	21.1	0.6	0.6	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.6	0.6	3.3	2.3	5.5	0.0	10.3	0.3	0.3	0.0
1977	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.4	0.4	0.0	0.0	1.1	0.6	4.8	3.4	5.3	0.0	0.0	0.4	0.4	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.0	1.3	0.0	2.2	1.7	3.2	5.0	0.2	0.2	0.0
1974-78	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.2	0.2	0.4	1.1	1.3	3.0	3.2	4.8	2.0	10.4	0.4	0.4	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.4	0.9	0.5	3.8	1.5	2.1	0.0	9.4	14.6	0.5	0.5	0.0
1980	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.9	2.2	1.0	4.8	6.0	0.0	0.3	0.3	0.0
1981	0.3	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	1.3	0.0	2.3	1.8	0.0	1.0	4.6	2.9	9.4	0.4	0.4	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.3	1.4	2.2	3.5	1.9	4.5	2.8	4.6	0.5	0.5	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.4	1.6	2.1	2.7	5.8	5.3	13.5	0.5	0.4	0.0
1979-83	0.1	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.1	0.5	0.5	1.3	2.2	1.9	1.8	4.0	5.2	8.5	0.4	0.4	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF GUM (ICD9 143)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	1	1	1	0	0	0	1	0	0	0	4
1965	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	1	1	0	5
1966	0	0	0	0	0	0	0	1	1	0	0	1	1	2	0	2	1	0	0	9
1967	0	0	0	0	0	0	0	0	0	0	1	1	2	1	1	1	0	0	0	7
1968	0	0	1	0	0	0	0	0	0	1	4	0	1	1	2	1	0	2	0	13
1964-68	0	0	1	0	0	0	0	1	1	3	6	4	5	4	3	5	2	3	0	38
1969	0	0	0	0	0	0	0	1	0	0	1	2	4	0	0	4	2	0	0	14
1970	0	0	0	0	0	0	0	0	1	2	0	1	2	0	3	2	1	1	0	13
1971	0	0	0	0	0	0	0	0	0	1	1	2	1	4	2	3	3	2	0	19
1972	0	0	0	0	0	0	0	0	0	0	1	0	1	2	1	0	2	0	0	7
1973	0	0	0	0	0	0	0	0	1	1	0	1	1	1	2	2	6	1	0	16
1969-73	0	0	0	0	0	0	0	1	2	4	3	6	9	7	8	11	14	4	0	69
1974	0	0	0	0	0	0	0	0	0	1	1	0	2	5	2	2	0	2	0	15
1975	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	4	2	0	0	8
1976	0	0	1	0	0	0	0	0	0	3	0	1	2	1	0	3	0	1	0	12
1977	0	0	0	0	0	0	0	0	0	0	1	2	2	2	1	1	1	0	0	10
1978	0	0	0	0	0	0	1	0	0	1	1	1	2	0	2	1	1	1	0	11
1974-78	0	0	1	0	0	0	1	0	0	5	4	5	8	8	5	11	4	4	0	56
1979	0	0	0	0	1	0	0	0	2	1	1	0	1	1	1	1	2	2	0	13
1980	0	0	0	0	0	0	0	0	0	0	1	2	3	3	1	5	2	0	1	18
1981	0	0	0	0	0	0	0	0	2	0	0	0	1	4	1	0	2	2	1	13
1982	0	0	0	0	0	0	1	0	0	1	0	3	2	3	0	3	0	3	0	16
1983	0	0	0	0	0	0	0	0	0	0	3	1	4	3	1	1	0	0	1	14
1979-83	0	0	0	0	1	0	1	0	4	2	5	6	11	14	4	10	6	7	3	74

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF GUM (ICD9 143)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.6	0.7	0.0	0.0	0.0	1.7	0.0	0.0	0.1	0.1	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.7	0.8	0.0	0.0	0.0	2.8	4.3	0.2	0.1	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.0	0.7	0.8	1.9	0.0	3.2	2.7	0.0	0.3	0.2	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	1.6	0.9	1.1	1.5	0.0	0.0	0.2	0.2	0.0
1968	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.5	2.2	0.0	0.8	0.9	2.2	1.5	0.0	7.5	0.4	0.3	0.0
1964-68	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.3	0.7	0.5	0.8	0.8	0.7	1.6	1.1	2.5	0.2	0.2	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	1.2	2.9	0.0	0.0	5.8	4.7	0.0	0.4	0.3	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.0	0.6	1.4	0.0	3.2	2.8	2.3	3.3	0.3	0.3	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	1.2	0.7	3.3	2.1	4.2	6.5	6.2	0.5	0.4	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.7	1.6	1.0	0.0	4.2	0.0	0.2	0.1	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.6	0.6	0.8	2.0	2.6	12.2	2.8	0.4	0.3	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.3	0.3	0.7	1.2	1.2	1.6	3.0	6.1	2.5	0.3	0.3	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	1.3	3.8	1.9	2.6	0.0	5.3	0.3	0.3	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.6	0.0	0.0	0.0	5.0	3.9	0.0	0.2	0.1	0.0
1976	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.0	0.5	1.2	0.7	0.0	3.6	0.0	2.4	0.3	0.2	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.0	1.2	1.4	0.9	1.2	1.8	0.0	0.2	0.2	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	0.5	1.2	0.0	1.7	1.1	1.7	2.2	0.2	0.2	0.0
1974-78	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.4	0.3	0.5	1.0	1.1	0.9	2.7	1.5	1.9	0.2	0.2	0.0
1979	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.8	0.4	0.4	0.0	0.6	0.6	0.8	1.1	3.4	4.2	0.3	0.2	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	1.7	1.9	0.8	5.4	3.2	0.0	0.3	0.3	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.0	0.5	2.4	0.8	0.0	3.1	3.9	0.2	0.2	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	1.3	1.0	1.8	0.0	3.0	0.0	5.6	0.3	0.2	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.4	1.9	1.8	0.7	1.0	0.0	0.0	0.2	0.2	0.0
1979-83	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.3	0.2	0.4	0.5	1.2	1.7	0.6	2.1	1.9	2.7	0.3	0.2	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF FLOOR OF MOUTH (ICD9 144)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	0	0	2	2	3	5	4	1	4	5	4	3	0	33
1965	0	0	0	0	0	0	0	1	1	1	6	7	4	8	4	5	3	0	0	40
1966	0	0	0	0	0	0	0	1	0	2	4	6	1	3	4	0	2	1	0	24
1967	0	0	0	0	0	0	0	0	1	2	4	6	1	2	6	2	0	2	0	26
1968	0	0	0	0	0	0	0	0	1	4	5	9	5	4	3	4	1	1	0	37
1964-68	0	0	0	0	0	0	0	2	5	11	22	33	15	18	21	16	10	7	0	160
1969	0	0	0	0	0	0	0	0	2	1	5	10	8	3	3	2	2	0	0	36
1970	0	0	0	0	0	1	0	2	2	3	2	5	6	10	4	4	2	2	0	43
1971	0	0	0	0	0	0	2	0	2	4	6	5	7	7	3	3	3	2	0	44
1972	0	0	0	0	0	0	0	1	3	8	3	5	8	4	2	3	0	2	0	39
1973	0	0	0	0	0	0	0	0	3	9	9	4	5	4	6	0	2	2	0	44
1969-73	0	0	0	0	0	1	2	3	12	25	25	29	34	28	18	12	9	8	0	206
1974	0	0	0	0	0	1	0	0	1	8	5	6	10	11	5	2	1	0	0	50
1975	0	0	0	0	0	0	0	1	0	4	5	9	10	6	2	3	3	3	0	46
1976	0	0	0	0	0	0	0	1	1	9	10	7	10	4	4	2	2	3	1	54
1977	0	0	0	0	0	0	0	0	3	7	9	13	8	6	6	3	2	3	0	60
1978	0	0	0	0	0	0	0	0	3	2	12	15	11	7	7	3	6	2	0	68
1974-78	0	0	0	0	0	1	0	2	8	30	41	50	49	34	24	13	14	11	1	278
1979	0	0	0	0	0	0	0	1	4	7	11	5	9	8	5	1	3	2	1	57
1980	1	0	0	0	0	0	1	0	1	7	15	9	7	9	7	3	0	1	0	61
1981	0	0	0	0	0	0	1	2	5	5	7	15	11	6	6	5	2	3	0	68
1982	0	0	0	0	0	0	0	1	0	11	6	11	16	6	4	4	2	1	0	62
1983	0	0	0	0	0	1	0	2	2	4	8	12	7	16	5	2	2	3	1	65
1979-83	1	0	0	0	0	1	2	6	12	34	47	52	50	45	27	15	9	10	2	313

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF FLOOR OF MOUTH (ICD9 144)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.1	1.8	3.6	3.5	1.1	5.9	11.0	15.7	22.1	1.0	0.9	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	3.4	4.9	3.4	8.7	5.8	10.9	11.6	0.0	1.2	1.1	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.0	2.2	4.1	0.8	3.2	5.7	0.0	7.7	7.1	0.7	0.7	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.0	2.2	3.9	0.8	2.1	8.5	4.2	0.0	13.8	0.7	0.7	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.9	2.7	5.7	4.0	4.1	4.2	8.4	3.7	6.7	1.0	1.0	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.4	1.1	2.5	4.5	2.5	3.9	6.0	6.9	7.7	9.9	0.9	0.9	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.5	2.7	6.2	6.2	3.0	4.2	4.2	7.4	0.0	1.0	1.0	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.8	0.8	1.3	1.1	3.0	4.5	9.7	5.5	8.2	7.2	12.2	1.1	1.1	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.8	1.7	3.2	2.9	5.1	6.6	4.0	6.1	10.6	11.6	1.1	1.1	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.2	3.4	1.5	2.9	5.7	3.7	2.6	6.1	0.0	11.2	1.0	1.0	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	3.8	4.3	2.4	3.4	3.6	7.5	0.0	6.9	11.0	1.1	1.1	0.1
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.3	1.0	2.2	2.6	3.5	5.0	5.3	4.8	4.9	6.4	9.4	1.1	1.1	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	3.4	2.3	3.6	6.7	9.7	6.1	4.0	3.4	0.0	1.2	1.2	0.2
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.7	2.2	5.2	6.4	5.2	2.4	5.7	10.3	15.8	1.1	1.1	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	3.8	4.4	3.9	6.3	3.3	4.6	3.7	6.7	15.5	1.2	1.2	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	2.9	4.0	6.9	5.1	4.8	6.8	5.3	6.6	15.2	1.4	1.4	0.2
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.8	5.2	7.5	7.0	5.4	7.7	5.1	19.5	10.0	1.5	1.5	0.2
1974-78	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.7	2.5	3.6	5.5	6.3	5.6	5.6	4.8	9.4	11.4	1.3	1.3	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	2.9	4.8	2.4	5.7	5.9	5.3	1.6	9.4	9.8	1.2	1.3	0.1
1980	0.3	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	2.9	6.4	4.2	4.3	6.5	7.2	4.8	0.0	4.8	1.3	1.3	0.2
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.7	2.0	2.1	3.0	7.0	6.5	4.3	6.0	7.7	5.9	14.2	1.4	1.4	0.2
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	4.7	2.5	5.0	9.0	4.2	3.8	6.0	5.6	4.6	1.3	1.3	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.6	0.8	1.7	3.4	5.4	3.7	11.2	4.6	2.9	5.3	13.5	1.3	1.3	0.2
1979-83	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.4	1.0	2.9	4.0	4.8	5.9	6.4	5.3	4.6	5.2	9.4	1.3	1.3	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF FLOOR OF MOUTH (ICD9 144)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	0	0	0	0	1	1	0	0	1	2	0	0	0	5
1965	0	0	0	0	0	1	0	0	0	3	1	1	3	1	1	0	3	1	0	15
1966	0	0	0	0	0	0	0	0	1	0	0	2	0	1	1	0	0	0	0	5
1967	0	0	0	0	0	0	0	0	0	0	0	1	2	0	1	2	1	0	0	7
1968	0	0	0	0	0	0	0	0	1	1	0	1	3	0	0	1	0	1	0	8
1964-68	0	0	0	0	0	1	0	0	2	4	2	6	8	2	4	5	4	2	0	40
1969	0	0	0	0	0	0	0	0	0	1	2	0	3	2	0	2	0	1	0	11
1970	0	0	0	0	0	0	0	0	1	0	0	1	1	1	0	0	0	0	0	4
1971	0	0	0	0	0	0	0	0	2	5	2	1	0	1	1	0	2	1	0	15
1972	0	0	0	0	0	0	0	0	1	1	1	0	4	0	0	2	1	0	0	10
1973	0	0	0	0	0	0	0	1	0	2	4	4	3	1	1	1	0	1	1	19
1969-73	0	0	0	0	0	0	0	1	4	9	9	6	11	5	2	5	3	3	1	59
1974	0	0	0	0	0	0	0	0	1	2	3	5	1	3	1	2	2	0	0	20
1975	0	0	0	0	0	0	0	0	1	1	1	3	5	3	1	2	0	0	0	17
1976	0	0	0	0	0	0	0	0	2	2	3	4	3	4	3	1	0	1	0	23
1977	0	0	0	0	0	0	0	0	2	1	3	4	3	3	2	1	0	2	0	21
1978	0	0	0	0	0	0	0	0	2	2	4	5	4	3	2	2	0	2	0	26
1974-78	0	0	0	0	0	0	0	0	8	8	14	21	16	16	9	8	2	5	0	107
1979	0	0	0	0	0	0	0	0	1	2	2	3	6	3	3	2	0	1	0	23
1980	0	0	0	0	0	0	0	0	1	1	2	2	4	4	3	1	0	3	0	21
1981	0	0	0	0	0	0	0	0	1	0	0	3	1	5	1	1	1	0	0	13
1982	0	0	0	0	0	0	0	0	1	0	2	4	5	6	1	1	1	6	0	27
1983	0	0	0	0	0	0	0	1	0	2	0	10	4	6	4	1	2	0	0	30
1979-83	0	0	0	0	0	0	0	1	4	5	6	22	20	24	12	6	4	10	0	114

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF FLOOR OF MOUTH (ICD9 144)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	0.0	0.0	1.2	3.4	0.0	0.0	0.2	0.1	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	1.6	0.6	0.7	2.5	1.0	1.2	0.0	8.4	4.3	0.5	0.4	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.4	0.0	0.9	1.1	0.0	0.0	0.0	0.1	0.1	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	1.6	0.0	1.1	3.1	2.6	0.0	0.2	0.2	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.0	0.6	2.3	0.0	0.0	1.5	0.0	3.8	0.2	0.2	0.0
1964-68	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.2	0.4	0.2	0.8	1.3	0.4	0.9	1.6	2.1	1.7	0.2	0.2	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.1	0.0	2.2	1.7	0.0	2.9	0.0	3.5	0.3	0.3	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.6	0.7	0.8	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	2.1	1.0	0.6	0.0	0.8	1.0	0.0	4.3	3.1	0.4	0.3	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.0	2.7	0.0	0.0	2.7	2.1	0.0	0.2	0.2	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.8	1.9	2.3	1.9	0.8	1.0	1.3	0.0	2.8	0.4	0.4	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.8	0.9	0.7	1.5	0.8	0.4	1.4	1.3	1.9	0.3	0.3	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.3	2.8	0.6	2.3	1.0	2.6	4.0	0.0	0.5	0.4	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	1.7	3.0	2.2	0.9	2.5	0.0	0.0	0.4	0.4	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.9	1.3	2.1	1.8	2.8	2.7	1.2	0.0	2.4	0.5	0.5	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	1.3	2.0	1.8	2.1	1.7	1.2	0.0	4.6	0.4	0.4	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.9	1.7	2.3	2.3	2.0	1.7	2.3	0.0	4.5	0.5	0.5	0.1
1974-78	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.7	1.2	2.2	1.9	2.3	1.6	1.9	0.7	2.4	0.5	0.4	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.9	1.3	3.5	1.9	2.4	2.2	0.0	2.1	0.5	0.5	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.9	0.9	2.2	2.5	2.4	1.1	0.0	6.1	0.4	0.4	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.3	0.5	3.0	0.8	1.0	1.6	0.0	0.2	0.2	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	1.7	2.5	3.6	0.7	1.0	1.5	11.1	0.5	0.4	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.9	0.0	4.3	1.9	3.6	2.9	1.0	2.9	0.0	0.5	0.5	0.1
1979-83	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.4	0.5	1.9	2.1	2.9	1.8	1.2	1.3	3.9	0.4	0.4	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF MOUTH (ICD9 145)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	1	0	0	0	0	1	5	9	7	4	4	9	6	4	0	50
1965	0	0	0	0	0	0	0	1	2	3	0	1	2	2	5	5	4	4	0	29
1966	0	0	0	0	0	0	0	1	0	2	5	9	4	3	5	3	2	3	0	37
1967	0	0	0	0	0	0	0	0	2	1	1	3	1	5	4	5	6	1	0	29
1968	0	0	0	0	0	0	0	0	3	5	2	3	6	9	5	6	7	2	0	48
1964-68	0	0	0	0	1	0	0	2	7	12	13	25	20	23	23	28	25	14	0	193
1969	0	0	0	1	0	1	1	0	2	2	2	3	2	5	7	2	2	3	0	33
1970	0	0	0	0	1	1	0	0	1	1	1	4	4	4	3	7	4	3	0	34
1971	0	0	0	1	1	0	0	0	0	3	5	5	3	2	5	2	3	3	0	33
1972	0	0	0	0	0	0	0	1	2	5	5	4	6	3	9	3	7	1	0	46
1973	0	0	1	0	0	0	0	0	1	3	5	3	8	10	10	3	0	3	0	47
1969-73	0	0	1	2	2	2	1	1	6	14	18	19	23	24	34	17	16	13	0	193
1974	0	0	0	0	0	1	0	0	0	0	3	6	11	6	5	3	4	2	0	41
1975	0	0	1	0	0	0	0	1	0	2	7	6	6	8	7	2	2	5	0	47
1976	0	0	1	0	0	0	0	0	0	3	8	9	8	2	6	7	6	8	0	58
1977	0	0	0	0	1	0	1	3	2	1	9	6	8	7	6	7	1	0	0	52
1978	0	0	0	0	1	0	0	2	0	3	7	8	11	8	4	5	2	3	0	54
1974-78	0	0	2	0	2	1	1	6	2	9	34	35	44	31	28	24	15	18	0	252
1979	0	0	0	0	1	0	1	1	0	4	5	7	6	11	3	6	2	4	0	51
1980	0	1	0	1	1	0	1	0	3	3	6	6	12	12	4	5	2	5	0	62
1981	0	0	0	0	0	1	0	2	1	4	6	8	9	7	9	4	3	2	2	58
1982	0	0	0	0	0	0	0	0	4	5	3	9	8	8	1	5	2	4	0	49
1983	0	0	0	0	0	0	1	1	3	3	7	7	8	6	4	5	6	3	1	55
1979-83	0	1	0	1	2	1	3	4	11	19	27	37	43	44	21	25	15	18	3	275

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF MOUTH (ICD9 145)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.5	2.9	6.4	6.1	4.4	5.9	19.8	23.6	29.4	1.5	1.4	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	1.6	0.0	0.7	1.7	2.2	7.2	10.9	15.4	28.8	0.9	0.8	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.0	2.8	6.1	3.3	3.2	7.1	6.4	7.7	21.4	1.1	1.1	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.5	0.5	2.0	0.8	5.3	5.6	10.5	22.7	6.9	0.8	0.7	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	2.4	1.1	1.9	4.8	9.3	7.0	12.6	26.1	13.3	1.3	1.3	0.1
1964-68	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.2	0.6	1.2	1.5	3.4	3.3	4.9	6.6	12.0	19.2	19.7	1.1	1.1	0.1
1969	0.0	0.0	0.0	0.3	0.0	0.4	0.4	0.0	0.8	0.9	1.1	1.8	1.6	5.0	9.7	4.2	7.4	19.1	0.9	0.9	0.1
1970	0.0	0.0	0.0	0.0	0.3	0.4	0.0	0.0	0.4	0.4	0.5	2.4	3.0	3.9	4.1	14.4	14.4	18.3	0.9	0.9	0.1
1971	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.0	0.0	1.3	2.6	2.9	2.2	1.9	6.7	4.1	10.6	17.3	0.9	0.8	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	2.1	2.5	2.3	4.3	2.8	11.7	6.1	24.4	5.6	1.2	1.1	0.1
1973	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.4	1.3	2.4	1.8	5.5	9.1	12.6	6.0	0.0	16.5	1.2	1.2	0.2
1969-73	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.5	1.2	1.9	2.3	3.4	4.6	9.1	6.9	11.3	15.2	1.0	1.0	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	1.4	3.6	7.3	5.3	6.1	5.9	13.7	10.8	1.0	1.0	0.1
1975	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.4	0.0	0.8	3.1	3.5	3.9	6.9	8.3	3.8	6.8	26.3	1.1	1.1	0.1
1976	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	1.3	3.5	5.0	5.1	1.7	6.9	12.9	20.2	41.2	1.4	1.3	0.1
1977	0.0	0.0	0.0	0.0	0.3	0.0	0.3	1.2	0.8	0.4	4.0	3.2	5.1	5.6	6.8	12.4	3.3	0.0	1.2	1.2	0.1
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	0.0	1.2	3.1	4.0	7.0	6.1	4.4	8.5	6.5	15.0	1.2	1.2	0.1
1974-78	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.5	0.2	0.8	3.0	3.9	5.7	5.1	6.5	8.8	10.1	18.6	1.2	1.2	0.1
1979	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.4	0.0	1.7	2.2	3.4	3.8	8.2	3.2	9.8	6.3	19.5	1.1	1.1	0.1
1980	0.0	0.3	0.0	0.2	0.3	0.0	0.3	0.0	1.2	1.3	2.6	2.8	7.4	8.6	4.1	7.9	6.0	24.2	1.3	1.4	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.7	0.4	1.7	2.5	3.7	5.3	5.0	8.9	6.2	8.8	9.4	1.2	1.2	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	2.1	1.3	4.1	4.5	5.6	1.0	7.5	5.6	18.5	1.0	1.0	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	1.1	1.3	3.0	3.2	4.3	4.2	3.7	7.3	15.9	13.5	1.1	1.0	0.1
1979-83	0.0	0.1	0.0	0.0	0.1	0.1	0.2	0.3	0.9	1.6	2.3	3.4	5.0	6.3	4.2	7.7	8.7	16.9	1.1	1.1	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF MOUTH (ICD9 145)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	1	1	0	2	0	4	3	3	1	4	4	4	2	0	29
1965	0	0	0	0	1	1	0	1	2	0	2	2	2	3	2	2	2	0	0	20
1966	0	0	0	0	0	0	1	0	0	1	0	1	3	3	0	2	0	3	0	14
1967	0	0	0	0	0	1	0	1	0	1	4	1	0	1	2	0	3	0	0	14
1968	0	0	0	0	1	0	0	0	0	0	0	1	0	0	3	1	1	0	0	7
1964-68	0	0	0	0	2	3	2	2	4	2	10	8	8	8	11	9	10	5	0	84
1969	0	0	0	1	0	0	1	1	0	1	4	0	0	3	3	4	2	0	0	20
1970	0	0	0	0	0	0	0	0	0	0	6	1	5	3	2	2	2	1	0	22
1971	0	0	0	0	0	0	1	2	0	4	0	4	1	1	2	3	3	1	0	22
1972	0	0	0	0	0	0	0	0	2	0	3	1	2	4	2	3	4	1	0	22
1973	0	0	0	0	0	1	0	1	0	0	2	4	3	2	0	4	5	4	0	26
1969-73	0	0	0	1	0	1	2	4	2	5	15	10	11	13	9	16	16	7	0	112
1974	0	0	0	0	0	1	0	0	1	4	3	3	2	6	2	4	0	2	0	28
1975	0	0	0	0	0	0	1	0	0	3	0	3	3	6	2	2	3	2	0	25
1976	0	0	0	0	0	0	0	3	1	3	3	5	3	8	4	1	1	0	0	32
1977	0	0	1	0	0	2	0	0	2	2	9	3	5	7	4	3	3	1	1	43
1978	0	0	0	1	0	0	0	0	0	5	4	5	3	3	7	3	2	2	0	35
1974-78	0	0	1	1	0	3	1	3	4	17	19	19	16	30	19	13	9	7	1	163
1979	0	0	0	0	0	0	0	1	2	1	4	3	6	5	6	5	1	3	0	37
1980	0	0	0	0	1	0	0	1	0	2	1	5	5	4	1	0	4	0	0	24
1981	0	0	0	0	0	0	2	1	2	5	5	2	6	4	3	2	2	3	1	38
1982	0	0	1	1	0	1	1	1	1	0	4	3	8	8	6	2	0	4	0	41
1983	0	0	0	0	0	2	0	1	1	0	7	4	5	7	5	4	2	6	1	45
1979-83	0	0	1	1	1	3	3	5	6	8	21	17	30	28	21	13	9	16	2	185

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF MOUTH (ICD9 145)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.5	0.5	0.0	0.9	0.0	2.4	2.2	2.5	1.0	4.8	6.8	11.6	9.1	0.9	0.7	0.1
1965	0.0	0.0	0.0	0.0	0.4	0.5	0.0	0.4	0.9	0.0	1.2	1.4	1.7	2.9	2.3	3.3	5.6	0.0	0.6	0.5	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.5	0.0	0.7	2.4	2.8	0.0	3.2	0.0	12.7	0.4	0.4	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.0	0.5	2.2	0.7	0.0	0.9	2.2	0.0	7.7	0.0	0.4	0.3	0.0
1968	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	3.3	1.5	2.5	0.0	0.2	0.1	0.0
1964-68	0.0	0.0	0.0	0.0	0.2	0.3	0.2	0.2	0.3	0.2	1.2	1.1	1.3	1.5	2.5	2.9	5.3	4.2	0.5	0.4	0.0
1969	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	0.0	0.5	2.2	0.0	0.0	2.6	3.2	5.8	4.7	0.0	0.5	0.4	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.2	0.6	3.6	2.5	2.1	2.8	4.5	3.3	0.6	0.5	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.0	1.7	0.0	2.3	0.7	0.8	2.1	4.2	6.5	3.1	0.5	0.5	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	1.5	0.6	1.3	3.2	2.0	4.0	8.4	2.9	0.5	0.4	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.0	0.0	0.9	2.3	1.9	1.6	0.0	5.3	10.2	11.1	0.6	0.5	0.0
1969-73	0.0	0.0	0.0	0.1	0.0	0.1	0.2	0.3	0.2	0.4	1.5	1.2	1.5	2.1	1.9	4.4	7.0	4.3	0.6	0.5	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	1.7	1.3	1.7	1.3	4.5	1.9	5.1	0.0	5.3	0.6	0.6	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.3	0.0	1.7	1.8	4.4	1.9	2.5	5.8	5.1	0.6	0.5	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.4	1.3	1.3	2.6	1.8	5.7	3.6	1.2	1.9	0.0	0.7	0.7	0.1
1977	0.0	0.0	0.3	0.0	0.0	0.6	0.0	0.0	0.9	0.9	3.8	1.5	2.9	4.8	3.5	3.5	5.4	2.3	0.9	0.8	0.1
1978	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	2.1	1.7	2.3	1.7	2.0	5.9	3.4	3.5	4.5	0.7	0.7	0.1
1974-78	0.0	0.0	0.1	0.1	0.0	0.2	0.1	0.2	0.3	1.5	1.6	2.0	1.9	4.3	3.4	3.1	3.4	3.4	0.7	0.6	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	0.4	1.7	1.3	3.5	3.2	4.9	5.5	1.7	6.4	0.7	0.7	0.1
1980	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	0.9	0.4	2.2	2.8	2.5	0.8	0.0	6.5	0.0	0.5	0.4	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.3	0.8	2.2	2.1	0.9	3.2	2.4	2.3	2.1	3.1	5.8	0.7	0.7	0.1
1982	0.0	0.0	0.3	0.3	0.0	0.3	0.3	0.3	0.4	0.0	1.7	1.3	4.0	4.8	4.5	2.0	0.0	7.4	0.8	0.7	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.3	0.4	0.0	3.0	1.7	2.4	4.2	3.6	3.9	2.9	10.5	0.8	0.7	0.1
1979-83	0.0	0.0	0.1	0.1	0.1	0.2	0.2	0.3	0.5	0.7	1.8	1.5	3.2	3.4	3.2	2.7	2.8	6.2	0.7	0.6	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OROPHARYNX (ICD9 146)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	1	6	8	4	5	3	4	1	2	0	34
1965	0	0	0	0	1	1	0	1	2	4	1	5	8	2	6	4	4	3	0	42
1966	0	0	0	0	0	0	0	0	0	1	4	3	9	4	3	6	2	0	0	32
1967	0	0	0	0	0	0	0	3	2	2	4	3	6	6	2	3	2	2	0	35
1968	0	0	0	1	1	0	0	0	5	5	4	12	4	3	5	4	3	2	0	49
1964-68	0	0	0	1	2	1	0	4	9	13	19	31	31	20	19	21	12	9	0	192
1969	0	0	0	0	1	0	0	3	1	6	7	8	13	10	3	2	2	0	1	57
1970	0	0	0	0	0	0	0	0	1	5	9	14	15	6	4	3	1	1	0	59
1971	0	0	0	0	0	0	1	1	2	9	7	11	10	10	3	3	1	1	0	59
1972	0	0	0	0	0	0	0	0	4	7	10	8	8	12	6	8	3	0	0	66
1973	0	0	0	0	1	0	0	2	0	6	11	9	11	10	8	6	3	2	0	69
1969-73	0	0	0	0	2	0	1	6	8	33	44	50	57	48	24	22	10	4	1	310
1974	1	0	0	0	0	0	0	1	2	4	7	9	9	10	12	2	4	1	0	62
1975	0	0	0	0	0	1	0	1	5	3	7	7	11	5	13	4	2	0	0	59
1976	0	0	0	0	0	0	0	1	4	6	13	14	16	8	9	5	3	0	0	79
1977	0	0	0	0	0	0	0	1	0	11	13	16	11	7	12	6	2	2	0	81
1978	0	0	0	0	0	0	0	1	3	10	14	19	8	7	8	7	1	0	0	78
1974-78	1	0	0	0	0	1	0	5	14	34	54	65	55	37	54	24	12	3	0	359
1979	0	0	0	0	0	0	0	4	2	5	4	14	9	6	7	6	2	3	0	62
1980	0	0	0	0	0	0	0	1	3	11	6	12	12	8	5	1	4	0	0	63
1981	0	1	0	0	0	1	1	1	1	4	7	19	7	12	4	1	2	2	0	63
1982	0	0	0	0	0	0	0	0	3	4	7	13	9	13	8	5	1	2	0	65
1983	0	0	0	0	1	0	0	1	1	8	7	11	16	7	5	1	0	0	0	58
1979-83	0	1	0	0	1	1	1	7	10	32	31	69	53	46	29	14	9	7	0	311

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OROPHARYNX (ICD9 146)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	3.5	5.7	3.5	5.6	4.4	8.8	3.9	14.7	1.0	1.0	0.1
1965	0.0	0.0	0.0	0.0	0.4	0.5	0.0	0.4	0.9	2.1	0.6	3.5	6.9	2.2	8.6	8.7	15.4	21.6	1.2	1.2	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	2.2	2.0	7.5	4.3	4.3	12.9	7.7	0.0	0.9	0.9	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.8	1.0	2.2	2.0	4.9	6.3	2.8	6.3	7.6	13.8	1.0	1.0	0.1
1968	0.0	0.0	0.0	0.3	0.4	0.0	0.0	0.0	2.1	2.4	2.2	7.6	3.2	3.1	7.0	8.4	11.2	13.3	1.4	1.3	0.1
1964-68	0.0	0.0	0.0	0.1	0.2	0.1	0.0	0.3	0.8	1.3	2.1	4.2	5.2	4.3	5.4	9.0	9.2	12.7	1.1	1.1	0.1
1969	0.0	0.0	0.0	0.0	0.3	0.0	0.0	1.3	0.4	2.8	3.8	4.9	10.1	10.0	4.2	4.2	7.4	0.0	1.5	1.5	0.2
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.2	4.9	8.4	11.3	5.8	5.5	6.2	3.6	6.1	1.6	1.6	0.2
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.8	3.9	3.7	6.5	7.3	9.4	4.0	6.1	3.5	5.8	1.5	1.5	0.2
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	3.0	5.0	4.7	5.7	11.1	7.8	16.2	10.5	0.0	1.7	1.6	0.2
1973	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	0.0	2.6	5.3	5.3	7.6	9.1	10.1	12.0	10.3	11.0	1.7	1.7	0.2
1969-73	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.5	0.7	2.9	4.6	6.0	8.3	9.1	6.4	9.0	7.1	4.7	1.6	1.6	0.2
1974	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.7	3.2	5.3	6.0	8.9	14.6	4.0	13.7	5.4	1.5	1.5	0.2
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	2.1	1.3	3.1	4.1	7.1	4.3	15.4	7.6	6.8	0.0	1.4	1.4	0.2
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	2.5	5.7	7.8	10.2	6.6	10.4	9.2	10.1	0.0	1.8	1.8	0.2
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	4.6	5.7	8.5	7.0	5.6	13.6	10.6	6.6	10.2	1.8	1.8	0.2
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	4.1	6.1	9.5	5.1	5.4	8.8	11.9	3.2	0.0	1.7	1.7	0.2
1974-78	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.4	1.2	2.8	4.8	7.2	7.1	6.1	12.5	8.8	8.0	3.1	1.7	1.7	0.2
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.5	0.8	2.1	1.7	6.7	5.7	4.5	7.5	9.8	6.3	14.6	1.4	1.3	0.2
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.2	4.6	2.6	5.6	7.4	5.8	5.2	1.6	12.1	0.0	1.3	1.4	0.2
1981	0.0	0.3	0.0	0.0	0.0	0.3	0.3	0.3	0.4	1.7	3.0	8.8	4.1	8.5	4.0	1.5	5.9	9.4	1.3	1.3	0.2
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	1.7	3.0	6.0	5.1	9.1	7.6	7.5	2.8	9.3	1.3	1.3	0.2
1983	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.3	0.4	3.4	3.0	5.0	8.5	4.9	4.6	1.5	0.0	0.0	1.2	1.2	0.2
1979-83	0.0	0.1	0.0	0.0	0.1	0.1	0.1	0.5	0.8	2.7	2.6	6.4	6.2	6.6	5.7	4.3	5.2	6.6	1.3	1.3	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OROPHARYNX (ICD9 146)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	1	0	2	1	3	1	1	3	1	1	1	0	15
1965	0	0	0	0	0	0	0	0	0	0	1	1	2	1	1	0	0	0	0	6
1966	0	0	0	0	0	0	0	1	0	0	1	3	2	1	1	0	1	0	0	10
1967	0	0	0	0	0	0	0	0	1	2	2	4	4	2	1	1	0	1	0	18
1968	0	0	0	0	1	0	0	0	0	1	5	2	3	1	1	1	1	0	0	16
1964-68	0	0	0	0	1	0	0	2	1	5	10	13	12	6	7	3	3	2	0	65
1969	0	0	0	0	0	0	0	1	0	1	4	3	3	2	1	2	0	0	0	17
1970	0	0	0	0	0	0	0	1	1	6	6	1	6	3	4	0	0	1	0	29
1971	0	0	0	0	0	1	0	0	1	2	1	3	3	7	5	0	0	0	0	23
1972	0	0	0	0	0	0	0	2	0	3	4	6	4	2	3	0	1	1	0	26
1973	0	0	0	0	0	0	0	0	0	4	3	4	5	4	5	1	1	2	0	29
1969-73	0	0	0	0	0	1	0	4	2	16	18	17	21	18	18	3	2	4	0	124
1974	1	0	0	0	0	0	0	0	0	3	3	3	5	7	2	1	0	1	0	26
1975	0	0	0	0	0	0	0	0	0	3	3	7	1	2	0	1	1	0	0	18
1976	0	0	0	0	0	0	0	0	3	1	5	2	6	3	3	0	1	2	0	26
1977	0	0	0	0	0	0	0	1	0	2	2	9	5	11	2	1	0	0	0	33
1978	0	0	0	0	0	0	1	0	1	2	1	3	7	7	3	1	2	2	0	30
1974-78	1	0	0	0	0	0	1	1	4	11	14	24	24	30	10	4	4	5	0	133
1979	0	0	0	1	0	0	0	0	1	1	3	5	12	9	4	3	1	1	0	41
1980	0	0	0	0	0	0	0	0	0	0	2	5	4	6	0	0	3	1	0	21
1981	0	0	0	0	0	1	0	0	2	1	7	4	6	3	3	3	1	0	0	31
1982	0	0	0	0	0	0	0	0	1	2	5	6	7	3	0	3	2	1	0	30
1983	0	0	0	0	0	1	0	0	2	3	1	7	6	3	5	4	0	1	0	33
1979-83	0	0	0	1	0	2	0	0	6	7	18	27	35	24	12	13	7	4	0	156

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OROPHARYNX (ICD9 146)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.1	0.6	2.2	0.8	1.0	3.6	1.7	2.9	4.5	0.5	0.4	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	1.7	1.0	1.2	0.0	0.0	0.0	0.2	0.2	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.6	2.1	1.6	0.9	1.1	0.0	2.7	0.0	0.3	0.3	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.0	1.1	2.6	3.1	1.8	1.1	1.5	0.0	4.0	0.5	0.5	0.1
1968	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.5	2.8	1.3	2.3	0.9	1.1	1.5	2.5	0.0	0.4	0.4	0.0
1964-68	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.2	0.1	0.5	1.2	1.8	1.9	1.1	1.6	1.0	1.6	1.7	0.4	0.3	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	2.2	1.8	2.2	1.7	1.1	2.9	0.0	0.0	0.5	0.4	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.6	3.2	0.6	4.3	2.5	4.2	0.0	0.0	3.3	0.7	0.7	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.4	0.8	0.5	1.7	2.1	5.8	5.1	0.0	0.0	0.0	0.6	0.6	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	1.3	2.0	3.4	2.7	1.6	3.0	0.0	2.1	2.9	0.6	0.6	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	1.4	2.3	3.2	3.1	4.9	1.3	2.0	5.6	0.7	0.6	0.1
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.3	0.2	1.4	1.8	2.0	2.9	3.0	3.7	0.8	0.9	2.5	0.6	0.6	0.1
1974	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	1.3	1.7	3.1	5.3	1.9	1.3	0.0	2.6	0.6	0.6	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	1.3	3.9	0.6	1.5	0.0	1.2	1.9	0.0	0.4	0.4	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.4	2.1	1.1	3.6	2.1	2.7	0.0	1.9	4.8	0.6	0.5	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	0.8	4.5	2.9	7.6	1.7	1.2	0.0	0.0	0.7	0.7	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.9	0.4	1.4	4.1	4.7	2.5	1.1	3.5	4.5	0.6	0.6	0.1
1974-78	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.3	0.9	1.2	2.5	2.9	4.3	1.8	1.0	1.5	2.4	0.6	0.6	0.1
1979	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.4	0.4	1.3	2.2	6.9	5.8	3.3	3.3	1.7	2.1	0.8	0.8	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	2.2	2.2	3.7	0.0	0.0	4.9	2.0	0.4	0.4	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	0.4	3.0	1.7	3.2	1.8	2.3	3.1	1.6	0.0	0.6	0.6	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	2.1	2.6	3.5	1.8	0.0	3.0	3.1	1.9	0.6	0.5	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	1.3	0.4	3.0	2.9	1.8	3.6	3.9	0.0	1.8	0.6	0.6	0.1
1979-83	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.5	0.6	1.5	2.3	3.7	2.9	1.8	2.7	2.2	1.5	0.6	0.6	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF NASOPHARYNX (ICD9 147)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	4	1	1	6	4	2	2	2	1	1	0	0	24
1965	0	0	1	0	0	0	3	2	1	1	4	2	4	1	2	3	0	1	0	25
1966	0	0	1	1	0	0	1	1	0	0	2	0	4	1	0	0	0	1	0	12
1967	0	0	0	0	0	1	1	1	0	4	2	2	1	1	0	0	0	0	0	13
1968	0	1	0	0	0	1	1	1	2	1	2	1	2	3	1	1	1	0	0	18
1964-68	0	1	2	1	0	2	6	9	4	7	16	9	13	8	5	5	2	2	0	92
1969	0	1	2	2	0	1	1	2	2	3	3	3	2	3	1	0	1	0	0	27
1970	0	0	2	0	2	1	1	3	4	4	4	3	2	1	0	1	1	0	0	29
1971	0	0	1	0	1	2	0	2	6	6	3	1	2	2	3	0	2	1	0	32
1972	0	0	1	1	0	2	2	2	2	4	3	6	4	1	1	0	1	0	0	30
1973	1	0	0	1	1	0	2	0	5	5	1	1	0	3	0	0	1	0	0	21
1969-73	1	1	6	4	4	6	6	9	19	22	14	14	10	10	5	1	6	1	0	139
1974	0	0	0	0	1	0	1	2	4	2	4	4	2	2	2	2	0	0	0	26
1975	0	0	0	3	1	2	1	5	3	4	4	2	4	4	2	0	0	0	0	35
1976	0	0	0	3	0	0	0	1	5	6	6	3	6	3	1	0	0	1	1	36
1977	1	1	0	0	2	1	2	2	3	1	2	6	3	4	7	0	1	0	0	36
1978	0	0	0	1	1	0	0	0	6	2	2	5	6	0	2	2	0	0	0	27
1974-78	1	1	0	7	5	3	4	10	21	15	18	20	21	13	14	4	1	1	1	160
1979	1	0	1	1	0	1	1	2	0	3	2	4	3	2	3	0	1	0	0	25
1980	0	0	0	0	3	0	3	3	4	3	0	4	3	3	2	0	0	0	0	28
1981	2	0	0	1	0	0	0	1	5	4	5	7	4	6	4	0	0	0	0	39
1982	1	0	0	1	1	0	3	0	7	1	5	5	5	5	2	1	1	0	0	38
1983	1	0	0	0	0	0	0	3	4	7	5	3	4	3	3	0	1	0	0	34
1979-83	5	0	1	3	4	1	7	9	20	18	17	23	19	19	14	1	3	0	0	164

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF NASOPHARYNX (ICD9 147)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	0.5	0.5	3.5	2.9	1.8	2.2	2.9	2.2	3.9	0.0	0.7	0.7	0.1
1965	0.0	0.0	0.3	0.0	0.0	0.0	1.3	0.8	0.4	0.5	2.3	1.4	3.4	1.1	2.9	6.5	0.0	7.2	0.7	0.7	0.1
1966	0.0	0.0	0.3	0.3	0.0	0.0	0.4	0.4	0.0	0.0	1.1	0.0	3.3	1.1	0.0	0.0	0.0	7.1	0.4	0.4	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.0	2.0	1.1	1.3	0.8	1.1	0.0	0.0	0.0	0.0	0.4	0.4	0.0
1968	0.0	0.2	0.0	0.0	0.0	0.4	0.4	0.4	0.8	0.5	1.1	0.6	1.6	3.1	1.4	2.1	3.7	0.0	0.5	0.5	0.1
1964-68	0.0	0.1	0.1	0.1	0.0	0.2	0.5	0.8	0.3	0.7	1.8	1.2	2.2	1.7	1.4	2.1	1.5	2.8	0.5	0.5	0.1
1969	0.0	0.2	0.5	0.6	0.0	0.4	0.4	0.8	0.8	1.4	1.6	1.8	1.6	3.0	1.4	0.0	3.7	0.0	0.7	0.7	0.1
1970	0.0	0.0	0.5	0.0	0.6	0.4	0.4	1.3	1.7	1.8	2.2	1.8	1.5	1.0	0.0	2.1	3.6	0.0	0.8	0.7	0.1
1971	0.0	0.0	0.2	0.0	0.3	0.7	0.0	0.8	2.5	2.6	1.6	0.6	1.5	1.9	4.0	0.0	7.1	5.8	0.8	0.8	0.1
1972	0.0	0.0	0.2	0.3	0.0	0.7	0.8	0.8	0.8	1.7	1.5	3.5	2.8	0.9	1.3	0.0	3.5	0.0	0.8	0.7	0.1
1973	0.3	0.0	0.0	0.3	0.3	0.0	0.8	0.0	2.1	2.1	0.5	0.6	0.0	2.7	0.0	0.0	3.4	0.0	0.5	0.5	0.0
1969-73	0.1	0.1	0.3	0.2	0.2	0.4	0.5	0.8	1.6	1.9	1.5	1.7	1.5	1.9	1.3	0.4	4.3	1.2	0.7	0.7	0.1
1974	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.8	1.6	0.8	1.8	2.4	1.3	1.8	2.4	4.0	0.0	0.0	0.6	0.6	0.1
1975	0.0	0.0	0.0	0.7	0.3	0.6	0.3	2.1	1.2	1.7	1.8	1.2	2.6	3.4	2.4	0.0	0.0	0.0	0.8	0.8	0.1
1976	0.0	0.0	0.0	0.7	0.0	0.0	0.0	0.4	2.1	2.5	2.6	1.7	3.8	2.5	1.2	0.0	0.0	5.2	0.8	0.8	0.1
1977	0.3	0.3	0.0	0.0	0.5	0.3	0.6	0.8	1.3	0.4	0.9	3.2	1.9	3.2	7.9	0.0	3.3	0.0	0.8	0.8	0.1
1978	0.0	0.0	0.0	0.2	0.3	0.0	0.0	0.0	2.5	0.8	0.9	2.5	3.8	0.0	2.2	3.4	0.0	0.0	0.6	0.6	0.1
1974-78	0.1	0.1	0.0	0.3	0.3	0.2	0.3	0.8	1.7	1.3	1.6	2.2	2.7	2.1	3.2	1.5	0.7	1.0	0.7	0.8	0.1
1979	0.3	0.0	0.3	0.2	0.0	0.3	0.3	0.7	0.0	1.2	0.9	1.9	1.9	1.5	3.2	0.0	3.1	0.0	0.6	0.6	0.1
1980	0.0	0.0	0.0	0.0	0.8	0.0	0.9	1.1	1.6	1.3	0.0	1.9	1.9	2.2	2.1	0.0	0.0	0.0	0.6	0.6	0.1
1981	0.7	0.0	0.0	0.2	0.0	0.0	0.0	0.3	2.0	1.7	2.1	3.2	2.4	4.3	4.0	0.0	0.0	0.0	0.8	0.9	0.1
1982	0.3	0.0	0.0	0.2	0.2	0.0	0.9	0.0	2.8	0.4	2.1	2.3	2.8	3.5	1.9	1.5	2.8	0.0	0.8	0.8	0.1
1983	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.5	3.0	2.1	1.4	2.1	2.1	2.7	0.0	2.6	0.0	0.7	0.7	0.1
1979-83	0.3	0.0	0.1	0.1	0.2	0.1	0.4	0.6	1.6	1.5	1.4	2.1	2.2	2.7	2.8	0.3	1.7	0.0	0.7	0.7	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF NASOPHARYNX (ICD9 147)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	0	0	1	0	0	1	0	0	0	0	1	0	2	0	0	1	0	0	7
1965	0	1	0	0	0	0	0	0	1	0	0	3	2	1	0	1	0	1	0	10
1966	0	0	1	1	0	0	1	0	0	2	2	3	1	3	2	1	0	0	0	17
1967	0	0	0	0	0	1	1	0	0	1	2	1	1	1	1	0	0	0	0	9
1968	0	0	0	1	0	0	0	0	0	0	1	3	1	0	1	0	2	0	0	9
1964-68	1	1	1	3	0	1	3	0	1	3	5	11	5	7	4	2	3	1	0	52
1969	0	2	1	0	1	0	2	3	1	0	1	1	0	1	3	3	1	0	0	20
1970	0	0	0	1	0	0	0	0	1	1	1	1	1	0	1	0	0	0	0	7
1971	0	0	0	0	0	1	0	0	0	1	0	0	1	2	0	2	1	0	0	8
1972	0	0	0	1	0	0	1	1	1	1	1	0	4	0	2	2	0	0	0	14
1973	0	0	0	0	0	0	0	0	1	4	2	0	2	0	3	2	0	0	0	14
1969-73	0	2	1	2	1	1	3	4	4	7	5	2	8	3	9	9	2	0	0	63
1974	0	0	0	1	1	0	0	1	0	1	0	0	1	2	0	1	0	1	0	9
1975	0	0	1	0	1	0	0	0	0	0	1	2	0	2	1	1	2	0	0	11
1976	1	0	1	0	1	0	1	1	4	1	1	0	4	0	0	1	2	0	0	18
1977	1	0	0	0	0	3	0	2	2	1	4	2	1	2	1	0	0	1	0	20
1978	0	0	0	0	0	0	1	0	2	1	1	0	2	1	1	0	0	0	0	9
1974-78	2	0	2	1	3	3	2	4	8	4	7	4	8	7	3	3	4	2	0	67
1979	1	0	0	0	0	0	0	0	3	1	3	2	1	0	0	0	0	0	0	11
1980	0	1	0	0	1	3	0	1	3	0	4	0	1	2	3	2	1	1	0	21
1981	1	0	0	0	0	1	1	2	0	0	1	3	4	4	1	2	0	1	0	21
1982	0	0	0	1	0	1	1	2	4	1	5	2	0	2	0	4	0	0	0	23
1983	0	1	0	2	0	0	1	0	1	2	4	1	1	1	1	1	0	0	0	16
1979-83	2	2	0	3	1	5	3	5	11	4	17	8	7	7	5	9	1	2	0	92

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF NASOPHARYNX (ICD9 147)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.0	0.0	0.4	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.7	0.0	2.0	0.0	0.0	2.9	0.0	0.2	0.2	0.0
1965	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	2.1	1.7	1.0	0.0	1.7	0.0	4.3	0.3	0.3	0.0
1966	0.0	0.0	0.3	0.3	0.0	0.0	0.5	0.0	0.0	1.0	1.1	2.1	0.8	2.8	2.3	1.6	0.0	0.0	0.5	0.5	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.0	0.5	1.1	0.7	0.8	0.9	1.1	0.0	0.0	0.0	0.3	0.3	0.0
1968	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.6	1.9	0.8	0.0	1.1	0.0	4.9	0.0	0.2	0.2	0.0
1964-68	0.1	0.1	0.1	0.2	0.0	0.1	0.3	0.0	0.1	0.3	0.6	1.5	0.8	1.3	0.9	0.6	1.6	0.8	0.3	0.3	0.0
1969	0.0	0.5	0.3	0.0	0.3	0.0	0.9	1.3	0.4	0.0	0.5	0.6	0.0	0.9	3.2	4.4	2.4	0.0	0.5	0.5	0.0
1970	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.6	0.7	0.0	1.1	0.0	0.0	0.0	0.2	0.2	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.4	0.0	0.0	0.7	1.6	0.0	2.8	2.2	0.0	0.2	0.2	0.0
1972	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	0.4	0.4	0.5	0.0	2.7	0.0	2.0	2.7	0.0	0.0	0.3	0.3	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	0.9	0.0	1.3	0.0	2.9	2.6	0.0	0.0	0.3	0.3	0.0
1969-73	0.0	0.1	0.1	0.1	0.1	0.1	0.3	0.3	0.3	0.6	0.5	0.2	1.1	0.5	1.9	2.5	0.9	0.0	0.3	0.3	0.0
1974	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.4	0.0	0.4	0.0	0.0	0.6	1.5	0.0	1.3	0.0	2.6	0.2	0.2	0.0
1975	0.0	0.0	0.3	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.4	1.1	0.0	1.5	0.9	1.2	3.9	0.0	0.3	0.2	0.0
1976	0.3	0.0	0.3	0.0	0.3	0.0	0.3	0.4	1.7	0.4	0.4	0.0	2.4	0.0	0.0	1.2	3.8	0.0	0.4	0.4	0.0
1977	0.3	0.0	0.0	0.0	0.0	0.8	0.0	0.8	0.9	0.4	1.7	1.0	0.6	1.4	0.9	0.0	0.0	2.3	0.4	0.5	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.9	0.4	0.4	0.0	1.2	0.7	0.8	0.0	0.0	0.0	0.2	0.2	0.0
1974-78	0.1	0.0	0.1	0.1	0.2	0.2	0.1	0.3	0.7	0.3	0.6	0.4	1.0	1.0	0.5	0.7	1.5	1.0	0.3	0.3	0.0
1979	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.4	1.3	0.9	0.6	0.0	0.0	0.0	0.0	0.0	0.2	0.3	0.0
1980	0.0	0.3	0.0	0.0	0.3	0.8	0.0	0.4	1.3	0.0	1.7	0.0	0.6	0.0	2.4	2.1	1.6	2.0	0.4	0.4	0.0
1981	0.3	0.0	0.0	0.0	0.0	0.3	0.3	0.7	0.0	0.0	0.4	1.3	2.1	2.4	0.8	2.1	0.0	1.9	0.4	0.4	0.0
1982	0.0	0.0	0.0	0.3	0.0	0.3	0.3	0.6	1.6	0.4	2.1	0.9	0.0	1.2	0.0	4.0	0.0	0.0	0.5	0.4	0.0
1983	0.0	0.3	0.0	0.5	0.0	0.0	0.3	0.0	0.4	0.9	1.7	0.4	0.5	0.6	0.7	1.0	0.0	0.0	0.3	0.3	0.0
1979-83	0.1	0.1	0.0	0.2	0.1	0.3	0.2	0.3	0.9	0.3	1.5	0.7	0.7	0.9	0.8	1.9	0.3	0.8	0.4	0.4	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF HYPOPHARYNX (ICD9 148)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	0	1	3	7	2	3	3	3	2	0	24
1965	0	0	0	0	0	0	0	0	1	0	3	6	3	1	2	2	3	1	0	22
1966	0	0	0	0	0	0	0	0	0	0	3	3	6	6	3	2	1	2	0	26
1967	0	0	0	0	0	0	0	1	0	1	6	4	5	3	7	2	2	0	0	31
1968	0	0	0	0	0	0	0	2	2	3	2	5	7	5	5	6	1	0	0	38
1964-68	0	0	0	0	0	0	0	3	3	4	15	21	28	17	20	15	10	5	0	141
1969	0	0	0	0	0	0	0	0	1	0	6	6	5	5	2	4	1	1	0	31
1970	0	0	0	0	0	0	0	0	1	4	2	8	7	2	5	3	1	0	0	33
1971	0	0	0	0	0	0	0	0	1	3	7	6	6	8	0	5	2	0	0	38
1972	0	0	0	0	0	0	0	0	1	2	2	5	5	6	7	3	1	0	0	32
1973	0	0	0	0	0	0	0	1	2	4	3	1	12	5	8	4	4	1	0	45
1969-73	0	0	0	0	0	0	0	1	6	13	20	26	35	26	22	19	9	2	0	179
1974	0	0	0	0	0	0	0	0	1	3	10	4	7	5	4	3	5	1	0	43
1975	0	0	0	0	0	0	0	2	2	5	7	14	6	12	7	4	1	0	0	60
1976	0	0	0	0	0	0	0	1	2	0	4	7	9	6	3	4	0	2	0	38
1977	0	0	0	0	0	0	0	0	1	9	7	13	9	7	6	2	1	0	0	57
1978	0	0	1	1	0	0	0	0	1	5	6	10	11	4	7	2	3	1	0	51
1974-78	0	0	0	0	0	1	0	0	1	5	6	10	11	4	7	2	3	1	0	51
1979	0	0	1	1	0	1	0	3	7	22	34	48	42	34	27	15	10	4	0	249
1980	0	0	0	0	0	0	0	0	2	9	7	9	8	5	10	3	2	4	0	59
1981	0	0	0	0	0	0	0	0	3	3	5	17	11	9	8	3	1	3	0	63
1982	0	0	0	0	0	0	0	0	2	3	5	8	7	13	2	2	0	0	0	43
1983	0	0	0	0	0	0	0	1	1	7	13	8	9	11	5	4	1	0	0	61
1979-83	0	0	0	0	0	1	0	1	1	7	13	8	9	11	5	4	1	0	0	61
	0	0	0	0	0	1	0	1	10	17	36	58	45	50	39	21	9	11	0	298

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF HYPOPHARYNX (ICD9 148)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	2.1	6.1	2.2	4.4	6.6	11.8	14.7	0.7	0.7	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.7	4.2	2.6	1.1	2.9	4.4	11.6	7.2	0.7	0.6	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	2.0	5.0	6.4	4.3	4.3	3.8	14.3	0.8	0.8	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	3.3	2.6	4.1	3.2	9.9	4.2	7.6	0.0	0.9	0.9	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.8	1.4	1.1	3.2	5.6	5.1	7.0	12.6	3.7	0.0	1.0	1.0	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	1.7	2.8	4.7	3.6	5.7	6.4	7.7	7.0	0.8	0.8	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	3.3	3.7	3.9	5.0	2.8	8.3	3.7	6.4	0.8	0.8	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.8	1.1	4.8	5.3	1.9	6.9	6.2	3.6	0.0	0.9	0.9	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	3.7	3.5	4.4	7.5	0.0	10.2	7.1	0.0	1.0	1.0	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	1.0	2.9	3.5	5.6	9.1	6.1	3.5	0.0	0.8	0.8	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.7	1.4	0.6	8.2	4.5	10.1	8.0	13.7	5.5	1.1	1.1	0.1
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.5	1.1	2.1	3.1	5.1	4.9	5.9	7.8	6.4	2.3	0.9	0.9	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	4.6	2.4	4.7	4.4	4.9	5.9	17.1	5.4	1.1	1.0	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.8	2.1	3.1	8.1	3.9	10.3	8.3	7.6	3.4	0.0	1.4	1.4	0.2
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	0.0	1.8	3.9	5.7	5.0	3.5	7.4	0.0	10.3	0.9	0.9	0.1
1977	0.0	0.0	0.3	0.2	0.0	0.0	0.0	0.0	0.4	3.7	3.1	6.9	5.7	5.6	6.8	3.5	3.3	0.0	1.3	1.3	0.2
1978	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	2.1	2.6	5.0	7.0	3.1	7.7	3.4	9.7	5.0	1.1	1.1	0.1
1974-78	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.6	1.8	3.0	5.3	5.4	5.6	6.2	5.5	6.7	4.1	1.2	1.2	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	3.7	3.0	4.3	5.1	3.7	10.7	4.9	6.3	19.5	1.3	1.3	0.2
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	1.3	2.1	8.0	6.8	6.5	8.2	4.8	3.0	14.5	1.3	1.3	0.2
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	1.3	2.1	3.7	4.1	9.2	2.0	3.1	0.0	0.0	0.9	0.9	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	3.0	6.0	4.5	6.3	10.5	7.5	11.2	4.6	1.2	1.2	0.2
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	5.1	5.0	5.9	9.8	7.3	11.7	5.3	13.5	1.4	1.4	0.2
1979-83	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.8	1.4	3.1	5.4	5.3	7.1	7.7	6.5	5.2	10.3	1.2	1.2	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF HYPOPHARYNX (ICD9 148)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	1	1	0	0	0	3	4	1	1	0	0	11
1965	0	0	0	0	0	0	0	0	1	1	1	1	0	2	1	1	0	0	0	8
1966	0	0	0	0	0	0	0	0	0	0	0	2	2	1	1	2	0	0	0	8
1967	0	0	0	0	0	0	0	0	0	0	2	0	2	1	1	0	0	1	0	7
1968	0	0	0	0	0	0	0	0	1	0	2	2	2	2	3	2	0	0	0	14
1964-68	0	0	0	0	0	0	0	0	3	2	5	5	6	9	10	6	1	1	0	48
1969	0	0	0	0	0	0	0	1	0	2	2	1	1	3	2	1	2	1	0	16
1970	0	0	0	0	0	0	0	0	1	1	2	2	2	1	2	2	2	0	0	15
1971	0	0	0	0	0	0	0	1	0	1	1	1	2	1	3	1	2	1	0	14
1972	0	0	0	0	0	0	0	0	0	1	3	4	4	3	1	0	0	0	0	16
1973	0	0	0	0	0	0	0	0	3	0	2	0	1	5	2	1	1	1	0	16
1969-73	0	0	0	0	0	0	0	2	4	5	10	8	10	13	10	5	7	3	0	77
1974	0	0	0	0	0	0	0	0	0	0	2	2	1	3	3	1	0	2	0	14
1975	0	0	0	0	0	0	0	0	0	1	0	4	4	1	2	1	0	1	1	15
1976	0	0	0	0	0	0	0	0	0	0	2	1	2	1	1	0	3	0	0	10
1977	0	0	0	0	0	0	2	0	1	1	1	0	1	4	1	2	0	1	0	14
1978	0	0	0	0	0	0	0	0	1	0	2	2	4	0	4	2	0	0	0	15
1974-78	0	0	0	0	0	0	2	0	2	2	7	9	12	9	11	6	3	4	1	68
1979	0	0	0	0	0	0	0	0	2	1	1	2	4	6	2	1	1	2	0	22
1980	0	0	0	0	0	0	0	0	0	0	3	6	3	1	1	3	2	0	0	19
1981	0	0	0	0	0	1	0	0	0	0	2	1	1	2	4	2	1	2	0	16
1982	0	0	0	0	0	0	0	0	1	1	1	2	2	1	2	5	2	1	0	18
1983	0	0	0	0	0	0	0	0	1	1	2	6	1	2	4	1	1	1	0	20
1979-83	0	0	0	0	0	1	0	0	4	3	9	17	11	12	13	12	7	6	0	95

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF HYPOPHARYNX (ICD9 148)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.5	0.0	0.0	0.0	3.0	4.8	1.7	2.9	0.0	0.3	0.3	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.6	0.7	0.0	1.9	1.2	1.7	0.0	0.0	0.2	0.2	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	1.6	0.9	1.1	3.2	0.0	0.0	0.2	0.2	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	1.6	0.9	1.1	0.0	0.0	4.0	0.2	0.2	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.1	1.3	1.5	1.8	3.3	3.0	0.0	0.0	0.4	0.3	0.0
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.2	0.6	0.7	1.0	1.7	2.3	1.9	0.5	0.8	0.3	0.2	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	1.1	0.6	0.7	2.6	2.2	1.5	4.7	3.5	0.4	0.4	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.1	1.2	1.4	0.8	2.1	2.8	4.5	0.0	0.4	0.3	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.5	0.6	1.4	0.8	3.1	1.4	4.3	3.1	0.3	0.3	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.5	2.3	2.7	2.4	1.0	0.0	0.0	0.0	0.4	0.4	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.0	0.9	0.0	0.6	3.9	2.0	1.3	2.0	2.8	0.4	0.3	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.3	0.4	1.0	0.9	1.4	2.1	2.1	1.4	3.1	1.9	0.4	0.3	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.1	0.6	2.3	2.9	1.3	0.0	5.3	0.3	0.3	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	2.2	2.4	0.7	1.9	1.2	0.0	2.5	0.3	0.3	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.5	1.2	0.7	0.9	0.0	5.6	0.0	0.2	0.2	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.4	0.4	0.4	0.0	0.6	2.7	0.9	2.4	0.0	2.3	0.3	0.3	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.8	0.9	2.3	0.0	3.4	2.3	0.0	0.0	0.3	0.3	0.0
1974-78	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.2	0.6	0.9	1.4	1.3	2.0	1.5	1.1	1.9	0.3	0.3	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	0.4	0.9	2.3	3.9	1.6	1.1	1.7	4.2	0.4	0.4	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	2.6	1.7	0.6	0.8	3.2	3.2	0.0	0.4	0.3	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.9	0.4	0.5	1.2	3.1	2.1	1.6	3.9	0.3	0.2	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.9	1.0	0.6	1.5	5.0	3.1	1.9	0.3	0.3	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.9	2.6	0.5	1.2	2.9	1.0	1.5	1.8	0.4	0.3	0.0
1979-83	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.3	0.3	0.8	1.5	1.2	1.5	2.0	2.5	2.2	2.3	0.4	0.3	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE LIP, ORAL CAVITY, AND PHARYNX (ICD9 149)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	1	3	2	4	6	2	6	3	5	2	0	34
1965	0	0	0	1	0	0	1	0	0	0	1	11	7	4	7	2	5	3	0	42
1966	0	0	0	0	0	0	0	0	1	2	5	3	9	7	9	5	2	2	0	45
1967	0	0	0	0	0	0	0	1	0	2	4	2	9	3	3	4	2	2	0	32
1968	0	0	0	0	0	0	0	0	2	1	1	6	4	1	1	0	1	2	0	19
1964-68	0	0	0	1	0	0	1	1	4	8	13	26	35	17	26	14	15	11	0	172
1969	0	0	0	0	0	0	0	0	0	1	6	1	2	2	1	0	2	1	0	16
1970	0	0	0	0	0	0	0	0	0	0	1	2	3	4	2	0	0	0	0	12
1971	0	0	0	0	0	0	0	1	0	5	0	2	2	1	3	2	0	0	0	16
1972	0	0	0	0	0	0	0	0	1	0	1	1	4	2	0	1	2	2	0	14
1973	0	0	0	0	0	0	0	0	1	3	3	4	1	3	1	2	1	1	0	20
1969-73	0	0	0	0	0	0	0	1	2	9	11	10	12	12	7	5	5	4	0	78
1974	0	0	0	0	0	0	0	0	2	4	7	3	3	1	3	2	2	1	0	28
1975	0	0	0	0	0	0	0	0	0	2	3	3	3	6	1	2	1	1	0	22
1976	0	0	0	0	0	0	0	0	0	3	6	2	2	3	2	1	0	0	0	19
1977	0	0	0	0	0	0	0	1	1	1	2	3	3	3	1	0	1	0	2	18
1978	0	0	0	0	0	0	0	1	0	1	2	1	3	4	2	4	1	0	0	19
1974-78	0	0	0	0	0	0	0	2	3	11	20	12	14	17	9	9	5	2	2	106
1979	0	0	0	0	0	0	0	0	0	1	3	6	3	5	1	2	1	1	0	23
1980	0	0	0	0	0	0	0	0	0	2	5	3	2	1	3	2	4	0	0	22
1981	0	0	0	0	0	0	0	0	2	3	1	6	1	4	4	3	0	1	0	25
1982	0	0	0	0	0	0	1	0	2	1	2	6	7	3	3	1	1	1	1	29
1983	0	0	0	0	0	0	0	0	0	0	4	7	6	3	4	1	1	3	0	29
1979-83	0	0	0	0	0	0	1	0	4	7	15	28	19	16	15	9	7	6	1	128

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE LIP, ORAL CAVITY, AND PHARYNX (ICD9 149)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.6	1.2	2.9	5.3	2.2	8.8	6.6	19.7	14.7	1.0	1.0	0.1
1965	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	0.0	0.0	0.6	7.7	6.0	4.4	10.1	4.4	19.3	21.6	1.3	1.2	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.0	2.8	2.0	7.5	7.5	12.8	10.7	7.7	14.3	1.3	1.3	0.2
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.0	2.2	1.3	7.3	3.2	4.2	8.4	7.6	13.8	0.9	0.9	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.5	0.5	3.8	3.2	1.0	1.4	0.0	3.7	13.3	0.5	0.5	0.1
1964-68	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.1	0.3	0.8	1.5	3.5	5.8	3.6	7.4	6.0	11.5	15.5	1.0	1.0	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	3.3	0.6	1.6	2.0	1.4	0.0	7.4	6.4	0.4	0.4	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.2	2.3	3.9	2.7	0.0	0.0	0.0	0.3	0.3	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	2.2	0.0	1.2	1.5	0.9	4.0	4.1	0.0	0.0	0.4	0.4	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	0.6	2.8	1.9	0.0	2.0	7.0	11.2	0.4	0.4	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	1.4	2.4	0.7	2.7	1.3	4.0	3.4	5.5	0.5	0.5	0.1
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.8	1.1	1.2	1.8	2.3	1.9	2.0	3.5	4.7	0.4	0.4	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.7	3.2	1.8	2.0	0.9	3.6	4.0	6.8	5.4	0.7	0.7	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.3	1.7	1.9	5.2	1.2	3.8	3.4	5.3	0.5	0.5	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	2.6	1.1	1.3	2.5	2.3	1.8	0.0	0.0	0.4	0.4	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.9	1.6	1.9	2.4	1.1	0.0	3.3	0.0	0.4	0.4	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.9	0.5	1.9	3.1	2.2	6.8	3.2	0.0	0.4	0.4	0.0
1974-78	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.9	1.8	1.3	1.8	2.8	2.1	3.3	3.4	2.1	0.5	0.5	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	2.9	1.9	3.7	1.1	3.3	3.1	4.9	0.5	0.5	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	2.1	1.4	1.2	0.7	3.1	3.2	12.1	0.0	0.5	0.4	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.3	0.4	2.8	0.6	2.8	4.0	4.6	0.0	4.7	0.5	0.5	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.8	0.4	0.8	2.7	3.9	2.1	2.9	1.5	2.8	4.6	0.6	0.6	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	3.2	3.2	2.1	3.7	1.5	2.6	13.5	0.6	0.6	0.1
1979-83	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.3	0.6	1.3	2.6	2.2	2.3	3.0	2.8	4.1	5.6	0.5	0.5	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE LIP, ORAL CAVITY, AND PHARYNX (ICD9 149)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	1	0	0	0	1	0	0	0	1	1	2	4	1	1	0	0	0	12
1965	0	0	0	0	0	0	0	3	0	1	1	1	0	0	4	0	2	1	0	13
1966	0	0	0	0	0	0	0	0	0	0	3	1	1	3	3	2	2	0	0	15
1967	0	0	0	0	0	0	0	0	2	0	0	1	2	4	0	1	2	2	0	14
1968	0	0	0	0	0	0	0	0	0	1	2	0	2	1	0	1	0	0	0	7
1964-68	0	0	1	0	0	0	1	3	2	2	7	4	7	12	8	5	6	3	0	61
1969	0	0	0	0	0	0	0	0	0	0	2	0	0	0	1	0	0	0	0	3
1970	0	0	0	0	0	1	0	0	0	0	0	3	1	2	1	1	1	0	0	10
1971	0	0	0	0	0	0	0	0	0	0	2	2	0	2	0	1	0	0	0	7
1972	0	0	0	0	0	0	0	0	1	1	1	0	2	2	0	0	0	2	0	9
1973	0	0	0	0	0	0	0	0	0	0	0	1	1	3	1	1	3	0	0	10
1969-73	0	0	0	0	0	1	0	0	1	1	5	6	4	9	3	3	4	2	0	39
1974	0	0	1	0	0	0	0	0	0	1	1	2	2	0	2	0	0	1	0	10
1975	0	0	0	0	0	0	0	0	0	1	2	2	1	1	1	0	1	0	0	9
1976	0	0	0	0	0	0	0	1	0	2	0	1	0	2	0	2	0	0	0	8
1977	0	0	0	0	0	0	0	0	1	0	1	1	2	0	1	1	0	1	0	8
1978	0	0	0	0	1	0	1	0	0	1	2	0	1	2	1	1	1	1	0	12
1974-78	0	0	1	0	1	0	1	1	1	5	6	6	6	5	5	4	2	3	0	47
1979	0	0	0	0	0	0	0	0	1	1	0	1	2	2	1	0	0	0	2	10
1980	0	0	0	0	0	0	0	0	0	2	0	2	0	2	1	1	0	0	0	8
1981	0	0	0	0	0	0	0	1	0	0	2	1	3	2	0	1	0	0	0	10
1982	0	0	0	0	0	0	0	0	0	0	1	1	1	0	2	0	1	2	0	8
1983	0	0	0	0	0	0	0	0	0	0	2	2	1	2	1	1	1	2	0	12
1979-83	0	0	0	0	0	0	0	1	1	3	5	7	7	8	5	3	2	4	2	48

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE LIP, ORAL CAVITY, AND PHARYNX (ICD9 149)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.3	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.6	0.7	1.7	3.9	1.2	1.7	0.0	0.0	0.4	0.3	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.0	0.5	0.6	0.7	0.0	0.0	4.7	0.0	5.6	4.3	0.4	0.3	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	0.7	0.8	2.8	3.4	3.2	5.4	0.0	0.4	0.4	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.7	1.6	3.7	0.0	1.5	5.1	0.0	0.4	0.3	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.1	0.0	1.5	0.9	0.0	1.5	0.0	0.0	0.2	0.2	0.0
1964-68	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.3	0.2	0.2	0.8	0.5	1.1	2.3	1.8	1.6	3.2	2.5	0.3	0.3	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0	0.0	1.1	0.0	0.0	0.0	0.1	0.1	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	1.8	0.7	1.7	1.1	1.4	2.3	0.0	0.3	0.2	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	1.2	0.0	1.6	0.0	1.4	0.0	0.0	0.2	0.2	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.0	1.3	1.6	0.0	0.0	0.0	5.8	0.2	0.2	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.6	2.3	1.0	1.3	6.1	0.0	0.2	0.2	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.1	0.5	0.7	0.6	1.5	0.6	0.8	1.7	1.2	0.2	0.2	0.0
1974	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.1	1.3	0.0	1.9	0.0	0.0	2.6	0.2	0.2	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	1.1	0.6	0.7	0.9	0.0	1.9	0.0	0.2	0.2	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	0.0	0.5	0.0	1.4	0.0	2.4	0.0	0.0	0.2	0.2	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.5	1.2	0.0	0.9	1.2	0.0	2.3	0.2	0.2	0.0
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.0	0.0	0.4	0.8	0.0	0.6	1.3	0.8	1.1	1.7	2.2	0.3	0.2	0.0
1974-78	0.0	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.4	0.5	0.6	0.7	0.7	0.9	1.0	0.7	1.4	0.2	0.2	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.4	1.2	1.3	0.8	0.0	0.0	0.0	0.2	0.2	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.9	0.0	1.2	0.8	1.1	0.0	0.0	0.2	0.2	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.9	0.4	1.6	1.2	0.0	1.0	0.0	0.0	0.2	0.2	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.0	1.5	0.0	1.5	3.7	0.1	0.1	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.9	0.5	1.2	0.7	1.0	1.5	3.5	0.2	0.2	0.0
1979-83	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.3	0.4	0.6	0.7	1.0	0.8	0.6	0.6	1.5	0.2	0.2	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OESOPHAGUS (ICD9 150)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	0	1	4	9	14	9	16	34	27	16	16	7	0	153
1965	0	0	0	0	0	0	0	0	4	4	7	17	21	20	19	28	16	4	0	140
1966	0	0	0	0	0	0	0	1	8	3	17	19	21	23	25	21	13	11	0	162
1967	0	0	0	0	0	0	0	3	2	16	11	25	22	27	26	23	7	12	0	174
1968	0	0	0	0	0	0	0	0	3	6	15	19	31	28	19	17	12	6	0	156
1964-68	0	0	0	0	0	0	0	5	21	38	64	89	111	132	116	105	64	40	0	785
1969	0	0	0	0	0	0	0	1	7	6	13	22	22	21	16	14	12	11	0	145
1970	0	0	0	0	0	0	0	0	3	8	17	22	20	25	17	23	13	9	0	157
1971	0	0	0	0	0	0	2	0	1	9	14	18	27	22	20	16	15	10	0	154
1972	0	0	0	0	0	0	0	1	4	9	14	29	26	27	15	22	20	10	0	177
1973	0	0	0	0	0	1	0	1	4	19	16	29	32	24	22	25	14	6	0	193
1969-73	0	0	0	0	0	1	2	3	19	51	74	120	127	119	90	100	74	46	0	826
1974	0	0	0	0	0	0	0	1	5	8	17	23	31	33	27	20	12	10	1	188
1975	0	0	0	0	0	0	1	2	3	11	15	27	25	38	29	16	14	11	1	193
1976	0	0	0	0	0	0	0	1	5	7	20	26	29	44	32	15	11	18	0	208
1977	0	0	0	0	0	0	2	0	1	10	16	27	35	34	23	20	16	6	0	190
1978	0	0	0	0	0	0	1	2	3	8	13	22	25	29	25	17	13	5	3	166
1974-78	0	0	0	0	0	0	4	6	17	44	81	125	145	178	136	88	66	50	5	945
1979	0	0	0	0	0	0	0	1	2	9	15	26	31	35	24	15	11	11	0	180
1980	0	0	0	0	0	0	1	1	4	12	20	29	35	41	30	26	10	6	0	215
1981	0	0	0	0	0	0	0	3	0	6	10	24	23	32	28	20	13	7	0	166
1982	0	0	0	0	1	0	0	1	1	8	11	37	32	38	30	26	13	9	1	208
1983	0	0	0	0	0	0	0	0	4	7	14	17	38	40	36	24	17	5	0	202
1979-83	0	0	0	0	1	0	1	6	11	42	70	133	159	186	148	111	64	38	1	971

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OESOPHAGUS (ICD9 150)

YEAR	AGE SPECIFIC RATES																	SUMMARY RATES		
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.8	4.8	8.2	6.4	14.8	37.8	39.5	35.2	63.0	51.5	4.6	4.5	0.6
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.8	2.1	4.0	11.9	18.0	21.9	27.4	61.8	61.8	28.8	4.2	3.9	0.4
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	3.4	1.5	9.5	12.9	17.6	24.7	35.5	45.1	50.8	78.6	4.7	4.6	0.5
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.8	7.9	6.8	16.4	17.9	28.4	36.7	48.4	26.5	82.8	5.0	4.9	0.6
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	2.9	8.2	12.1	24.6	28.8	26.7	35.6	44.8	40.0	4.4	4.3	0.5
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.8	3.9	7.2	12.0	18.5	28.3	33.1	45.0	49.0	56.3	4.6	4.4	0.5
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.9	2.8	7.1	13.5	17.1	21.0	22.3	29.1	44.1	70.1	4.0	3.9	0.4
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	3.5	9.2	13.2	15.1	24.3	23.3	47.4	46.8	54.9	4.2	4.1	0.4
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.4	3.9	7.4	10.6	19.8	20.7	26.9	32.7	53.2	57.8	4.0	3.9	0.5
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.6	3.9	7.0	17.0	18.4	25.0	19.6	44.5	69.7	55.9	4.6	4.3	0.5
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	1.6	8.1	7.7	17.1	22.0	21.8	27.6	50.2	48.1	33.0	4.9	4.7	0.5
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.3	1.6	4.5	7.7	14.3	18.6	22.6	24.8	40.9	52.5	53.8	4.3	4.2	0.5
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.0	3.4	7.8	13.6	20.6	29.2	32.8	39.5	41.1	54.1	4.6	4.5	0.5
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	1.2	4.6	6.7	15.7	16.1	32.8	34.4	30.5	47.9	57.9	4.6	4.5	0.6
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.1	2.9	8.8	14.5	18.4	36.5	37.8	27.6	37.0	92.8	4.9	4.8	0.6
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.4	4.2	7.0	14.3	22.2	27.0	26.0	35.4	53.0	30.5	4.3	4.2	0.5
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	1.3	3.3	5.7	11.1	15.9	22.3	27.6	28.9	42.2	25.0	3.6	3.5	0.4
1974-78	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.5	1.4	3.7	7.2	13.8	18.6	29.4	31.5	32.3	44.3	51.8	4.4	4.3	0.5
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	3.7	6.5	12.5	19.8	26.0	25.6	24.5	34.6	53.7	3.9	3.9	0.5
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	1.6	5.0	8.6	13.6	21.7	29.5	30.9	41.3	30.2	29.0	4.5	4.5	0.6
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	0.0	2.5	4.2	11.1	13.6	22.7	27.8	30.9	38.1	33.0	3.4	3.3	0.4
1982	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.3	0.4	3.4	4.6	17.0	18.0	26.6	28.7	39.1	36.4	41.7	4.2	4.1	0.5
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.5	3.0	5.9	7.7	20.2	28.1	33.0	35.0	45.0	22.4	4.0	3.9	0.5
1979-83	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.4	0.9	3.5	6.0	12.4	18.7	26.6	29.3	34.3	37.1	35.7	4.0	3.9	0.5

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OESOPHAGUS (ICD9 150)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	3	5	6	4	13	11	8	10	7	0	67
1965	0	0	0	0	0	0	0	0	1	3	7	7	7	8	11	14	9	8	0	75
1966	0	0	0	0	0	0	1	0	1	2	3	5	6	6	6	11	8	4	0	53
1967	0	0	0	0	0	0	0	1	2	5	2	5	6	6	12	11	6	7	0	63
1968	0	0	0	0	0	1	1	0	2	2	4	8	10	10	9	15	9	7	0	78
1964-68	0	0	0	0	0	1	2	1	6	15	21	31	33	43	49	59	42	33	0	336
1969	0	0	0	0	1	0	0	2	2	2	6	13	8	14	11	9	15	8	0	91
1970	0	0	0	0	0	0	0	0	3	4	3	6	14	9	7	9	9	9	0	73
1971	0	0	0	0	0	0	0	0	1	4	5	7	5	11	10	7	8	5	0	63
1972	0	0	0	0	0	0	1	0	1	2	6	10	10	12	11	12	11	14	0	90
1973	2	0	0	0	0	1	0	0	3	4	4	4	10	12	16	15	15	5	0	91
1969-73	2	0	0	0	1	1	1	2	10	16	24	40	47	58	55	52	58	41	0	408
1974	0	0	1	0	0	0	1	1	3	0	8	9	6	11	14	12	13	6	0	85
1975	0	0	0	1	1	0	1	0	1	6	5	6	9	13	16	8	16	19	0	102
1976	0	0	1	0	0	1	0	3	0	2	6	5	11	14	14	13	16	15	0	101
1977	0	0	0	0	0	0	2	0	2	0	7	9	8	13	12	18	11	13	1	96
1978	0	0	0	0	1	0	0	1	0	3	4	12	11	12	8	12	16	10	0	90
1974-78	0	0	2	1	2	1	4	5	6	11	30	41	45	63	64	63	72	63	1	474
1979	0	0	0	0	0	0	0	0	2	0	5	8	13	13	11	13	11	17	1	94
1980	0	0	0	0	0	0	0	1	0	2	5	9	9	21	13	9	12	23	0	104
1981	0	0	0	0	0	0	2	0	1	0	1	12	14	13	22	18	9	13	0	105
1982	0	0	1	0	0	0	1	0	0	6	5	4	6	18	12	22	10	13	1	99
1983	0	0	0	0	0	0	0	0	3	5	4	9	9	9	18	15	11	14	0	97
1979-83	0	0	1	0	0	0	3	1	6	13	20	42	51	74	76	77	53	80	2	499

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OESOPHAGUS (ICD9 150)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	3.0	4.3	3.4	12.8	13.3	13.7	28.9	31.8	2.0	1.6	0.2
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.6	4.1	4.9	5.8	7.7	12.9	23.2	25.1	34.8	2.2	1.8	0.2
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.4	1.0	1.7	3.4	4.8	5.7	6.8	17.8	21.5	16.9	1.5	1.2	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	2.5	1.1	3.3	4.7	5.5	13.3	17.0	15.4	27.9	1.8	1.4	0.2
1968	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.8	0.9	2.2	5.1	7.6	9.0	9.9	22.4	22.1	26.4	2.1	1.7	0.2
1964-68	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.5	1.5	2.4	4.2	5.3	8.1	11.2	18.9	22.4	27.4	1.9	1.6	0.2
1969	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.9	0.8	0.9	3.3	8.0	5.9	12.2	11.9	13.1	35.4	28.3	2.4	2.0	0.2
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	1.7	1.6	3.6	10.0	7.6	7.4	12.8	20.4	29.7	1.9	1.6	0.2
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	2.6	4.0	3.5	9.0	10.3	9.7	17.3	15.4	1.5	1.3	0.2
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.8	2.9	5.7	6.7	9.6	11.1	16.2	23.2	40.8	2.1	1.7	0.2
1973	0.7	0.0	0.0	0.0	0.0	0.3	0.0	0.0	1.3	1.7	1.9	2.3	6.5	9.3	15.7	19.7	30.6	13.9	2.1	1.7	0.2
1969-73	0.1	0.0	0.0	0.0	0.1	0.1	0.1	0.2	0.9	1.4	2.4	4.7	6.5	9.5	11.3	14.4	25.3	25.4	2.0	1.7	0.2
1974	0.0	0.0	0.3	0.0	0.0	0.0	0.4	0.4	1.3	0.0	3.5	5.1	3.8	8.3	13.3	15.4	25.9	15.8	1.9	1.6	0.2
1975	0.0	0.0	0.0	0.3	0.3	0.0	0.4	0.0	0.4	2.6	2.1	3.3	5.4	9.6	14.8	10.0	31.0	48.2	2.2	1.8	0.2
1976	0.0	0.0	0.3	0.0	0.0	0.3	0.0	1.2	0.0	0.9	2.5	2.6	6.5	10.0	12.6	15.7	30.0	36.2	2.1	1.7	0.2
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.9	0.0	3.0	4.5	4.7	8.9	10.5	21.3	19.9	30.0	1.9	1.5	0.2
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	1.3	1.7	5.6	6.4	8.0	6.7	13.7	28.0	22.3	1.8	1.4	0.2
1974-78	0.0	0.0	0.1	0.1	0.1	0.1	0.3	0.4	0.5	0.9	2.6	4.3	5.4	9.0	11.5	15.3	26.9	30.4	2.0	1.6	0.2
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	2.1	3.6	7.5	8.4	9.0	14.4	18.5	36.1	1.8	1.4	0.2
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	2.1	3.9	5.0	13.0	10.3	9.6	19.4	46.8	1.9	1.6	0.2
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.4	0.0	0.4	5.1	7.5	7.9	16.9	18.7	14.1	25.2	1.9	1.5	0.2
1982	0.0	0.0	0.3	0.0	0.0	0.0	0.3	0.0	0.0	2.6	2.1	1.7	3.0	10.8	8.9	22.0	15.3	24.1	1.8	1.4	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	2.2	1.7	3.8	4.3	5.4	12.9	14.5	16.2	24.6	1.7	1.4	0.2
1979-83	0.0	0.0	0.1	0.0	0.0	0.0	0.2	0.1	0.5	1.1	1.7	3.6	5.4	9.1	11.6	15.9	16.6	30.9	1.8	1.5	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF STOMACH (ICD9 151)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	1	0	0	0	2	4	9	22	23	43	72	100	116	108	123	71	41	0	736
1965	0	0	0	0	0	2	5	10	14	18	36	54	80	97	97	92	57	33	0	595
1966	0	0	0	0	0	3	1	8	13	31	49	58	65	94	111	94	60	35	0	622
1967	0	0	0	0	0	1	0	4	13	29	33	59	93	98	95	83	55	27	0	590
1968	0	0	0	0	0	1	1	10	8	25	38	54	96	90	90	72	63	39	0	587
1964-68	1	1	0	0	0	9	11	41	70	126	199	297	434	495	501	464	306	175	0	3130
1969	0	1	0	0	1	0	2	6	17	27	36	55	70	92	99	94	68	39	1	608
1970	0	0	0	0	1	2	4	7	13	27	40	60	92	72	85	96	61	40	1	601
1971	0	0	0	0	0	0	2	8	21	23	27	54	89	87	80	89	74	27	0	581
1972	1	0	0	0	1	0	5	3	19	25	39	65	76	96	109	77	43	42	1	602
1973	1	0	0	0	1	1	0	6	14	33	37	58	89	114	105	87	61	34	1	642
1969-73	2	1	0	0	4	3	13	30	84	135	179	292	416	461	478	443	307	182	4	3034
1974	0	0	0	0	0	5	3	7	22	24	46	55	81	85	104	90	39	36	3	600
1975	0	0	0	0	0	3	2	5	19	28	44	57	78	83	95	102	52	32	1	601
1976	0	0	0	0	1	2	4	6	12	25	46	67	72	111	111	87	59	43	3	649
1977	0	0	0	1	0	1	5	6	15	24	37	62	64	76	84	77	47	29	1	529
1978	0	0	0	0	0	0	5	6	10	25	49	65	91	89	87	81	45	33	1	587
1974-78	0	0	0	1	1	11	19	30	78	126	222	306	386	444	481	437	242	173	9	2966
1979	0	0	0	0	0	1	2	8	13	26	37	57	75	78	83	98	51	40	0	569
1980	0	0	0	0	0	0	6	5	10	22	39	62	68	83	99	84	51	44	4	577
1981	0	0	0	0	0	0	4	4	8	22	47	65	93	97	88	76	62	28	2	596
1982	0	0	0	0	0	1	3	4	14	18	37	57	69	75	84	85	61	38	2	548
1983	0	0	0	0	0	2	2	14	10	21	43	56	99	86	89	85	46	44	1	598
1979-83	0	0	0	0	0	4	17	35	55	109	203	297	404	419	443	428	271	194	9	2888

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF STOMACH (ICD9 151)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.3	0.0	0.0	0.0	1.0	1.8	3.8	10.1	12.2	25.3	51.5	87.8	128.9	158.1	270.9	279.5	301.5	22.3	21.3	2.4
1965	0.0	0.0	0.0	0.0	0.0	1.0	2.2	4.2	6.2	9.4	20.6	37.7	68.6	106.0	139.8	200.4	220.1	237.4	17.7	16.9	2.0
1966	0.0	0.0	0.0	0.0	0.0	1.4	0.4	3.3	5.6	15.9	27.4	39.4	54.3	101.0	157.7	201.7	230.8	250.0	18.1	17.3	2.0
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.7	5.5	14.4	18.1	38.8	75.7	103.2	134.0	174.7	208.3	186.2	16.8	16.3	1.9
1968	0.0	0.0	0.0	0.0	0.0	0.4	0.4	4.2	3.3	12.0	20.8	34.4	76.3	92.6	126.6	150.6	235.1	260.0	16.5	16.0	1.8
1964-68	0.1	0.1	0.0	0.0	0.0	0.8	1.0	3.4	6.1	12.8	22.4	40.2	72.5	106.0	143.1	199.0	234.5	246.5	18.2	17.5	2.0
1969	0.0	0.2	0.0	0.0	0.3	0.0	0.9	2.5	7.0	12.4	19.7	33.9	54.3	92.0	137.9	195.4	250.0	248.4	16.7	15.9	1.8
1970	0.0	0.0	0.0	0.0	0.3	0.7	1.7	2.9	5.4	11.9	21.6	35.9	69.5	69.9	116.6	197.9	219.4	243.9	16.1	15.4	1.7
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.8	3.3	8.7	9.9	14.3	31.7	65.4	82.0	107.5	182.0	262.4	156.1	15.3	14.5	1.6
1972	0.3	0.0	0.0	0.0	0.3	0.0	2.0	1.3	7.8	10.7	19.6	38.1	53.9	88.9	142.1	155.9	149.8	234.6	15.4	15.0	1.8
1973	0.3	0.0	0.0	0.0	0.3	0.3	0.0	2.5	5.7	14.0	17.7	34.2	61.2	103.5	131.9	174.7	209.6	186.8	16.2	15.6	1.8
1969-73	0.1	0.1	0.0	0.0	0.2	0.2	1.1	2.5	6.9	11.8	18.6	34.7	60.8	87.4	127.3	181.0	217.7	212.9	16.0	15.3	1.7
1974	0.0	0.0	0.0	0.0	0.0	1.5	1.1	2.9	9.0	10.2	21.0	32.6	53.9	75.3	126.4	177.9	133.6	194.6	14.8	14.2	1.7
1975	0.0	0.0	0.0	0.0	0.0	0.9	0.7	2.1	7.8	11.8	19.7	33.1	50.3	71.6	112.6	194.7	178.1	168.4	14.6	13.8	1.5
1976	0.0	0.0	0.0	0.0	0.3	0.6	1.4	2.4	5.0	10.5	20.3	37.4	45.7	92.1	128.5	159.9	198.7	221.6	15.2	14.6	1.7
1977	0.0	0.0	0.0	0.2	0.0	0.3	1.6	2.4	6.3	10.0	16.3	32.8	40.6	60.4	95.1	136.3	155.6	147.2	12.1	11.6	1.3
1978	0.0	0.0	0.0	0.0	0.0	0.0	1.5	2.3	4.2	10.3	21.4	32.7	57.8	68.4	95.9	137.5	146.1	165.0	13.1	12.7	1.5
1974-78	0.0	0.0	0.0	0.0	0.1	0.6	1.3	2.4	6.5	10.5	19.7	33.7	49.6	73.3	111.3	160.2	162.3	179.1	13.9	13.4	1.5
1979	0.0	0.0	0.0	0.0	0.0	0.3	0.6	2.9	5.4	10.8	16.1	27.3	47.9	58.0	88.6	160.1	160.4	195.1	12.4	11.9	1.3
1980	0.0	0.0	0.0	0.0	0.0	0.0	1.7	1.8	4.1	9.2	16.7	29.1	42.2	59.8	102.1	133.5	154.1	212.6	12.2	11.7	1.3
1981	0.0	0.0	0.0	0.0	0.0	0.0	1.1	1.4	3.2	9.3	20.0	30.1	55.1	68.7	87.4	117.5	181.8	132.1	12.3	11.9	1.4
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.9	1.3	5.5	7.6	15.6	26.1	38.8	52.6	80.3	127.8	170.9	175.9	11.1	10.5	1.1
1983	0.0	0.0	0.0	0.0	0.0	0.5	0.6	4.3	3.8	8.9	16.1	25.3	52.7	60.4	81.6	124.1	121.7	197.3	11.8	11.4	1.3
1979-83	0.0	0.0	0.0	0.0	0.0	0.2	1.0	2.4	4.4	9.2	17.3	27.6	47.4	59.9	87.7	132.2	157.1	182.5	12.0	11.5	1.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF STOMACH (ICD9 151)

YEAR OF DIAGNOSIS	AGE (YEARS)																	ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	2	17	10	12	21	28	43	47	57	83	52	42	0	414
1965	0	0	0	0	0	2	3	5	9	16	20	17	34	39	56	71	54	31	0	357
1966	1	0	0	0	0	1	4	12	13	8	12	21	51	45	50	65	49	47	0	379
1967	0	0	0	0	1	1	1	4	8	14	19	23	36	51	64	59	45	35	0	361
1968	0	0	0	0	0	0	4	4	9	14	18	22	28	36	44	69	36	36	1	321
1964-68	1	0	0	0	1	4	14	42	49	64	90	111	192	218	271	347	236	191	1	1832
1969	0	0	0	0	1	1	3	3	5	12	21	24	28	48	63	57	36	44	0	346
1970	0	0	0	0	1	0	2	8	8	17	12	27	30	46	42	53	42	45	0	333
1971	0	0	0	0	1	4	2	2	7	16	17	32	34	50	70	66	55	51	0	407
1972	0	0	0	1	1	0	1	5	13	10	15	22	29	46	52	58	47	63	0	363
1973	1	0	0	0	1	1	2	5	8	8	24	16	36	41	60	61	61	44	0	369
1969-73	1	0	0	1	5	6	10	23	41	63	89	121	157	231	287	295	241	247	0	1818
1974	0	0	0	0	0	1	4	11	4	23	20	22	34	47	66	54	51	49	0	386
1975	0	0	0	0	2	0	1	5	5	18	15	24	33	47	53	83	66	44	0	396
1976	0	0	0	0	1	0	0	4	6	18	24	26	28	50	57	46	47	45	1	353
1977	0	0	0	1	1	3	4	6	8	11	20	21	22	43	63	48	44	40	0	335
1978	0	0	0	1	1	2	7	8	6	11	20	33	22	42	52	65	59	46	0	375
1974-78	0	0	0	2	5	6	16	34	29	81	99	126	139	229	291	296	267	224	1	1845
1979	0	0	0	0	2	2	3	6	10	15	14	22	26	42	60	54	52	38	0	346
1980	0	0	0	1	2	0	2	4	5	10	18	27	40	50	50	62	45	49	3	368
1981	0	0	0	0	0	5	2	4	8	17	19	33	36	47	50	73	55	54	0	403
1982	0	0	1	0	0	2	3	4	8	15	13	24	26	46	56	73	60	60	1	392
1983	0	0	0	1	0	2	4	2	8	21	11	29	34	41	50	60	56	67	1	387
1979-83	0	0	1	2	4	11	14	20	39	78	75	135	162	226	266	322	268	268	5	1896

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF STOMACH (ICD9 151)

YEAR	AGE SPECIFIC RATES																	SUMMARY RATES		
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.9	7.3	4.5	6.5	12.8	20.3	36.4	46.3	68.7	141.6	150.3	190.9	12.6	9.9	1.0
1965	0.0	0.0	0.0	0.0	0.0	0.9	1.4	2.1	3.9	8.5	11.8	12.0	28.1	37.6	65.4	117.5	150.4	134.8	10.6	8.3	0.9
1966	0.3	0.0	0.0	0.0	0.0	0.5	1.8	5.2	5.5	4.1	6.9	14.4	41.0	42.4	56.4	105.0	131.7	198.3	10.9	8.7	0.9
1967	0.0	0.0	0.0	0.0	0.4	0.4	0.4	1.7	3.3	6.9	10.6	15.2	28.0	47.0	71.0	91.2	115.4	139.4	10.0	8.1	0.9
1968	0.0	0.0	0.0	0.0	0.0	0.0	1.8	1.7	3.7	6.6	9.9	14.0	21.2	32.3	48.2	103.0	88.5	135.8	8.6	6.8	0.7
1964-68	0.1	0.0	0.0	0.0	0.1	0.4	1.3	3.6	4.2	6.5	10.4	15.1	30.8	41.0	61.8	111.0	125.9	158.8	10.5	8.3	0.9
1969	0.0	0.0	0.0	0.0	0.3	0.4	1.3	1.3	2.1	5.4	11.5	14.7	20.6	41.9	68.0	82.8	84.9	155.5	9.0	7.3	0.8
1970	0.0	0.0	0.0	0.0	0.3	0.0	0.9	3.5	3.4	7.4	6.4	16.0	21.5	39.0	44.4	75.2	95.0	148.5	8.4	6.8	0.7
1971	0.0	0.0	0.0	0.0	0.3	1.4	0.8	0.9	3.0	6.7	8.8	18.5	23.6	41.1	72.1	91.5	119.0	156.9	9.9	7.9	0.9
1972	0.0	0.0	0.0	0.3	0.3	0.0	0.4	2.2	5.6	4.2	7.4	12.6	19.5	36.7	52.5	78.1	98.9	183.7	8.6	6.8	0.7
1973	0.3	0.0	0.0	0.0	0.3	0.3	0.8	2.2	3.4	3.4	11.1	9.1	23.4	31.9	58.9	80.3	124.5	122.6	8.5	6.7	0.7
1969-73	0.1	0.0	0.0	0.1	0.3	0.4	0.8	2.0	3.5	5.4	9.1	14.2	21.7	38.0	59.1	81.6	105.1	153.1	8.9	7.1	0.8
1974	0.0	0.0	0.0	0.0	0.0	0.3	1.5	4.7	1.7	9.8	8.8	12.5	21.3	35.5	62.9	69.3	101.8	129.3	8.7	7.0	0.8
1975	0.0	0.0	0.0	0.0	0.5	0.0	0.4	2.1	2.1	7.7	6.4	13.2	19.9	34.8	49.0	103.7	127.9	111.7	8.7	6.7	0.7
1976	0.0	0.0	0.0	0.0	0.3	0.0	0.0	1.6	2.6	7.7	10.1	13.7	16.6	35.6	51.4	55.6	88.2	108.7	7.5	6.1	0.7
1977	0.0	0.0	0.0	0.3	0.3	0.8	1.3	2.4	3.4	4.7	8.5	10.4	12.9	29.5	54.9	56.7	79.4	92.4	6.9	5.6	0.6
1978	0.0	0.0	0.0	0.2	0.3	0.6	2.1	3.1	2.6	4.7	8.5	15.4	12.8	28.0	43.8	74.4	103.1	102.4	7.5	5.9	0.6
1974-78	0.0	0.0	0.0	0.1	0.3	0.3	1.1	2.8	2.5	6.9	8.5	13.1	16.6	32.5	52.2	71.7	99.8	108.3	7.8	6.2	0.7
1979	0.0	0.0	0.0	0.0	0.5	0.5	0.9	2.2	4.2	6.4	6.0	9.8	15.0	27.1	48.8	59.9	87.2	80.7	6.7	5.4	0.6
1980	0.0	0.0	0.0	0.2	0.5	0.8	0.6	1.4	2.1	4.3	7.7	11.7	22.4	31.0	39.6	66.5	72.8	99.8	6.9	5.6	0.6
1981	0.0	0.0	0.0	0.0	0.0	1.3	0.5	1.4	3.3	7.3	8.1	14.2	19.2	28.6	38.4	75.8	86.1	104.9	7.4	5.9	0.6
1982	0.0	0.0	0.3	0.0	0.0	0.5	0.8	1.3	3.2	6.5	5.6	10.3	13.1	27.5	41.7	72.9	91.7	111.1	6.9	5.4	0.6
1983	0.0	0.0	0.0	0.3	0.0	0.5	1.1	0.6	3.0	9.0	4.7	12.4	16.2	24.6	35.8	57.9	82.2	117.8	6.6	5.3	0.5
1979-83	0.0	0.0	0.1	0.1	0.2	0.6	0.8	1.3	3.2	6.7	6.4	11.7	17.1	27.7	40.7	66.6	84.1	103.6	6.9	5.5	0.6

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF SMALL INTESTINE, INCLUDING DUODENUM (ICD9 152)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	1	0	1	3	3	8	2	2	2	3	1	1	0	27
1965	0	0	0	0	0	0	0	0	1	3	1	1	3	0	3	0	2	1	0	15
1966	0	0	0	0	0	0	0	0	1	1	5	3	3	1	3	1	1	2	0	21
1967	0	0	0	1	0	0	0	0	1	1	1	0	1	2	1	2	2	0	0	12
1968	0	0	0	0	0	0	1	0	1	0	1	1	5	1	1	1	1	0	0	13
1964-68	0	0	0	1	0	0	2	0	5	8	11	13	14	6	10	7	7	4	0	88
1969	0	0	1	0	0	0	0	0	0	2	2	1	2	4	0	2	3	0	0	17
1970	0	0	0	0	0	0	0	0	1	0	5	0	4	3	3	4	1	1	0	22
1971	0	0	0	0	0	0	0	0	2	4	2	4	1	4	3	4	1	0	0	25
1972	0	0	0	0	0	0	1	0	0	2	2	3	4	3	2	1	1	0	0	20
1973	0	0	0	0	1	0	0	0	2	4	3	3	0	5	1	3	2	1	0	25
1969-73	0	0	1	0	1	0	1	1	5	12	14	11	11	19	9	14	8	2	0	109
1974	0	0	0	0	0	0	0	0	0	2	2	2	5	2	4	4	0	3	0	24
1975	0	0	0	0	0	0	1	0	0	2	5	2	10	5	4	2	2	1	0	34
1976	0	0	0	0	0	0	1	3	1	1	6	3	9	5	3	3	1	0	0	36
1977	1	0	0	0	3	0	0	0	1	0	1	3	5	4	3	0	1	4	0	23
1978	0	0	0	1	0	0	0	0	0	0	1	3	7	7	6	2	3	0	0	41
1974-78	1	0	0	0	0	1	1	0	3	3	2	7	7	6	6	2	3	0	0	41
1979	0	0	0	1	0	1	3	3	4	8	16	17	36	22	20	11	7	8	0	158
1980	0	0	0	0	0	2	1	1	1	4	3	9	4	5	7	1	3	1	0	42
1981	0	0	0	0	0	1	0	2	2	2	3	3	2	3	2	3	3	0	0	26
1982	0	0	0	0	0	1	0	0	2	2	6	4	6	6	8	5	2	1	0	43
1983	0	0	0	0	0	0	0	0	2	4	4	7	9	4	11	5	1	1	1	49
1979-83	0	0	0	0	0	4	1	6	8	13	17	28	28	26	36	21	9	4	1	202

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF SMALL INTESTINE, INCLUDING DUODENUM (ICD9 152)

YEAR	AGE SPECIFIC RATES																	SUMMARY RATES		
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	1.6	1.8	5.7	1.8	2.2	2.9	6.6	3.9	7.4	0.8	0.8	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.6	0.6	0.7	2.6	0.0	4.3	0.0	7.7	7.2	0.4	0.4	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	2.8	2.0	2.5	1.1	4.3	2.1	3.8	14.3	0.6	0.6	0.1
1967	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.5	0.5	0.0	0.8	2.1	1.4	4.2	7.6	0.0	0.3	0.3	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.0	0.5	0.6	4.0	1.0	1.4	2.1	3.7	0.0	0.4	0.4	0.0
1964-68	0.0	0.0	0.0	0.1	0.0	0.0	0.2	0.0	0.4	0.8	1.2	1.8	2.3	1.3	2.9	3.0	5.4	5.6	0.5	0.5	0.1
1969	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.1	0.6	1.6	4.0	0.0	4.2	11.0	0.0	0.5	0.4	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	2.7	0.0	3.0	2.9	4.1	8.2	3.6	6.1	0.6	0.6	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.7	1.1	2.3	0.7	3.8	4.0	8.2	3.5	0.0	0.6	0.6	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.9	1.0	1.8	2.8	2.8	2.6	2.0	3.5	0.0	0.5	0.5	0.1
1973	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.8	1.7	1.4	1.8	0.0	4.5	1.3	6.0	6.9	5.5	0.6	0.6	0.1
1969-73	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.4	1.0	1.5	1.3	1.6	3.6	2.4	5.7	5.7	2.3	0.6	0.5	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.9	1.2	3.3	1.8	4.9	7.9	0.0	16.2	0.6	0.6	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	2.8	1.2	6.4	4.3	4.7	3.8	6.8	5.3	0.8	0.8	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.3	1.2	0.4	0.4	2.6	1.7	5.7	4.1	3.5	5.5	3.4	0.0	0.8	0.8	0.1
1977	0.3	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.6	3.2	3.2	3.4	0.0	3.3	20.3	0.8	0.8	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0	1.3	1.2	0.9	3.5	4.4	4.6	6.6	3.4	9.7	0.0	0.5	0.6	0.1
1974-78	0.1	0.0	0.0	0.0	0.0	0.1	0.2	0.2	0.3	0.7	1.4	1.9	4.6	3.6	4.6	4.0	4.7	8.3	0.9	0.9	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.6	0.3	0.4	0.4	1.7	1.3	4.3	2.6	3.7	7.5	1.6	9.4	4.9	0.7	0.7	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.7	0.8	0.8	1.3	1.4	1.2	2.2	2.1	4.8	9.1	0.0	0.9	0.9	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	0.8	2.5	1.9	3.6	4.3	7.9	7.7	5.9	4.7	0.6	0.5	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.7	1.7	3.2	5.1	2.8	10.5	7.5	2.8	4.6	0.9	0.9	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	0.4	0.4	2.3	3.7	5.6	7.3	10.2	0.0	4.5	1.0	1.0	0.1
1979-83	0.0	0.0	0.0	0.0	0.0	0.2	0.1	0.4	0.6	1.1	1.4	2.6	3.3	3.7	7.1	6.5	5.2	3.8	0.8	0.8	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF SMALL INTESTINE, INCLUDING DUODENUM (ICD9 152)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	0	0	4	2	5	3	1	4	3	2	4	0	0	28
1965	0	0	0	0	0	0	0	1	1	1	0	3	5	2	2	6	3	0	0	24
1966	0	0	0	0	0	1	0	0	1	2	2	2	1	4	1	1	3	1	0	19
1967	0	0	0	0	0	0	1	0	0	1	0	2	1	1	3	1	3	0	0	13
1968	0	0	0	0	0	0	0	1	0	3	2	4	4	4	4	1	1	0	0	24
1964-68	0	0	0	0	0	1	1	2	6	9	9	14	12	15	13	11	14	1	0	108
1969	0	0	0	1	0	0	2	0	0	0	2	4	4	2	3	2	0	3	0	23
1970	0	0	0	0	0	0	0	0	2	3	2	4	3	0	3	1	2	0	0	20
1971	0	0	0	0	0	0	0	1	1	0	1	5	3	5	4	4	1	1	0	26
1972	0	0	0	0	0	0	0	0	1	1	4	3	4	4	3	6	2	1	0	29
1973	0	0	0	0	0	1	1	1	0	0	0	2	2	1	2	6	2	3	0	21
1969-73	0	0	0	1	0	1	3	2	4	4	9	18	16	12	15	19	7	8	0	119
1974	0	0	0	0	0	1	0	1	1	1	2	3	3	5	10	2	3	2	0	34
1975	0	0	0	0	0	0	0	1	1	2	0	1	2	4	7	5	2	2	0	27
1976	0	0	0	0	0	0	0	0	0	1	5	5	3	2	5	2	2	0	0	25
1977	0	0	0	0	0	0	0	0	0	4	2	1	3	6	7	3	3	2	0	31
1978	0	0	0	0	0	0	0	0	1	2	3	4	5	8	3	2	2	2	0	32
1974-78	0	0	0	0	0	1	0	2	3	10	12	14	16	25	32	14	12	8	0	149
1979	0	0	0	0	0	1	0	1	0	0	1	4	5	5	4	5	2	5	0	33
1980	0	0	0	0	0	0	0	1	1	3	1	3	4	7	4	7	4	2	0	37
1981	0	0	0	0	0	0	0	1	2	3	1	5	5	5	4	5	2	4	0	37
1982	0	0	0	0	1	0	0	0	4	0	3	4	4	7	8	7	3	2	0	43
1983	0	0	0	0	0	1	0	2	1	2	2	6	5	6	3	4	5	5	0	42
1979-83	0	0	0	0	1	2	0	5	8	8	8	22	23	30	23	28	16	18	0	192

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF SMALL INTESTINE, INCLUDING DUODENUM (ICD9 152)

YEAR	AGE SPECIFIC RATES																	SUMMARY RATES		
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.8	1.1	3.0	2.2	0.8	3.9	3.6	3.4	11.6	0.0	0.8	0.7	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.0	2.1	4.1	1.9	2.3	9.9	8.4	0.0	0.7	0.6	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.4	1.0	1.1	1.4	0.8	3.8	1.1	1.6	8.1	4.2	0.5	0.5	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.6	0.5	0.0	1.3	0.8	0.9	3.3	1.5	7.7	0.0	0.4	0.3	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.4	1.1	2.5	3.0	3.6	4.4	1.5	2.5	0.0	0.7	0.6	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.5	0.9	1.0	1.9	1.9	2.8	3.0	3.5	7.5	0.8	0.6	0.5	0.1
1969	0.0	0.0	0.0	0.3	0.0	0.0	0.9	0.0	0.0	0.0	1.1	2.5	2.9	1.7	3.2	2.9	0.0	10.6	0.6	0.6	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.3	1.1	2.4	2.1	0.0	3.2	1.4	4.5	0.0	0.5	0.5	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.5	2.9	2.1	4.1	4.1	5.5	2.2	3.1	0.6	0.6	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.0	1.7	2.7	3.2	3.0	8.1	4.2	2.9	0.7	0.6	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.4	0.0	0.0	0.0	1.1	1.3	0.8	2.0	7.9	4.1	8.4	0.5	0.4	0.0
1969-73	0.0	0.0	0.0	0.1	0.0	0.1	0.3	0.2	0.3	0.3	0.9	2.1	2.2	2.0	3.1	5.3	3.1	5.0	0.6	0.5	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.4	0.4	0.9	1.7	1.9	3.8	9.5	2.6	6.0	5.3	0.8	0.7	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.9	0.0	0.6	1.2	3.0	6.5	6.3	3.9	5.1	0.6	0.5	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.1	2.6	1.8	1.4	4.5	2.4	3.8	0.0	0.6	0.5	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	0.8	0.5	1.8	4.1	6.1	3.5	5.4	4.6	0.6	0.6	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	1.3	1.9	2.9	5.3	2.5	2.3	3.5	4.5	0.7	0.6	0.1
1974-78	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.3	0.9	1.0	1.5	1.9	3.6	5.7	3.4	4.5	3.9	0.6	0.6	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.0	0.0	0.4	1.8	2.9	3.2	3.3	5.5	3.4	10.6	0.6	0.5	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.3	0.4	1.3	2.2	4.3	3.2	7.5	6.5	4.1	0.7	0.6	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	1.3	0.4	2.1	2.7	3.0	3.1	5.2	3.1	7.8	0.7	0.6	0.1
1982	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	1.6	0.0	1.3	1.7	2.0	4.2	6.0	7.0	4.6	3.7	0.8	0.7	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.6	0.4	0.9	0.9	2.6	2.4	3.6	2.1	3.9	7.3	8.8	0.7	0.6	0.1
1979-83	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.3	0.6	0.7	0.7	1.9	2.4	3.7	3.5	5.8	5.0	7.8	0.7	0.6	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF COLON (ICD9 153)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	2	2	3	7	14	19	41	66	89	115	132	126	129	86	54	0	885
1965	0	0	0	3	1	3	6	11	10	34	59	93	83	121	122	117	100	51	0	814
1966	0	0	0	0	0	4	5	12	25	25	47	88	101	115	110	96	84	45	0	757
1967	0	0	0	1	0	2	5	10	9	25	65	89	104	119	133	124	85	46	0	817
1968	0	0	0	1	1	6	8	12	24	28	43	83	117	117	128	95	66	46	0	775
1964-68	0	0	0	7	4	18	31	59	87	153	280	442	520	604	619	561	421	242	0	4048
1969	0	0	2	2	2	2	6	11	16	47	62	101	126	142	139	104	86	54	0	902
1970	0	0	1	1	5	2	5	13	27	33	64	108	114	122	130	108	82	68	0	883
1971	0	0	0	1	3	5	3	9	26	42	59	92	108	137	123	117	75	53	0	853
1972	1	1	0	1	1	3	5	6	30	50	54	84	136	139	137	116	71	63	2	900
1973	0	1	0	2	1	8	6	11	21	52	86	104	163	174	176	127	98	60	3	1093
1969-73	1	2	3	7	12	20	25	50	120	224	325	489	647	714	705	572	412	298	5	4631
1974	0	1	1	0	1	3	7	15	26	41	76	105	132	167	177	143	106	66	2	1069
1975	1	0	0	1	3	3	12	11	25	41	68	112	163	174	168	152	77	69	3	1083
1976	0	0	0	3	5	2	4	10	23	52	83	157	173	195	202	160	129	69	2	1269
1977	0	0	0	1	2	7	11	13	29	51	90	133	167	181	192	140	102	74	3	1196
1978	1	1	0	4	3	6	13	5	18	48	101	154	178	187	214	147	119	98	3	1300
1974-78	2	2	1	9	14	21	47	54	121	233	418	661	813	904	953	742	533	376	13	5917
1979	0	0	0	1	2	3	13	15	20	57	87	137	164	228	234	165	141	63	2	1332
1980	0	0	0	1	0	5	9	23	27	40	83	147	166	264	191	196	148	74	1	1375
1981	0	0	0	1	1	2	11	11	33	55	96	155	190	239	244	195	125	84	4	1446
1982	0	0	0	0	1	6	10	13	30	41	87	149	217	264	272	194	152	89	3	1528
1983	1	0	0	0	1	5	14	19	21	45	104	146	221	243	294	209	158	99	7	1587
1979-83	1	0	0	3	5	21	57	81	131	238	457	734	958	1238	1235	959	724	409	17	7268

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF COLON (ICD9 153)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.7	0.9	1.5	3.1	5.9	8.7	21.7	38.8	63.7	101.0	146.7	184.5	284.1	338.6	397.1	26.8	25.8	2.8
1965	0.0	0.0	0.0	1.0	0.4	1.4	2.7	4.6	4.4	17.8	33.8	64.9	71.1	132.2	175.8	254.9	386.1	366.9	24.2	22.9	2.5
1966	0.0	0.0	0.0	0.0	0.0	1.9	2.2	5.0	10.7	12.8	26.3	59.8	84.4	123.5	156.2	206.8	323.1	321.4	22.1	21.2	2.4
1967	0.0	0.0	0.0	0.3	0.0	0.9	2.2	4.2	3.8	12.4	35.7	58.6	84.6	125.3	187.6	261.1	322.0	317.2	23.4	22.3	2.5
1968	0.0	0.0	0.0	0.3	0.4	2.5	3.5	5.0	10.0	13.4	23.5	52.8	92.9	120.4	180.0	198.7	246.3	306.7	21.7	21.1	2.5
1964-68	0.0	0.0	0.0	0.5	0.3	1.6	2.7	4.9	7.5	15.5	31.5	59.8	86.8	129.4	176.8	240.6	322.6	340.8	23.6	22.6	2.6
1969	0.0	0.0	0.5	0.6	0.7	0.8	2.6	4.6	6.6	21.6	33.9	62.2	97.8	142.0	193.6	216.2	316.2	343.9	24.8	24.0	2.8
1970	0.0	0.0	0.3	0.3	1.6	0.7	2.1	5.4	11.2	14.6	34.6	64.6	86.1	118.4	178.3	222.7	295.0	414.6	23.7	22.9	2.6
1971	0.0	0.0	0.0	0.3	0.9	1.7	1.2	3.8	10.7	18.1	31.3	54.0	79.4	129.1	165.3	239.3	266.0	306.4	22.4	21.6	2.4
1972	0.3	0.3	0.0	0.3	0.3	1.0	2.0	2.5	12.3	21.4	27.2	49.2	96.5	128.7	178.6	234.8	247.4	352.0	23.1	22.4	2.6
1973	0.0	0.3	0.0	0.5	0.3	2.5	2.3	4.6	8.6	22.1	41.2	61.3	112.0	158.8	221.1	255.0	336.8	329.7	27.5	26.6	3.1
1969-73	0.1	0.1	0.1	0.4	0.7	1.4	2.1	4.2	9.9	19.6	33.7	58.2	94.6	135.4	187.8	233.8	292.2	348.5	24.3	23.5	2.7
1974	0.0	0.3	0.2	0.0	0.3	0.9	2.5	6.2	10.6	17.3	34.7	62.3	87.8	147.9	215.1	282.6	363.0	356.8	26.5	25.3	2.9
1975	0.3	0.0	0.0	0.2	0.8	0.9	4.2	4.5	10.3	17.2	30.4	65.1	105.0	150.0	199.1	290.1	263.7	363.2	26.2	25.2	2.9
1976	0.0	0.0	0.0	0.7	1.4	0.6	1.4	4.0	9.6	21.8	36.5	87.7	109.8	161.8	233.8	294.1	434.3	355.7	29.9	28.6	3.3
1977	0.0	0.0	0.0	0.2	0.5	2.0	3.5	5.1	12.1	21.2	39.5	70.4	105.8	143.8	217.4	247.8	337.7	375.6	27.4	26.5	3.1
1978	0.3	0.3	0.0	1.0	0.8	1.7	3.9	1.9	7.5	19.9	44.1	77.4	113.0	143.7	235.9	249.6	386.4	490.0	29.0	28.2	3.2
1974-78	0.1	0.1	0.0	0.4	0.8	1.2	3.1	4.3	10.0	19.5	37.1	72.8	104.5	149.3	220.6	272.8	357.5	389.2	27.8	26.7	3.1
1979	0.0	0.0	0.0	0.2	0.5	0.8	3.8	5.5	8.3	23.7	37.8	65.7	104.7	169.5	249.7	269.6	443.4	307.3	28.9	27.8	3.3
1980	0.0	0.0	0.0	0.2	0.0	1.4	2.6	8.1	11.1	16.8	35.6	69.1	103.0	190.2	196.9	311.6	447.1	357.5	29.2	27.9	3.1
1981	0.0	0.0	0.0	0.2	0.3	0.6	3.1	3.8	13.3	23.3	40.8	71.8	112.6	169.4	242.3	301.4	366.6	396.2	29.9	28.9	3.3
1982	0.0	0.0	0.0	0.0	0.2	1.6	2.8	4.2	11.8	17.4	36.7	68.3	122.0	185.8	260.0	291.7	425.8	412.0	30.7	29.6	3.5
1983	0.3	0.0	0.0	0.0	0.2	1.3	4.0	5.8	8.8	19.1	43.9	66.1	117.7	170.8	269.5	305.1	418.0	443.9	31.1	29.8	3.5
1979-83	0.1	0.0	0.0	0.1	0.3	1.2	3.3	5.4	10.5	20.0	39.0	68.2	112.4	177.0	244.5	296.2	419.7	384.8	30.0	28.8	3.4

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF COLON (ICD9 153)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	2	2	6	4	18	31	59	81	105	137	158	168	145	99	88	0	1103
1965	1	1	1	0	2	5	6	20	37	42	64	94	120	128	167	126	111	72	0	997
1966	0	0	0	1	3	3	5	17	26	46	62	79	105	145	154	146	116	65	0	973
1967	1	0	0	1	2	0	5	14	29	39	73	92	92	141	126	138	103	80	0	936
1968	0	0	0	0	1	2	6	16	32	45	66	93	120	124	153	137	128	78	0	1001
1964-68	2	1	1	4	10	16	26	85	155	231	346	463	574	696	768	692	557	383	0	5010
1969	0	1	0	0	3	3	5	13	24	38	80	104	131	150	155	141	110	96	0	1054
1970	0	0	0	1	5	5	12	16	32	43	92	117	120	146	159	170	132	98	0	1148
1971	0	0	1	0	0	5	6	17	34	52	81	110	112	142	136	182	130	99	0	1107
1972	1	0	0	2	1	0	9	8	30	58	78	124	138	147	188	168	126	100	1	1179
1973	0	0	0	3	2	4	8	17	32	50	87	101	145	191	204	197	151	159	2	1353
1969-73	1	1	1	6	11	17	40	71	152	241	418	556	646	776	842	858	649	552	3	5841
1974	1	2	0	1	3	1	8	13	26	42	96	126	146	193	204	219	144	137	3	1365
1975	1	0	0	1	3	5	12	25	20	47	87	92	158	171	211	172	150	130	4	1289
1976	0	0	0	1	1	8	12	16	21	59	91	118	159	204	226	210	178	153	1	1458
1977	0	0	0	2	0	6	7	12	31	42	96	121	181	190	244	205	153	150	0	1440
1978	0	0	0	2	1	5	6	19	29	63	105	138	150	233	205	219	168	153	3	1539
1974-78	2	2	0	7	8	25	45	85	127	253	475	595	834	991	1090	1025	793	723	11	7091
1979	0	0	1	1	2	1	6	18	15	53	81	166	172	217	224	264	190	167	0	1578
1980	0	0	1	2	0	3	3	21	26	47	94	155	185	217	251	216	201	167	5	1594
1981	0	1	1	1	4	5	10	18	26	59	89	138	186	186	239	223	207	172	2	1567
1982	0	0	0	2	1	3	9	15	32	48	89	141	179	269	257	241	192	210	1	1689
1983	0	0	0	2	3	4	8	18	37	46	95	142	198	214	294	291	204	222	6	1784
1979-83	0	1	3	8	10	16	36	90	136	253	448	742	920	1103	1265	1235	994	938	14	8212

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF COLON (ICD9 153)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.8	0.9	2.9	1.8	7.7	14.0	31.9	49.2	76.1	116.1	155.5	202.4	247.4	286.1	400.0	33.6	28.5	3.2
1965	0.3	0.3	0.3	0.0	0.9	2.4	2.7	8.6	16.1	22.2	37.7	66.2	99.1	123.6	195.1	208.6	309.2	313.0	29.4	24.6	2.8
1966	0.0	0.0	0.0	0.3	1.2	1.4	2.3	7.3	11.0	23.5	35.6	54.0	84.3	136.7	173.8	235.9	311.8	274.3	27.9	23.1	2.6
1967	0.3	0.0	0.0	0.3	0.8	0.0	2.2	6.1	12.1	19.2	40.9	60.8	71.7	130.0	139.8	213.3	264.1	318.7	26.0	21.6	2.4
1968	0.0	0.0	0.0	0.0	0.4	0.8	2.7	7.0	13.3	21.2	36.4	59.2	90.8	111.4	167.8	204.5	314.5	294.3	27.0	22.4	2.5
1964-68	0.1	0.1	0.1	0.3	0.8	1.5	2.4	7.3	13.3	23.5	39.8	63.0	92.0	131.0	175.1	221.4	297.2	318.4	28.7	23.9	2.7
1969	0.0	0.3	0.0	0.0	1.0	1.2	2.2	5.7	10.1	17.1	43.6	63.9	96.5	130.9	167.4	204.9	259.4	339.2	27.6	23.2	2.7
1970	0.0	0.0	0.0	0.3	1.6	1.9	5.2	7.0	13.7	18.6	49.2	69.4	86.0	123.6	167.9	241.1	298.6	323.4	29.1	24.2	2.7
1971	0.0	0.0	0.3	0.0	0.0	1.8	2.5	7.4	14.6	21.9	42.1	63.5	77.8	116.7	140.1	252.4	281.4	304.6	27.2	22.5	2.4
1972	0.3	0.0	0.0	0.6	0.3	0.0	3.7	3.5	12.9	24.5	38.3	71.0	92.6	117.4	189.7	226.1	265.3	291.5	28.2	23.6	2.7
1973	0.0	0.0	0.0	0.8	0.6	1.3	3.1	7.4	13.8	21.1	40.3	57.7	94.2	148.5	200.4	259.2	308.2	442.9	31.4	25.8	2.9
1969-73	0.1	0.1	0.1	0.3	0.7	1.2	3.4	6.2	13.0	20.7	42.5	65.1	89.4	127.6	173.5	237.2	283.0	342.2	28.8	23.9	2.7
1974	0.3	0.6	0.0	0.3	0.8	0.3	3.0	5.5	11.1	17.8	42.2	71.5	91.5	145.8	194.5	281.1	287.4	361.5	30.9	25.4	2.9
1975	0.3	0.0	0.0	0.3	0.8	1.4	4.3	10.5	8.6	20.0	37.2	50.7	95.2	126.5	195.2	215.0	290.7	329.9	28.4	23.5	2.7
1976	0.0	0.0	0.0	0.3	0.3	2.2	4.1	6.6	9.0	25.3	38.3	62.0	94.2	145.3	203.8	253.9	334.0	369.6	31.1	25.6	2.9
1977	0.0	0.0	0.0	0.5	0.0	1.7	2.2	4.8	13.3	18.0	40.6	59.8	105.7	130.5	212.5	242.0	276.2	346.4	29.9	24.8	2.9
1978	0.0	0.0	0.0	0.5	0.3	1.4	1.8	7.3	12.4	27.0	44.5	64.3	110.3	155.1	172.7	250.6	293.7	340.8	31.0	26.1	2.9
1974-78	0.1	0.1	0.0	0.4	0.4	1.4	3.0	6.9	10.9	21.6	40.5	61.7	99.5	140.8	195.6	248.4	296.3	349.4	30.3	25.1	2.9
1979	0.0	0.0	0.3	0.2	0.5	0.3	1.8	6.6	6.4	22.7	34.7	73.7	99.2	140.1	182.3	292.7	318.8	354.6	30.8	25.2	2.8
1980	0.0	0.0	0.3	0.5	0.0	0.8	0.8	7.5	10.8	20.2	40.2	67.3	103.5	134.7	198.9	231.5	325.2	340.1	30.2	25.0	2.9
1981	0.0	0.3	0.3	0.3	1.0	1.3	2.7	6.2	10.6	25.5	38.1	59.2	99.1	113.1	183.4	231.6	323.9	334.0	29.0	23.9	2.7
1982	0.0	0.0	0.0	0.5	0.2	0.8	2.5	4.8	12.7	20.7	38.1	60.4	90.3	160.9	191.2	240.8	293.6	388.9	30.1	25.0	2.9
1983	0.0	0.0	0.0	0.5	0.7	1.0	2.2	5.5	14.1	19.8	40.7	60.5	94.2	128.1	210.6	280.6	299.6	390.2	30.9	25.2	2.8
1979-83	0.0	0.1	0.2	0.4	0.5	0.8	2.0	6.1	11.0	21.8	38.3	64.2	97.0	135.4	193.6	255.4	311.8	362.7	30.2	24.8	2.8

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF RECTUM, RECTOSIGMOID JUNCTION AND ANUS (ICD9 154)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	2	0	3	2	9	30	50	72	72	97	89	78	56	15	0	575
1965	0	0	0	0	0	1	3	6	14	29	54	46	58	78	63	83	35	21	0	491
1966	0	0	0	0	1	1	2	11	23	24	51	64	86	75	98	70	47	35	0	588
1967	0	0	0	1	0	1	1	3	7	25	32	55	59	78	84	63	43	17	0	469
1968	0	0	0	0	1	2	1	2	19	27	34	48	60	72	63	49	38	24	0	440
1964-68	0	0	0	1	4	5	10	24	72	135	221	285	335	400	397	343	219	112	0	2563
1969	0	0	0	0	1	1	6	10	17	24	50	69	80	88	78	66	46	21	1	558
1970	0	0	0	0	1	1	3	2	17	25	45	76	62	89	85	59	38	30	0	533
1971	0	0	0	1	1	1	3	7	15	33	44	75	82	90	80	66	37	33	0	568
1972	0	0	0	1	0	2	3	6	17	26	53	65	77	81	84	69	55	29	0	568
1973	0	0	0	1	2	0	4	12	13	29	58	77	80	88	108	77	59	45	2	655
1969-73	0	0	0	3	5	5	19	37	79	137	250	362	381	436	435	337	235	158	3	2882
1974	0	0	1	1	1	2	4	6	15	31	52	60	75	111	87	70	56	36	2	610
1975	0	0	0	0	1	1	2	6	16	30	53	51	97	98	86	64	47	27	4	583
1976	0	0	0	0	0	2	7	8	11	35	65	77	82	130	94	70	65	44	2	692
1977	0	0	0	1	0	2	3	10	14	33	73	67	97	91	91	81	55	33	0	651
1978	0	0	0	0	0	7	4	6	13	21	61	68	80	127	112	85	55	26	2	667
1974-78	0	0	1	2	2	14	20	36	69	150	304	323	431	557	470	370	278	166	10	3203
1979	0	0	0	0	3	1	3	5	16	28	58	101	100	125	109	99	60	31	3	742
1980	0	0	0	1	0	4	3	4	18	33	52	90	83	118	107	83	55	44	1	696
1981	0	0	0	0	1	2	4	8	15	31	54	89	97	108	112	84	48	35	1	689
1982	0	0	0	0	1	2	3	6	7	22	61	80	93	125	105	87	54	41	3	690
1983	1	0	0	0	0	2	3	8	9	19	48	99	101	120	110	91	61	35	4	711
1979-83	1	0	0	1	5	11	16	31	65	133	273	459	474	596	543	444	278	186	12	3528

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF RECTUM, RECTOSIGMOID JUNCTION AND ANUS (ICD9 154)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-72	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.9	0.0	1.3	0.8	4.1	15.9	29.4	51.5	63.2	107.8	130.3	171.8	220.5	110.3	17.4	16.7	2.0
1965	0.0	0.0	0.0	0.0	0.0	0.5	1.3	2.5	6.2	15.2	30.9	32.1	49.7	85.2	90.8	180.8	135.1	151.1	14.5	14.0	1.6
1966	0.0	0.0	0.0	0.0	0.4	0.5	0.9	4.6	9.9	12.3	28.5	43.5	71.9	80.6	139.2	150.2	180.8	250.0	17.1	16.6	1.9
1967	0.0	0.0	0.0	0.3	0.0	0.4	0.4	1.2	2.9	12.4	17.6	36.2	48.0	82.1	118.5	132.6	162.9	117.2	13.4	12.9	1.6
1968	0.0	0.0	0.0	0.0	0.4	0.8	0.4	0.8	7.9	12.9	18.6	30.5	47.7	74.1	88.6	102.5	141.8	160.0	12.3	12.0	1.4
1964-68	0.0	0.0	0.0	0.1	0.3	0.5	0.9	2.0	6.2	13.7	24.9	38.6	55.9	85.7	113.4	147.1	167.8	157.7	14.9	14.4	1.7
1969	0.0	0.0	0.0	0.0	0.3	0.4	2.6	4.2	7.0	11.8	27.3	42.5	62.1	88.0	108.6	137.2	169.1	133.8	15.2	14.8	1.8
1970	0.0	0.0	0.0	0.0	0.3	0.4	1.3	0.8	7.0	11.1	24.3	45.5	46.8	86.4	116.6	121.6	136.7	182.9	14.3	13.9	1.7
1971	0.0	0.0	0.0	0.3	0.3	0.3	1.2	2.9	6.2	14.2	23.3	44.0	60.2	84.8	107.5	135.0	131.2	190.8	14.8	14.5	1.7
1972	0.0	0.0	0.0	0.3	0.0	0.7	1.2	2.5	7.0	11.1	26.7	38.1	54.6	75.0	109.5	139.7	191.6	162.0	14.6	14.0	1.6
1973	0.0	0.0	0.0	0.3	0.6	0.0	1.5	5.1	5.3	12.3	27.8	45.4	55.0	79.9	135.7	154.6	202.7	247.3	16.5	15.8	1.8
1969-73	0.0	0.0	0.0	0.2	0.3	0.3	1.6	3.1	6.5	12.0	25.9	43.1	55.7	82.7	115.9	137.7	166.7	184.8	15.1	14.6	1.7
1974	0.0	0.0	0.2	0.3	0.3	0.6	1.5	2.5	6.1	13.1	23.8	35.6	49.9	98.3	105.7	138.3	191.8	194.6	15.0	14.5	1.7
1975	0.0	0.0	0.0	0.0	0.3	0.3	0.7	2.5	6.6	12.6	23.7	29.7	62.5	84.5	101.9	122.1	161.0	142.1	14.0	13.6	1.6
1976	0.0	0.0	0.0	0.0	0.0	0.6	2.4	3.2	4.6	14.6	28.6	43.0	52.1	107.9	108.8	128.7	218.9	226.8	16.2	15.7	1.8
1977	0.0	0.0	0.0	0.2	0.0	0.6	0.9	3.9	5.8	13.7	32.1	35.5	61.5	72.3	103.1	143.4	182.1	167.5	14.9	14.4	1.6
1978	0.0	0.0	0.0	0.0	0.0	2.0	1.2	2.3	5.4	8.7	26.6	34.2	50.8	97.6	123.5	144.3	178.6	130.0	14.8	14.3	1.7
1974-78	0.0	0.0	0.0	0.1	0.1	0.8	1.3	2.9	5.7	12.5	27.0	35.6	55.4	92.0	108.8	135.6	186.5	171.8	15.0	14.5	1.7
1979	0.0	0.0	0.0	0.0	0.8	0.3	0.9	1.8	6.6	11.6	25.2	48.4	63.8	92.9	116.3	161.8	188.7	151.2	16.1	15.5	1.8
1980	0.0	0.0	0.0	0.2	0.0	1.1	0.9	1.4	7.4	13.8	22.3	42.3	51.5	85.0	110.3	132.0	166.2	212.6	14.8	14.4	1.7
1981	0.0	0.0	0.0	0.0	0.3	0.6	1.1	2.7	6.1	13.1	22.9	41.2	57.5	76.5	111.2	129.8	140.8	165.1	14.3	13.9	1.7
1982	0.0	0.0	0.0	0.0	0.2	0.5	0.9	1.9	2.8	9.3	25.7	36.7	52.3	87.6	100.4	130.8	151.3	189.8	13.9	13.5	1.6
1983	0.3	0.0	0.0	0.0	0.0	0.5	0.9	2.4	3.4	8.1	20.2	44.8	53.8	84.3	100.8	132.8	161.4	157.0	13.9	13.4	1.6
1979-83	0.1	0.0	0.0	0.0	0.3	0.6	0.9	2.1	5.2	11.2	23.3	42.6	55.6	85.2	107.5	137.1	161.2	175.0	14.6	14.1	1.7

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF RECTUM, RECTOSIGMOID JUNCTION AND ANUS (ICD9 154)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	2	1	3	3	18	20	34	39	56	72	65	51	46	28	0	438
1965	0	0	0	0	0	1	2	5	13	30	31	31	44	52	54	49	26	34	0	372
1966	0	0	0	0	0	1	4	11	18	14	40	52	49	63	64	46	48	37	0	447
1967	0	0	0	0	1	5	2	9	8	29	27	45	57	46	57	54	39	26	0	405
1968	0	0	0	2	0	1	3	4	11	29	30	35	50	48	58	42	34	25	0	372
1964-68	0	0	0	[illegible]	3	9	14	32	68	122	162	202	256	281	298	242	193	150	0	2034
1969	0	0	0	0	1	1	2	8	14	30	47	49	50	59	65	50	38	32	0	446
1970	0	0	0	1	2	1	3	8	8	19	21	50	53	56	73	41	54	43	1	434
1971	0	0	0	0	1	0	7	9	11	22	29	52	55	57	57	51	32	37	0	420
1972	0	0	0	1	0	0	4	5	9	23	30	48	49	53	49	52	39	36	0	398
1973	0	0	0	0	1	3	2	8	20	27	34	55	56	62	79	63	52	35	2	499
1969-73	0	0	0	2	5	5	18	38	62	121	161	254	263	287	323	257	215	183	3	2197
1974	0	0	0	1	3	1	2	10	12	22	27	48	35	78	54	79	58	41	2	473
1975	0	0	0	0	1	2	4	2	12	32	45	43	56	56	72	57	31	58	2	473
1976	0	0	0	0	1	0	3	5	9	11	39	40	72	66	84	59	59	48	3	499
1977	0	0	0	0	1	1	0	3	8	24	46	65	81	71	77	66	56	49	3	551
1978	0	0	0	0	3	0	2	7	17	24	35	55	62	69	69	71	80	56	2	552
1974-78	0	0	0	1	9	4	11	27	58	113	192	251	306	340	356	332	284	252	12	2548
1979	0	0	0	0	2	0	8	8	11	23	49	64	65	71	79	86	58	48	3	575
1980	0	0	0	0	0	5	4	7	13	39	43	62	80	78	71	61	75	61	3	602
1981	0	0	0	1	0	1	6	4	7	20	35	53	63	79	82	89	62	52	0	554
1982	0	0	1	0	0	2	3	8	13	19	30	68	62	73	66	79	71	65	4	564
1983	0	0	0	0	1	3	5	6	5	24	47	39	66	81	69	62	72	63	3	546
1979-83	0	0	1	1	3	11	26	33	49	125	204	286	336	382	367	377	338	289	13	2841

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF RECTUM, RECTOSIGMOID JUNCTION AND ANUS (ICD9 154)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.9	0.5	1.4	1.3	8.1	10.8	20.7	28.3	47.5	70.9	78.3	87.0	132.9	127.3	13.3	11.3	1.3
1965	0.0	0.0	0.0	0.0	0.0	0.5	0.9	2.1	5.7	15.8	18.3	21.8	36.3	50.2	63.1	81.1	72.4	147.8	11.0	9.4	1.1
1966	0.0	0.0	0.0	0.0	0.0	0.5	1.8	4.7	7.6	7.2	22.9	35.6	39.4	59.4	72.2	74.3	129.0	156.1	12.8	10.9	1.2
1967	0.0	0.0	0.0	0.0	0.4	2.2	0.9	3.9	3.3	14.3	15.1	29.7	44.4	42.4	63.3	83.5	100.0	103.6	11.3	9.7	1.1
1968	0.0	0.0	0.0	0.6	0.0	0.4	1.3	1.7	4.6	13.7	16.5	22.3	37.9	43.1	63.6	62.7	83.5	94.3	10.1	8.7	1.0
1964-68	0.0	0.0	0.0	0.1	0.2	0.8	1.3	2.8	5.8	12.4	18.7	27.5	41.0	52.9	68.0	77.4	103.0	124.7	11.7	10.0	1.2
1969	0.0	0.0	0.0	0.0	0.3	0.4	0.9	3.5	5.9	13.5	25.6	30.1	36.8	51.5	70.2	72.7	89.6	113.1	11.7	10.1	1.2
1970	0.0	0.0	0.0	0.3	0.6	0.4	1.3	3.5	3.4	8.2	11.2	29.7	38.0	47.4	77.1	58.2	122.2	141.9	11.0	9.2	1.1
1971	0.0	0.0	0.0	0.0	0.3	0.0	3.0	3.9	4.7	9.3	15.1	30.0	38.2	46.3	58.7	70.7	69.3	113.8	10.4	9.0	1.0
1972	0.0	0.0	0.0	0.3	0.0	0.0	1.6	2.2	3.9	9.7	14.7	27.5	32.9	42.3	49.4	70.0	82.1	105.0	9.6	8.1	0.9
1973	0.0	0.0	0.0	0.0	0.3	0.9	0.8	3.5	8.6	11.4	15.7	31.4	36.4	48.2	77.6	82.9	106.1	97.5	11.7	9.9	1.2
1969-73	0.0	0.0	0.0	0.1	0.3	0.4	1.5	3.3	5.3	10.4	16.4	29.7	36.4	47.2	66.6	71.1	93.8	113.5	10.8	9.3	1.1
1974	0.0	0.0	0.0	0.3	0.8	0.3	0.7	4.3	5.1	9.3	11.9	27.2	21.9	58.9	51.5	101.4	115.8	108.2	10.7	8.8	1.0
1975	0.0	0.0	0.0	0.0	0.3	0.6	1.4	0.8	5.1	13.6	19.2	23.7	33.7	41.4	66.6	71.2	60.1	147.2	10.4	8.9	1.0
1976	0.0	0.0	0.0	0.0	0.3	0.0	1.0	2.1	3.8	4.7	16.4	21.0	42.7	47.0	75.7	71.3	110.7	115.9	10.6	8.9	1.1
1977	0.0	0.0	0.0	0.0	0.3	0.3	0.0	1.2	3.4	10.3	19.5	32.1	47.3	48.8	67.1	77.9	101.1	113.2	11.5	9.7	1.1
1978	0.0	0.0	0.0	0.0	0.8	0.0	0.6	2.7	7.3	10.3	14.8	25.6	36.0	45.9	58.1	81.2	139.9	124.7	11.2	9.2	1.0
1974-78	0.0	0.0	0.0	0.1	0.5	0.2	0.7	2.2	5.0	9.7	16.4	26.0	36.5	48.3	63.9	80.4	106.1	121.8	10.9	9.1	1.0
1979	0.0	0.0	0.0	0.0	0.5	0.0	2.3	2.9	4.7	9.9	21.0	28.4	37.5	45.8	64.3	95.3	97.3	101.9	11.3	9.5	1.1
1980	0.0	0.0	0.0	0.0	0.0	1.3	1.1	2.5	5.4	16.8	18.4	26.9	44.7	48.4	56.3	65.4	121.4	124.2	11.5	9.9	1.1
1981	0.0	0.0	0.0	0.3	0.0	0.3	1.6	1.4	2.9	8.6	15.0	22.7	33.6	48.1	62.9	92.4	97.0	101.0	10.3	8.5	1.0
1982	0.0	0.0	0.3	0.0	0.0	0.5	0.8	2.6	5.2	8.2	12.8	29.1	31.3	43.7	49.1	78.9	108.6	120.4	10.1	8.4	0.9
1983	0.0	0.0	0.0	0.0	0.2	0.8	1.4	1.8	1.9	10.3	20.1	16.6	31.4	48.5	49.4	59.8	105.7	110.7	9.6	8.1	0.9
1979-83	0.0	0.0	0.1	0.1	0.2	0.6	1.5	2.2	4.0	10.8	17.5	24.7	35.4	46.9	56.2	78.0	106.0	111.8	10.5	8.9	1.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF LIVER AND INTRAHEPATIC BILE DUCTS (ICD9 155)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	3	0	0	0	1	0	0	1	1	0	2	8	6	8	11	6	2	3	0	52
1965	2	3	0	1	0	0	1	0	2	0	3	11	5	7	9	7	4	2	0	57
1966	1	0	0	0	1	0	0	0	3	4	4	9	12	15	14	6	4	3	0	76
1967	4	1	1	0	0	0	0	1	1	4	7	8	6	11	11	9	5	4	0	73
1968	1	0	0	2	0	0	1	2	2	2	8	7	14	16	13	9	8	4	0	89
1964-68	11	4	1	3	2	0	2	4	9	10	24	43	43	57	58	37	23	16	0	347
1969	1	0	1	0	0	0	2	2	1	2	9	7	12	15	12	10	6	4	0	84
1970	0	0	0	0	1	0	1	0	0	1	6	11	9	10	7	8	5	1	0	68
1971	0	0	0	0	1	1	0	3	2	5	3	12	7	12	20	9	9	7	0	91
1972	2	0	1	0	0	1	1	0	4	6	9	10	17	22	21	11	7	5	0	117
1973	2	0	0	0	0	0	1	3	3	2	8	14	13	14	11	8	5	4	1	89
1969-73	5	0	2	0	2	2	5	8	10	16	35	54	58	73	71	46	32	21	1	441
1974	1	0	1	1	0	3	1	0	1	6	11	11	10	14	14	6	6	3	1	90
1975	2	0	0	1	0	2	1	2	2	5	4	12	12	12	11	6	6	2	1	81
1976	0	0	0	0	0	2	1	2	6	7	9	9	14	10	14	6	9	4	0	93
1977	2	0	0	0	0	1	0	4	0	6	12	10	15	19	22	11	14	4	0	120
1978	0	0	0	0	1	0	1	1	5	3	7	16	11	13	9	12	2	5	0	86
1974-78	5	0	1	2	1	8	4	9	14	27	43	58	62	68	70	41	37	18	2	470
1979	3	0	0	1	1	0	1	4	2	3	5	12	12	13	21	12	2	5	0	97
1980	4	0	0	0	1	0	2	0	1	8	6	16	9	18	12	12	4	6	0	99
1981	2	0	0	0	1	1	2	3	4	4	8	11	16	14	11	15	7	5	0	104
1982	2	0	0	1	0	1	2	0	2	2	4	15	14	21	21	13	8	3	0	109
1983	1	1	1	0	3	0	3	1	3	7	7	16	15	19	14	10	7	7	0	115
1979-83	12	1	1	2	6	2	10	8	12	24	30	70	66	85	79	62	28	26	0	524

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF LIVER AND INTRAHEPATIC BILE DUCTS (ICD9 155)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.8	0.0	0.0	0.0	0.5	0.0	0.0	0.4	0.5	0.0	1.2	5.7	5.3	8.9	16.1	13.2	7.9	22.1	1.6	1.6	0.2
1965	0.5	0.8	0.0	0.3	0.0	0.0	0.4	0.0	0.9	0.0	1.7	7.7	4.3	7.7	13.0	15.3	15.4	14.4	1.7	1.6	0.2
1966	0.3	0.0	0.0	0.0	0.4	0.0	0.0	0.0	1.3	2.0	2.2	6.1	10.0	16.1	19.9	12.9	15.4	21.4	2.2	2.2	0.3
1967	1.1	0.3	0.3	0.0	0.0	0.0	0.0	0.4	0.4	2.0	3.8	5.3	4.9	11.6	15.5	18.9	18.9	27.6	2.1	2.0	0.2
1968	0.3	0.0	0.0	0.6	0.0	0.0	0.4	0.8	0.8	1.0	4.4	4.5	11.1	16.5	18.3	18.8	29.9	26.7	2.5	2.4	0.3
1964-68	0.6	0.2	0.1	0.2	0.2	0.0	0.2	0.3	0.8	1.0	2.7	5.8	7.2	12.2	16.6	15.9	17.6	22.5	2.0	2.0	0.2
1969	0.3	0.0	0.3	0.0	0.0	0.0	0.9	0.8	0.4	0.9	4.9	4.3	9.3	15.0	16.7	20.8	22.1	25.5	2.3	2.3	0.3
1970	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.0	0.0	0.4	3.2	6.6	6.8	9.7	9.6	16.5	18.0	6.1	1.6	1.5	0.2
1971	0.0	0.0	0.0	0.0	0.3	0.3	0.0	1.3	0.8	2.2	1.6	7.0	5.1	11.3	26.9	18.4	31.9	40.5	2.4	2.3	0.3
1972	0.6	0.0	0.2	0.0	0.0	0.3	0.4	0.0	1.6	2.6	4.5	5.9	12.1	20.4	27.4	22.3	24.4	27.9	3.0	3.0	0.4
1973	0.6	0.0	0.0	0.0	0.0	0.0	0.4	1.3	1.2	0.9	3.8	8.3	8.9	12.7	13.8	16.1	17.2	22.0	2.2	2.2	0.3
1969-73	0.3	0.0	0.1	0.0	0.1	0.1	0.4	0.7	0.8	1.4	3.6	6.4	8.5	13.8	18.9	18.8	22.7	24.6	2.3	2.3	0.3
1974	0.3	0.0	0.2	0.3	0.0	0.9	0.4	0.0	0.4	2.5	5.0	6.5	6.7	12.4	17.0	11.9	20.5	16.2	2.2	2.1	0.3
1975	0.6	0.0	0.0	0.2	0.0	0.6	0.3	0.8	0.8	2.1	1.8	7.0	7.7	10.3	13.0	11.5	20.5	10.5	1.9	1.9	0.2
1976	0.0	0.0	0.0	0.0	0.0	0.6	0.3	0.8	2.5	2.9	4.0	5.0	8.9	8.3	16.2	11.0	30.3	20.6	2.2	2.1	0.2
1977	0.6	0.0	0.0	0.0	0.0	0.3	0.0	1.6	0.0	2.5	5.3	5.3	9.5	15.1	24.9	19.5	46.4	20.3	2.8	2.7	0.3
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.4	2.1	1.2	3.1	8.0	7.0	10.0	9.9	20.4	6.5	25.0	1.9	1.9	0.2
1974-78	0.3	0.0	0.0	0.1	0.1	0.5	0.3	0.7	1.2	2.3	3.8	6.4	8.0	11.2	16.2	15.0	24.8	18.6	2.2	2.2	0.3
1979	1.0	0.0	0.0	0.2	0.3	0.0	0.3	1.5	0.8	1.2	2.2	5.8	7.7	9.7	22.4	19.6	6.3	24.4	2.1	2.1	0.3
1980	1.3	0.0	0.0	0.0	0.3	0.0	0.6	0.0	0.4	3.4	2.6	7.5	5.6	13.0	12.4	19.1	12.1	29.0	2.1	2.1	0.2
1981	0.7	0.0	0.0	0.0	0.3	0.3	0.6	1.0	1.6	1.7	3.4	5.1	9.5	9.9	10.9	23.2	20.5	23.6	2.2	2.1	0.2
1982	0.7	0.0	0.0	0.2	0.0	0.3	0.6	0.0	0.8	0.8	1.7	6.9	7.9	14.7	20.1	19.5	22.4	13.9	2.2	2.1	0.3
1983	0.3	0.3	0.3	0.0	0.7	0.0	0.9	0.3	1.1	3.0	3.0	7.2	8.0	13.4	12.8	14.6	18.5	31.4	2.3	2.3	0.3
1979-83	0.8	0.1	0.1	0.1	0.3	0.1	0.6	0.5	1.0	2.0	2.6	6.5	7.7	12.2	15.6	19.1	16.2	24.5	2.2	2.2	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF LIVER AND INTRAHEPATIC BILE DUCTS (ICD9 155)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	4	0	0	0	0	1	0	1	1	1	3	3	6	5	9	7	12	8	0	61
1965	1	0	0	0	0	0	2	0	1	5	1	4	5	9	6	6	7	5	0	52
1966	1	0	0	0	0	1	1	0	1	1	0	7	4	2	7	7	9	6	0	47
1967	1	0	0	0	2	1	0	0	3	2	4	7	7	7	7	4	9	4	0	58
1968	1	0	0	0	1	0	0	0	4	0	2	5	5	5	5	4	2	5	0	39
1964-68	8	0	0	0	3	3	3	1	10	9	10	26	27	28	34	28	39	28	0	257
1969	1	0	0	0	1	0	0	0	1	5	2	4	4	6	5	6	10	1	0	46
1970	1	1	1	1	1	0	0	1	1	4	4	4	4	3	8	5	6	1	0	46
1971	1	0	0	0	2	1	1	1	3	0	7	2	4	3	6	13	7	5	0	56
1972	1	0	0	0	0	0	1	0	1	1	4	5	6	8	5	10	4	8	0	54
1973	1	0	0	1	1	0	0	1	0	4	2	4	5	6	11	6	12	11	0	65
1969-73	5	1	1	2	5	1	2	3	6	14	19	19	23	26	35	40	39	26	0	267
1974	3	0	0	1	1	1	0	1	3	3	6	4	6	4	11	15	10	5	1	75
1975	0	0	0	0	0	1	1	1	0	2	6	4	13	6	6	6	3	11	0	60
1976	3	0	0	0	0	0	0	1	2	5	3	3	10	7	4	10	10	4	0	62
1977	1	0	1	0	0	1	3	1	0	5	6	8	11	12	10	11	8	11	1	90
1978	0	1	0	0	0	0	0	1	0	1	1	4	4	7	9	12	5	7	1	53
1974-78	7	1	1	1	1	3	4	5	5	16	22	23	44	36	40	54	36	38	3	340
1979	0	0	0	0	0	0	2	1	0	1	4	4	5	6	8	13	5	5	0	54
1980	1	0	0	1	0	1	2	0	1	1	2	5	5	7	10	8	5	6	0	55
1981	1	0	0	0	2	0	1	2	1	1	2	3	6	13	11	10	4	11	1	69
1982	1	1	0	0	0	2	1	0	4	2	2	5	8	10	11	7	9	4	1	68
1983	0	0	0	1	0	3	0	0	0	3	5	8	6	9	4	11	8	2	0	60
1979-83	3	1	0	2	2	6	6	3	6	8	15	25	30	45	44	49	31	28	2	306

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF LIVER AND INTRAHEPATIC BILE DUCTS (ICD9 155)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	1.1	0.0	0.0	0.0	0.0	0.5	0.0	0.4	0.5	0.5	1.8	2.2	5.1	4.9	10.8	11.9	34.7	36.4	1.9	1.5	0.1
1965	0.3	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.4	2.6	0.6	2.8	4.1	8.7	7.0	9.9	19.5	21.7	1.5	1.3	0.1
1966	0.3	0.0	0.0	0.0	0.0	0.5	0.5	0.0	0.4	0.5	0.0	4.8	3.2	1.9	7.9	11.3	24.2	25.3	1.4	1.0	0.1
1967	0.3	0.0	0.0	0.0	0.8	0.4	0.0	0.0	1.2	1.0	2.2	4.6	5.5	6.5	7.8	6.2	23.1	15.9	1.6	1.4	0.2
1968	0.3	0.0	0.0	0.0	0.4	0.0	0.0	0.0	1.7	0.0	1.1	3.2	3.8	4.5	5.5	6.0	4.9	18.9	1.1	0.9	0.1
1964-68	0.4	0.0	0.0	0.0	0.2	0.3	0.3	0.1	0.9	0.9	1.2	3.5	4.3	5.3	7.8	9.0	20.8	23.3	1.5	1.2	0.1
1969	0.3	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	2.3	1.1	2.5	2.9	5.2	5.4	8.7	23.6	3.5	1.2	1.0	0.1
1970	0.3	0.3	0.3	0.3	0.3	0.0	0.0	0.8	0.4	1.7	2.1	2.4	2.9	2.5	8.4	7.1	13.6	3.3	1.2	1.0	0.1
1971	0.3	0.0	0.0	0.0	0.6	0.4	0.4	0.4	1.3	0.0	3.6	1.2	2.8	2.5	6.2	18.0	15.2	15.4	1.4	1.1	0.1
1972	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.4	2.0	2.9	4.0	6.4	5.0	13.5	8.4	23.3	1.3	1.1	0.1
1973	0.3	0.0	0.0	0.3	0.3	0.0	0.0	0.4	0.0	1.7	0.9	2.3	3.2	4.7	10.8	7.9	24.5	30.6	1.5	1.2	0.1
1969-73	0.3	0.1	0.1	0.1	0.3	0.1	0.2	0.3	0.5	1.2	1.9	2.2	3.2	4.3	7.2	11.1	17.0	16.1	1.3	1.1	0.1
1974	1.0	0.0	0.0	0.3	0.3	0.3	0.0	0.4	1.3	1.3	2.6	2.3	3.8	3.0	10.5	19.3	20.0	13.2	1.7	1.4	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.4	0.0	0.9	2.6	2.2	7.8	4.4	5.6	7.5	5.8	27.9	1.3	1.1	0.1
1976	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	2.1	1.3	1.6	5.9	5.0	3.6	12.1	18.8	9.7	1.4	1.2	0.1
1977	0.3	0.0	0.3	0.0	0.0	0.3	1.0	0.4	0.0	2.1	2.5	4.0	6.4	8.2	8.7	13.0	14.4	25.4	1.9	1.6	0.2
1978	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.4	1.9	2.3	4.7	7.6	13.7	8.7	15.6	1.0	0.8	0.1
1974-78	0.5	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.4	1.4	1.9	2.4	5.3	5.1	7.2	13.1	13.5	18.4	1.4	1.2	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.4	0.0	0.4	1.7	1.8	2.9	3.9	6.5	14.4	8.4	10.6	1.1	0.8	0.1
1980	0.3	0.0	0.0	0.2	0.0	0.3	0.6	0.0	0.4	0.4	0.9	2.2	2.8	4.3	7.9	8.6	8.1	12.2	1.0	0.9	0.1
1981	0.3	0.0	0.0	0.0	0.5	0.0	0.3	0.7	0.4	0.4	0.9	1.3	3.2	7.9	8.4	10.4	6.3	21.4	1.3	1.1	0.1
1982	0.3	0.3	0.0	0.0	0.0	0.5	0.3	0.0	1.6	0.9	0.9	2.1	4.0	6.0	8.2	7.0	13.8	7.4	1.2	1.1	0.1
1983	0.0	0.0	0.0	0.3	0.0	0.8	0.0	0.0	0.0	1.3	2.1	3.4	2.9	5.4	2.9	10.6	11.7	3.5	1.1	0.9	0.1
1979-83	0.2	0.1	0.0	0.1	0.1	0.3	0.3	0.2	0.5	0.7	1.3	2.2	3.2	5.5	6.7	10.1	9.7	10.8	1.1	1.0	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF GALLBLADDER AND EXTRAHEPATIC BILE DUCTS (ICD9 156)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	1	2	1	1	5	6	5	16	13	8	4	0	62
1965	0	0	0	0	0	0	1	0	1	3	6	6	8	13	15	10	9	4	0	76
1966	0	0	0	0	0	0	0	0	0	1	5	4	6	8	15	4	6	2	0	51
1967	0	0	0	0	0	0	0	1	3	3	4	8	12	9	8	13	6	3	0	70
1968	0	0	0	0	0	0	0	1	2	0	5	3	7	11	16	9	5	4	0	64
1964-68	0	0	0	0	0	0	1	3	8	8	21	26	39	46	70	49	34	17	0	322
1969	0	0	0	0	0	0	0	1	1	0	4	3	10	6	12	12	11	4	0	64
1970	0	0	0	0	0	0	1	0	1	4	10	6	10	13	14	7	9	3	0	78
1971	0	0	0	0	0	0	0	1	1	5	2	3	10	12	15	8	3	3	0	63
1972	0	0	0	0	0	0	0	1	1	1	3	12	10	12	13	4	6	4	0	67
1973	0	0	0	0	0	0	0	1	5	3	5	6	11	11	13	13	8	4	0	80
1969-73	0	0	0	0	0	0	1	4	9	13	24	30	51	54	67	44	37	18	0	352
1974	0	0	0	0	0	0	0	2	1	3	5	3	10	13	10	6	5	10	0	68
1975	0	0	0	0	0	1	0	1	3	0	2	2	9	10	14	8	7	5	0	62
1976	0	0	0	0	0	0	0	1	2	3	11	6	9	13	14	11	6	8	0	84
1977	0	0	0	0	0	2	0	0	2	7	9	9	15	16	11	13	8	6	0	98
1978	0	0	1	0	0	0	1	0	2	5	7	6	11	17	14	12	13	7	1	97
1974-78	0	0	1	0	0	3	1	4	10	18	34	26	54	69	63	50	39	36	1	409
1979	0	0	0	1	1	1	2	1	3	1	4	13	13	23	17	11	12	9	0	112
1980	0	0	0	0	0	0	0	0	2	3	7	9	16	17	17	14	6	9	0	100
1981	0	0	0	0	0	0	0	0	1	1	8	7	19	24	14	15	9	4	1	103
1982	0	0	0	0	1	0	0	2	2	3	5	12	8	15	14	20	7	10	0	99
1983	0	0	0	0	0	0	1	0	2	0	5	11	13	15	14	13	8	8	0	90
1979-83	0	0	0	1	2	1	3	3	10	8	29	52	69	94	76	73	42	40	1	504

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF GALLBLADDER AND EXTRAHEPATIC BILE DUCTS (ICD9 156)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.5	0.6	3.6	5.3	5.6	23.4	28.6	31.5	29.4	1.9	1.7	0.2
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	1.6	3.4	4.2	6.9	14.2	21.6	21.8	34.7	28.8	2.3	2.2	0.3
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	2.8	2.7	5.0	8.6	21.3	8.6	23.1	14.3	1.5	1.4	0.2
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	1.5	2.2	5.3	9.8	9.5	11.3	27.4	22.7	20.7	2.0	1.9	0.2
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	0.0	2.7	1.9	5.6	11.3	22.5	18.8	18.7	26.7	1.8	1.7	0.2
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.7	0.8	2.4	3.5	6.5	9.9	20.0	21.0	26.1	25.9	1.9	1.8	0.2
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	2.2	1.8	7.8	6.0	16.7	24.9	40.4	25.5	1.8	1.6	0.2
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	1.8	5.4	3.6	7.6	12.6	19.2	14.4	32.4	18.3	2.1	2.0	0.3
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.2	1.1	1.8	7.3	11.3	20.2	16.4	10.6	17.3	1.6	1.6	0.2
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	1.5	7.0	7.1	11.1	16.9	8.1	20.9	22.3	1.7	1.7	0.2
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.1	1.3	2.4	3.5	7.6	10.0	16.3	26.1	27.5	22.0	2.0	1.9	0.2
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.7	1.1	2.5	3.6	7.5	10.2	17.8	18.0	26.2	21.1	1.9	1.8	0.2
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	1.3	2.3	1.8	6.7	11.5	12.2	11.9	17.1	54.1	1.7	1.7	0.2
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	1.2	0.0	0.9	1.2	5.8	8.6	16.6	15.3	24.0	26.3	1.5	1.4	0.2
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.3	4.8	3.4	5.7	10.8	16.2	20.2	20.2	41.2	2.0	1.9	0.2
1977	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.8	2.9	4.0	4.8	9.5	12.7	12.5	23.0	26.5	30.5	2.2	2.2	0.2
1978	0.0	0.0	0.3	0.0	0.0	0.0	0.3	0.0	0.8	2.1	3.1	3.0	7.0	13.1	15.4	20.4	42.2	35.0	2.2	2.1	0.2
1974-78	0.0	0.0	0.0	0.0	0.0	0.2	0.1	0.3	0.8	1.5	3.0	2.9	6.9	11.4	14.6	18.3	26.2	37.3	1.9	1.9	0.2
1979	0.0	0.0	0.0	0.2	0.3	0.3	0.6	0.4	1.2	0.4	1.7	6.2	8.3	17.1	18.1	18.0	37.7	43.9	2.4	2.4	0.3
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.3	3.0	4.2	9.9	12.2	17.5	22.3	18.1	43.5	2.1	2.1	0.2
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	3.4	3.2	11.3	17.0	13.9	23.2	26.4	18.9	2.1	2.0	0.2
1982	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.6	0.8	1.3	2.1	5.5	4.5	10.5	13.4	30.1	19.6	46.3	2.0	1.9	0.2
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.8	0.0	2.1	5.0	6.9	10.5	12.8	19.0	21.2	35.9	1.8	1.7	0.2
1979-83	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.2	0.8	0.7	2.5	4.8	8.1	13.4	15.0	22.5	24.3	37.6	2.1	2.0	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF GALLBLADDER AND EXTRAHEPATIC BILE DUCTS (ICD9 156)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	2	2	4	9	17	20	31	25	15	2	0	127
1965	0	0	0	0	0	0	0	0	2	4	5	9	13	20	22	21	11	9	0	116
1966	0	0	0	0	0	0	1	1	1	5	7	10	16	26	13	18	16	9	0	123
1967	0	0	0	0	0	0	0	2	2	1	6	11	16	25	15	21	18	4	0	121
1968	0	0	0	0	0	0	0	1	1	5	2	12	12	15	27	23	12	8	0	116
1964-68	0	0	0	0	0	0	1	4	8	17	24	51	74	104	108	108	72	32	0	603
1969	0	0	0	0	0	0	1	0	3	1	9	13	8	20	17	23	12	9	0	116
1970	0	0	0	0	0	0	0	0	1	1	4	8	18	18	26	21	19	12	0	128
1971	0	0	0	0	1	1	0	1	1	1	7	5	16	18	15	18	17	12	0	113
1972	0	0	0	0	0	0	1	0	1	3	7	9	21	23	25	21	11	11	1	134
1973	0	0	0	0	0	1	0	3	3	1	11	8	18	18	33	23	17	11	0	147
1969-73	0	0	0	0	1	2	2	4	9	7	38	43	81	97	116	106	76	55	1	638
1974	0	0	0	0	0	0	1	0	2	3	5	4	16	17	23	20	12	14	0	117
1975	0	0	0	0	0	0	0	0	1	6	11	11	11	16	19	21	11	16	0	123
1976	0	0	0	0	0	1	0	0	1	6	2	14	14	17	15	23	22	13	0	124
1977	0	0	0	0	0	0	0	1	3	4	9	13	11	15	22	21	21	9	0	129
1978	0	0	0	0	0	0	0	0	4	4	9	18	23	18	29	20	17	21	1	164
1974-78	0	0	0	0	0	1	1	1	11	23	36	60	75	79	108	105	83	73	1	657
1979	0	0	0	0	0	0	0	0	1	7	5	11	19	18	19	28	25	17	0	150
1980	0	0	0	0	0	0	0	1	4	3	7	12	24	19	26	23	28	13	0	160
1981	0	0	0	0	0	1	0	0	1	3	8	11	11	28	31	32	22	17	0	165
1982	0	1	0	0	0	0	0	2	1	3	6	14	18	21	26	30	30	20	0	172
1983	0	0	0	0	0	1	0	1	1	2	7	12	14	19	22	27	26	21	0	153
1979-83	0	1	0	0	0	2	0	4	8	18	33	60	86	105	124	140	131	88	0	800

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF GALLBLADDER AND EXTRAHEPATIC BILE DUCTS (ICD9 156)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.1	2.4	6.5	14.4	19.7	37.3	42.7	43.4	9.1	3.8	3.1	0.4
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	2.1	2.9	6.3	10.7	19.3	25.7	34.8	30.6	39.1	3.4	2.8	0.3
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.4	0.4	2.6	4.0	6.8	12.9	24.5	14.7	29.1	43.0	38.0	3.5	2.9	0.3
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.8	0.5	3.4	7.3	12.5	23.0	16.6	32.5	46.2	15.9	3.4	2.7	0.3
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.4	1.1	7.6	9.1	11.7	29.6	34.3	29.5	30.2	3.1	2.5	0.3
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.7	1.7	2.8	6.9	11.9	19.6	24.6	34.5	38.4	26.6	3.4	2.8	0.3
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.3	0.5	4.9	8.0	5.9	17.5	18.4	33.4	28.3	31.8	3.0	2.5	0.3
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.1	4.7	12.9	15.2	27.5	29.8	43.0	39.6	3.2	2.6	0.3
1971	0.0	0.0	0.0	0.0	0.3	0.4	0.0	0.4	0.4	0.4	3.6	2.9	11.1	14.8	15.4	25.0	36.8	36.9	2.8	2.2	0.2
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	1.3	3.4	5.2	14.1	18.4	25.2	28.3	23.2	32.1	3.2	2.7	0.3
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.0	1.3	1.3	0.4	5.1	4.6	11.7	14.0	32.4	30.3	34.7	30.6	3.4	2.8	0.4
1969-73	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.3	0.8	0.6	3.9	5.0	11.2	15.9	23.9	29.3	33.1	34.1	3.1	2.6	0.3
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	1.3	2.2	2.3	10.0	12.8	21.9	25.7	24.0	36.9	2.6	2.1	0.3
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.6	4.7	6.1	6.6	11.8	17.6	26.2	21.3	40.6	2.7	2.2	0.2
1976	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	2.6	0.8	7.4	8.3	9.3	13.5	27.8	41.3	31.4	2.6	2.1	0.2
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	1.7	3.8	6.4	6.4	10.3	19.2	24.8	37.9	20.8	2.7	2.1	0.2
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	1.7	3.8	8.4	13.3	12.0	24.4	22.9	29.7	46.8	3.3	2.7	0.3
1974-78	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.9	2.0	3.1	6.2	9.0	11.2	19.4	25.4	31.0	35.3	2.8	2.3	0.3
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	3.0	2.1	4.9	11.0	11.6	15.5	31.0	41.9	36.1	2.9	2.3	0.2
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	1.3	3.0	5.2	13.4	11.8	20.6	24.7	45.3	26.5	3.0	2.5	0.3
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	1.3	3.4	4.7	5.9	17.0	23.8	33.2	34.4	33.0	3.0	2.4	0.3
1982	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.6	0.4	1.3	2.6	6.0	9.1	12.6	19.3	30.0	45.9	37.0	3.0	2.4	0.3
1983	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.4	0.9	3.0	5.1	6.7	11.4	15.8	26.0	38.2	36.9	2.6	2.0	0.2
1979-83	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.3	0.6	1.6	2.8	5.2	9.1	12.9	19.0	28.9	41.1	34.0	2.9	2.3	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF PANCREAS (ICD9 157)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	3	4	6	26	26	35	42	70	67	41	30	20	0	370
1965	0	0	0	0	0	0	1	2	4	17	19	40	44	41	49	35	31	9	0	292
1966	0	0	0	0	1	0	2	5	6	16	31	30	47	54	56	32	26	15	0	321
1967	0	0	0	0	0	1	0	0	9	14	21	29	48	63	52	30	31	7	0	306
1968	0	0	0	0	0	0	1	8	10	19	21	36	41	55	50	35	22	20	0	318
1964-68	0	0	0	0	1	1	7	19	35	92	118	170	222	283	274	173	140	71	0	1606
1969	0	0	0	0	0	0	1	3	14	15	27	46	56	69	55	55	41	20	0	402
1970	0	0	0	0	0	1	2	3	10	20	27	36	57	69	78	47	19	23	0	392
1971	0	0	0	0	1	1	1	2	9	22	24	55	50	72	56	45	34	23	0	395
1972	0	0	0	1	1	2	1	5	13	21	37	41	57	62	55	66	38	19	0	419
1973	0	0	0	0	0	0	3	2	7	11	22	41	57	46	79	34	35	23	1	361
1969-73	0	0	0	1	2	4	8	15	53	89	137	219	277	318	323	247	167	108	1	1969
1974	0	0	0	0	2	3	2	5	10	17	36	41	56	57	60	38	27	31	0	388
1975	0	0	0	0	1	3	0	5	9	14	27	43	48	59	59	43	33	17	0	361
1976	0	0	0	1	1	0	1	2	7	19	28	38	74	70	73	60	27	21	1	423
1977	0	0	0	0	0	0	2	3	12	19	43	45	62	65	72	67	23	27	0	440
1978	0	0	1	0	0	2	1	4	8	14	37	45	68	66	71	52	29	23	0	421
1974-78	0	0	1	1	4	5	6	19	46	83	171	212	308	317	335	260	139	119	1	2033
1979	0	0	0	0	2	0	0	4	5	24	36	54	69	62	63	45	35	24	0	423
1980	0	0	0	0	1	0	1	7	9	24	37	54	52	67	62	40	42	27	0	423
1981	1	0	0	1	1	2	1	4	5	7	23	47	50	56	81	45	48	19	0	391
1982	0	0	0	0	0	1	2	3	6	18	30	54	66	78	91	62	43	17	0	463
1983	0	0	0	0	0	1	4	1	11	15	25	46	84	68	80	54	43	21	0	455
1979-83	1	0	0	1	4	4	8	19	36	80	151	255	323	331	377	246	211	108	0	2155

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF PANCREAS (ICD9 157)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	1.3	1.7	2.8	13.8	15.3	25.1	36.9	77.8	98.1	90.3	118.1	147.1	11.2	10.9	1.4
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.8	8.9	10.9	27.9	37.7	44.8	70.6	76.3	119.7	64.7	8.7	8.3	1.0
1966	0.0	0.0	0.0	0.0	0.4	0.0	0.9	2.1	2.6	8.2	17.3	20.4	39.3	58.0	79.5	68.7	100.0	107.1	9.3	9.2	1.1
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	3.8	7.0	11.5	19.1	39.1	66.3	73.3	63.2	117.4	48.3	8.7	8.5	1.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.4	3.3	4.2	9.1	11.5	22.9	32.6	56.6	70.3	73.2	82.1	133.3	8.9	8.7	1.0
1964-68	0.0	0.0	0.0	0.0	0.1	0.1	0.6	1.6	3.0	9.3	13.3	23.0	37.1	60.6	78.3	74.2	107.3	100.0	9.3	9.1	1.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	5.8	6.9	14.7	28.3	43.4	69.0	76.6	114.3	150.7	127.4	11.0	10.6	1.2
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.3	4.1	8.8	14.6	21.5	43.1	67.0	107.0	96.9	68.3	140.2	10.5	10.4	1.3
1971	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.8	3.7	9.5	12.7	32.3	36.7	67.9	75.3	92.0	120.6	132.9	10.3	10.0	1.2
1972	0.0	0.0	0.0	0.3	0.3	0.7	0.4	2.1	5.3	9.0	18.6	24.0	40.4	57.4	71.7	133.6	132.4	106.1	10.8	10.3	1.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.8	2.9	4.7	10.5	24.2	39.2	41.8	99.2	68.3	120.3	126.4	9.1	8.8	1.1
1969-73	0.0	0.0	0.0	0.1	0.1	0.3	0.7	1.3	4.4	7.8	14.2	26.1	40.5	60.3	86.0	100.9	118.4	126.3	10.3	10.0	1.2
1974	0.0	0.0	0.0	0.0	0.6	0.9	0.7	2.1	4.1	7.2	16.4	24.3	37.3	50.5	72.9	75.1	92.5	167.6	9.5	9.3	1.1
1975	0.0	0.0	0.0	0.0	0.3	0.9	0.0	2.1	3.7	5.9	12.1	25.0	30.9	50.9	69.9	82.1	113.0	89.5	8.7	8.4	1.0
1976	0.0	0.0	0.0	0.2	0.3	0.0	0.3	0.8	2.9	7.9	12.3	21.2	47.0	58.1	84.5	110.3	90.9	108.2	9.9	9.6	1.2
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.6	1.2	5.0	7.9	18.9	23.8	39.3	51.6	81.5	118.6	76.2	137.1	10.1	9.8	1.1
1978	0.0	0.0	0.3	0.0	0.0	0.6	0.3	1.5	3.4	5.8	16.2	22.6	43.2	50.7	78.3	88.3	94.2	115.0	9.4	9.2	1.1
1974-78	0.0	0.0	0.0	0.0	0.2	0.5	0.4	1.5	3.8	6.9	15.2	23.4	39.6	52.4	77.5	95.3	93.2	123.2	9.5	9.3	1.1
1979	0.0	0.0	0.0	0.0	0.5	0.0	0.0	1.5	2.1	10.0	15.6	25.9	44.0	46.1	67.2	73.5	110.1	117.1	9.2	9.0	1.1
1980	0.0	0.0	0.0	0.0	0.3	0.0	0.3	2.5	3.7	10.1	15.9	25.4	32.3	48.3	63.9	63.4	126.9	130.4	9.0	8.8	1.0
1981	0.3	0.0	0.0	0.2	0.3	0.6	0.3	1.4	2.0	3.0	9.8	21.8	29.6	39.7	80.4	69.6	140.8	89.6	8.1	7.7	9.9
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.6	1.0	2.4	4.2	12.7	24.7	37.1	54.7	87.0	93.2	120.4	78.7	9.3	8.9	1.1
1983	0.0	0.0	0.0	0.0	0.0	0.3	1.1	0.3	4.2	6.4	10.5	20.8	45.8	47.8	73.3	78.8	113.8	94.2	8.9	8.7	1.0
1979-83	0.1	0.0	0.0	0.0	0.2	0.2	0.5	1.3	2.9	6.7	12.9	23.7	37.9	47.3	74.6	76.0	122.3	101.6	8.9	8.6	1.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF PANCREAS (ICD9 157)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	2	1	7	14	24	17	27	35	47	36	40	15	0	265
1965	0	0	0	1	0	0	1	3	3	2	11	15	24	40	32	37	23	18	0	210
1966	0	1	0	0	0	0	0	2	4	12	10	23	30	25	37	44	24	21	0	233
1967	0	0	0	0	0	0	0	2	2	8	19	21	23	26	32	49	31	20	0	233
1968	0	0	0	0	0	0	2	1	3	12	15	18	30	38	36	45	29	24	0	253
1964-68	0	1	0	1	0	0	5	9	19	48	79	94	134	164	184	211	147	98	0	1194
1969	0	0	0	0	0	0	0	2	9	10	23	29	42	32	46	51	39	31	0	314
1970	0	0	0	0	1	1	2	7	3	12	13	21	28	36	27	45	35	25	0	256
1971	0	0	0	0	0	0	2	2	10	9	15	33	36	42	46	35	26	19	0	275
1972	0	0	0	0	0	0	0	0	6	12	16	29	38	30	47	50	59	37	0	304
1973	0	0	0	0	0	0	0	2	7	11	11	33	45	36	54	51	30	42	0	322
1969-73	0	0	0	0	1	1	4	13	35	54	78	145	189	176	220	232	169	154	0	1471
1974	0	0	0	0	1	0	1	4	5	7	13	22	40	52	40	41	44	33	1	304
1975	0	0	0	0	0	0	1	1	5	18	19	26	27	42	47	47	34	50	0	317
1976	1	0	0	1	0	1	1	0	5	11	19	19	37	58	58	42	45	27	0	325
1977	1	0	0	0	2	0	3	2	6	16	29	28	37	54	48	42	40	38	0	346
1978	0	0	0	0	0	0	0	0	4	9	19	23	34	38	70	48	43	40	0	328
1974-78	2	0	0	1	3	1	6	7	25	61	99	118	175	244	263	220	206	188	1	1620
1979	0	0	0	0	0	1	0	1	8	9	18	28	42	55	63	68	50	51	0	394
1980	0	0	0	1	0	0	2	2	8	7	24	26	37	49	55	63	39	48	0	361
1981	0	1	0	0	0	0	1	0	5	13	27	39	42	67	63	57	50	49	0	414
1982	0	0	0	0	0	0	3	4	8	8	24	25	40	53	72	62	40	47	0	386
1983	0	0	0	0	0	0	1	2	5	13	17	29	41	56	73	47	50	39	0	373
1979-83	0	1	0	1	0	1	7	9	34	50	110	147	202	280	326	297	229	234	0	1928

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF PANCREAS (ICD9 157)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	3.2	7.6	14.6	12.3	22.9	34.4	56.6	61.4	115.6	68.2	8.0	6.6	0.8
1965	0.0	0.0	0.0	0.4	0.0	0.0	0.5	1.3	1.3	1.1	6.5	10.6	19.8	38.6	37.4	61.3	64.1	78.3	6.2	5.0	0.6
1966	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.9	1.7	6.1	5.7	15.7	24.1	23.6	41.8	71.1	64.5	88.6	6.7	5.4	0.6
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.8	3.9	10.6	13.9	17.9	24.0	35.5	75.7	79.5	79.7	6.5	5.1	0.5
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	1.2	5.7	8.3	11.5	22.7	34.1	39.5	67.2	71.3	90.6	6.8	5.6	0.6
1964-68	0.0	0.1	0.0	0.1	0.0	0.0	0.5	0.8	1.6	4.9	9.1	12.8	21.5	30.9	42.0	67.5	78.4	81.5	6.9	5.5	0.6
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	3.8	4.5	12.5	17.8	31.0	27.9	49.7	74.1	92.0	109.5	8.2	6.7	0.7
1970	0.0	0.0	0.0	0.0	0.3	0.4	0.9	3.1	1.3	5.2	7.0	12.5	20.1	30.5	28.5	63.8	79.2	82.5	6.5	5.3	0.5
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.9	4.3	3.8	7.8	19.1	25.0	34.5	47.4	48.5	56.3	58.5	6.8	5.8	0.7
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.6	5.1	7.8	16.6	25.5	24.0	47.4	67.3	82.1	107.9	7.2	5.8	0.6
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	3.0	4.6	5.1	18.9	29.2	28.0	53.0	67.1	61.2	117.0	7.5	6.2	0.7
1969-73	0.0	0.0	0.0	0.0	0.1	0.1	0.3	1.1	3.0	4.6	7.9	17.0	26.2	28.9	45.3	64.1	73.7	95.5	7.2	5.9	0.7
1974	0.0	0.0	0.0	0.0	0.3	0.0	0.4	1.7	2.1	3.0	5.7	12.5	25.1	39.3	38.1	52.6	87.8	87.1	6.9	5.6	0.6
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.1	7.7	8.1	14.3	16.3	31.1	43.5	58.7	65.9	126.9	6.9	5.6	0.6
1976	0.3	0.0	0.0	0.3	0.0	0.3	0.3	0.0	2.1	4.7	8.0	10.0	21.9	41.3	52.3	50.8	84.4	65.2	6.9	5.7	0.7
1977	0.3	0.0	0.0	0.0	0.5	0.0	1.0	0.8	2.6	6.9	12.3	13.8	21.6	37.1	41.8	49.6	72.2	87.8	7.2	6.0	0.7
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	3.9	8.1	10.7	19.7	25.3	59.0	54.9	75.2	89.1	6.5	5.3	0.6
1974-78	0.1	0.0	0.0	0.1	0.2	0.1	0.4	0.6	2.1	5.2	8.5	12.2	20.9	34.7	47.2	53.3	77.0	90.9	6.9	5.6	0.7
1979	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	3.4	3.9	7.7	12.4	24.2	35.5	51.3	75.4	83.9	108.3	7.6	6.1	0.7
1980	0.0	0.0	0.0	0.2	0.0	0.0	0.6	0.7	3.3	3.0	10.3	11.3	20.7	30.4	43.6	67.5	63.1	97.8	6.8	5.5	0.6
1981	0.0	0.3	0.0	0.0	0.0	0.0	0.3	0.0	2.0	5.6	11.6	16.7	22.4	40.8	48.3	59.2	78.2	95.1	7.6	6.3	0.7
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.3	3.2	3.5	10.3	10.7	20.2	31.7	53.6	61.9	61.2	87.0	6.9	5.7	0.7
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.6	1.9	5.6	7.3	12.4	19.5	33.5	52.3	45.3	73.4	68.5	6.5	5.4	0.7
1979-83	0.0	0.1	0.0	0.1	0.0	0.1	0.4	0.6	2.8	4.3	9.4	12.7	21.3	34.4	49.9	61.4	71.8	90.5	7.1	5.8	0.7

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF RETROPERITONEUM AND PERITONEUM (ICD9 158)

YEAR OF DIAGNOSIS	AGE (YEARS)																			ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	0	0	2	1	0	1	2	0	3	0	0	0	0	9
1965	1	0	0	0	0	0	1	0	1	1	1	1	1	2	3	0	0	0	0	12
1966	0	0	0	0	0	0	0	0	2	1	0	1	1	2	1	1	0	3	0	12
1967	0	0	0	1	0	0	0	0	2	1	1	0	0	0	1	1	1	2	0	10
1968	0	0	0	1	0	0	0	0	1	2	1	1	0	4	2	1	0	0	0	13
1964-68	1	0	0	2	0	0	1	0	8	6	3	4	4	8	10	3	1	5	0	56
1969	0	3	0	0	0	0	0	0	1	0	0	0	0	1	2	0	1	0	0	8
1970	0	0	0	0	0	0	0	0	0	1	0	1	1	3	1	1	2	1	0	11
1971	0	0	0	0	0	0	0	2	0	1	0	1	0	2	2	3	0	1	0	12
1972	0	0	0	0	0	0	0	1	0	0	1	1	1	0	1	0	1	0	0	6
1973	0	0	0	0	0	1	0	1	1	1	0	0	1	2	0	0	0	0	0	7
1969-73	0	3	0	0	0	1	0	4	2	3	1	3	3	8	6	4	4	2	0	44
1974	0	0	0	0	0	0	1	0	0	3	4	4	4	1	1	2	0	0	0	20
1975	0	0	0	0	0	0	0	0	1	1	2	2	3	0	1	0	0	1	0	11
1976	0	0	0	0	0	0	0	0	0	1	1	0	0	2	3	0	0	0	0	7
1977	0	0	0	0	0	0	0	0	0	1	2	1	0	3	2	0	3	0	1	13
1978	0	0	0	0	0	0	0	0	0	1	1	2	1	1	2	0	1	2	0	11
1974-78	0	0	0	0	0	0	1	0	1	7	10	9	8	7	9	2	4	3	1	62
1979	0	0	0	0	0	0	0	0	0	1	1	0	5	0	1	1	0	0	0	9
1980	0	0	0	0	0	0	2	0	0	0	3	1	4	3	1	0	1	0	0	15
1981	1	0	0	0	0	0	1	1	1	2	0	3	2	4	2	1	0	0	0	18
1982	1	0	0	0	0	0	0	1	2	1	2	0	2	3	1	0	0	1	0	14
1983	1	0	0	0	0	0	0	0	0	0	2	1	1	4	0	1	1	0	0	11
1979-83	3	0	0	0	0	0	3	2	3	4	8	5	14	14	5	3	2	1	0	67

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF RETROPERITONEUM AND PERITONEUM (ICD9 158)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.5	0.0	0.7	1.8	0.0	4.4	0.0	0.0	0.0	0.3	0.3	0.0
1965	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.5	0.6	0.7	0.9	2.2	4.3	0.0	0.0	0.0	0.4	0.4	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.5	0.0	0.7	0.8	2.1	1.4	2.1	0.0	21.4	0.4	0.4	0.0
1967	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.8	0.5	0.5	0.0	0.0	0.0	1.4	2.1	3.8	13.8	0.3	0.3	0.0
1968	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	1.0	0.5	0.6	0.0	4.1	2.8	2.1	0.0	0.0	0.4	0.4	0.0
1964-68	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.7	0.6	0.3	0.5	0.7	1.7	2.9	1.3	0.8	7.0	0.3	0.3	0.0
1969	0.0	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	1.0	2.8	0.0	3.7	0.0	0.2	0.2	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.6	0.8	2.9	1.4	2.1	7.2	6.1	0.3	0.3	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.4	0.0	0.6	0.0	1.9	2.7	6.1	0.0	5.8	0.3	0.3	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	0.6	0.7	0.0	1.3	0.0	3.5	0.0	0.2	0.1	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.4	0.4	0.0	0.0	0.7	1.8	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1969-73	0.0	0.2	0.0	0.0	0.0	0.1	0.0	0.3	0.2	0.3	0.1	0.4	0.4	1.5	1.6	1.6	2.8	2.3	0.2	0.2	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.3	1.8	2.4	2.7	0.9	1.2	4.0	0.0	0.0	0.5	0.5	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.9	1.2	1.9	0.0	1.2	0.0	0.0	5.3	0.3	0.3	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.0	1.7	3.5	0.0	0.0	0.0	0.2	0.2	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.5	0.0	2.4	2.3	0.0	9.9	0.0	0.3	0.3	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.0	0.6	0.8	2.2	0.0	3.2	10.0	0.2	0.2	0.0
1974-78	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.6	0.9	1.0	1.0	1.2	2.1	0.7	2.7	3.1	0.3	0.3	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	3.2	0.0	1.1	1.6	0.0	0.0	0.2	0.2	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	1.3	0.5	2.5	2.2	1.0	0.0	3.0	0.0	0.3	0.3	0.0
1981	0.3	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.8	0.0	1.4	1.2	2.8	2.0	1.5	0.0	0.0	0.4	0.4	0.0
1982	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	0.4	0.8	0.0	1.1	2.1	1.0	0.0	0.0	4.6	0.3	0.3	0.0
1983	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.5	0.5	2.8	0.0	1.5	2.6	0.0	0.2	0.2	0.0
1979-83	0.2	0.0	0.0	0.0	0.0	0.0	0.2	0.1	0.2	0.3	0.7	0.5	1.6	2.0	1.0	0.9	1.2	0.9	0.3	0.3	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF RETROPERITONEUM AND PERITONEUM (ICD9 158)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	1	0	0	0	0	0	1	2	0	1	3	2	2	3	1	3	1	0	20
1965	0	0	1	0	0	0	0	0	3	2	1	0	2	4	1	1	2	1	0	18
1966	1	0	0	0	0	1	0	0	1	1	0	2	1	2	0	1	2	1	0	13
1967	0	0	0	0	0	0	0	0	0	0	1	0	0	0	2	1	1	0	0	5
1968	0	0	0	0	0	0	0	0	0	2	1	1	2	1	0	3	0	1	0	11
1964-68	1	1	1	0	0	1	0	1	6	5	4	6	7	9	6	7	8	4	0	67
1969	0	0	0	0	0	0	0	0	0	0	0	2	1	1	1	2	1	1	0	9
1970	0	0	0	0	0	0	2	0	0	0	0	0	3	3	1	1	1	1	0	12
1971	0	0	0	0	0	0	0	1	2	0	1	3	0	2	2	2	0	1	0	14
1972	0	0	0	0	0	0	1	0	0	0	0	0	0	0	2	2	1	1	1	8
1973	0	0	0	0	0	0	0	2	0	0	1	2	1	2	0	3	1	1	0	13
1969-73	0	0	0	0	0	0	3	3	2	0	2	7	5	8	6	10	4	5	1	56
1974	0	0	0	0	0	0	1	1	0	1	0	1	0	2	2	2	0	0	0	10
1975	0	0	0	0	0	0	1	0	1	1	0	1	0	2	0	2	2	0	1	11
1976	0	0	0	0	0	0	1	0	1	1	0	1	0	1	2	1	1	0	1	10
1977	0	0	0	0	0	0	0	1	1	2	2	1	1	0	0	3	3	0	0	14
1978	0	0	0	0	1	0	0	0	0	0	0	2	0	1	2	1	1	2	0	10
1974-78	0	0	0	0	1	0	3	2	3	5	2	6	1	6	6	9	7	2	2	55
1979	0	0	0	0	0	0	1	1	0	2	2	3	0	0	1	2	2	1	0	15
1980	0	0	0	0	0	0	0	1	1	1	0	0	2	1	4	0	1	0	0	11
1981	1	1	0	0	0	0	0	0	1	0	4	0	1	0	3	1	0	2	0	14
1982	0	1	0	0	0	0	0	0	0	0	0	1	1	1	1	2	2	0	0	9
1983	1	0	0	1	1	1	2	0	0	0	0	2	2	0	1	1	1	0	0	13
1979-83	2	2	0	1	1	1	3	2	2	3	6	6	6	2	10	6	6	3	0	62

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF RETROPERITONEUM AND PERITONEUM (ICD9 158)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.0	0.6	2.2	1.7	2.0	3.6	1.7	8.7	4.5	0.6	0.5	0.1
1965	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	1.3	1.1	0.6	0.0	1.7	3.9	1.2	1.7	5.6	4.3	0.5	0.5	0.0
1966	0.3	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.4	0.5	0.0	1.4	0.8	1.9	0.0	1.6	5.4	4.2	0.4	0.3	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	2.2	1.5	2.6	0.0	0.1	0.1	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.6	0.6	1.5	0.9	0.0	4.5	0.0	3.8	0.3	0.3	0.0
1964-68	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.5	0.5	0.5	0.8	1.1	1.7	1.4	2.2	4.3	3.3	0.4	0.3	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.7	0.9	1.1	2.9	2.4	3.5	0.2	0.2	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.0	2.1	2.5	1.1	1.4	2.3	3.3	0.3	0.3	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.0	0.5	1.7	0.0	1.6	2.1	2.8	0.0	3.1	0.3	0.3	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.0	2.7	2.1	2.9	0.2	0.1	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.5	1.1	0.6	1.6	0.0	3.9	2.0	2.8	0.3	0.3	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.2	0.0	0.2	0.8	0.7	1.3	1.2	2.8	1.7	3.1	0.3	0.2	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.4	0.0	0.6	0.0	1.5	1.9	2.6	0.0	0.0	0.2	0.2	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.4	0.0	0.6	0.0	1.5	0.0	2.5	3.9	0.0	0.2	0.2	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.4	0.0	0.5	0.0	0.7	1.8	1.2	1.9	0.0	0.2	0.2	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.9	0.8	0.5	0.6	0.0	0.0	3.5	5.4	0.0	0.3	0.2	0.0
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.7	1.7	1.1	1.7	4.5	0.2	0.2	0.0
1974-78	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.2	0.3	0.4	0.2	0.6	0.1	0.9	1.1	2.2	2.6	1.0	0.2	0.2	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.0	0.9	0.9	1.3	0.0	0.0	0.8	2.2	3.4	2.1	0.3	0.3	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.0	0.0	1.1	0.6	3.2	0.0	1.6	0.0	0.2	0.2	0.0
1981	0.3	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.7	0.0	0.5	0.0	2.3	1.0	0.0	3.9	0.3	0.3	0.0
1982	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.6	0.7	2.0	3.1	0.0	0.2	0.1	0.0
1983	0.3	0.0	0.0	0.3	0.2	0.3	0.5	0.0	0.0	0.0	0.0	0.9	1.0	0.0	0.7	1.0	1.5	0.0	0.3	0.2	0.0
1979-83	0.1	0.1	0.0	0.1	0.1	0.1	0.2	0.1	0.2	0.3	0.5	0.5	0.6	0.2	1.5	1.2	1.9	1.2	0.2	0.2	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE DIGESTIVE ORGANS AND PERITONEUM (ICD9 159)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	1	2	4	3	3	6	8	8	10	4	4	0	53
1965	0	0	0	0	0	0	1	0	0	5	4	5	5	3	6	10	4	7	0	50
1966	0	0	0	0	0	0	1	2	3	4	3	2	6	12	4	10	8	1	0	56
1967	0	0	0	0	0	0	0	0	0	3	3	4	6	6	11	9	5	5	0	52
1968	0	0	0	0	0	0	1	0	1	1	1	3	7	5	3	8	6	3	0	39
1964-68	0	0	0	0	0	0	3	3	6	17	14	17	30	34	32	47	27	20	0	250
1969	0	0	0	0	0	0	0	0	3	3	3	1	2	5	6	10	6	2	0	41
1970	0	0	0	0	0	0	1	0	1	3	1	2	5	9	8	3	2	4	0	39
1971	0	0	0	0	0	1	0	0	2	4	2	2	11	8	6	4	0	5	0	45
1972	0	0	0	0	0	1	0	1	2	1	5	1	9	9	12	5	6	10	0	62
1973	0	0	0	0	0	0	1	0	3	4	3	4	10	9	14	10	13	10	0	81
1969-73	0	0	0	0	0	2	2	1	11	15	14	10	37	40	46	32	27	31	0	268
1974	0	1	0	0	1	0	0	1	1	3	7	5	12	9	12	14	7	6	0	79
1975	0	0	0	0	0	1	0	0	2	0	2	2	5	16	8	11	13	8	0	68
1976	0	0	0	0	0	0	0	1	0	2	4	5	13	16	15	15	9	4	0	84
1977	0	0	0	0	0	1	1	2	3	2	5	7	12	10	17	9	8	5	0	82
1978	0	0	0	0	0	0	1	0	0	4	8	11	10	12	18	26	14	6	0	110
1974-78	0	1	0	0	1	2	2	4	6	11	26	30	52	63	70	75	51	29	0	423
1979	0	0	0	0	0	0	0	1	3	6	4	4	10	20	15	13	8	9	0	93
1980	0	0	0	0	0	1	0	0	4	1	7	6	12	22	10	17	15	14	0	109
1981	0	0	0	0	0	0	0	2	3	7	5	8	15	22	24	14	11	11	0	122
1982	0	0	0	0	0	0	1	0	0	2	6	10	14	15	13	19	8	7	1	96
1983	0	0	0	0	0	0	1	3	4	3	4	20	14	18	20	20	11	5	0	123
1979-83	0	0	0	0	0	1	2	6	14	19	26	48	65	97	82	83	53	46	1	543

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE DIGESTIVE ORGANS AND PERITONEUM (ICD9 159)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	2.1	1.8	2.1	5.3	8.9	11.7	22.0	15.7	29.4	1.6	1.5	0.2
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	2.6	2.3	3.5	4.3	3.3	8.6	21.8	15.4	50.4	1.5	1.4	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.3	2.0	1.7	1.4	5.0	12.9	5.7	21.5	30.8	7.1	1.6	1.5	0.2
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.5	1.6	2.6	4.9	6.3	15.5	18.9	18.9	34.5	1.5	1.4	0.2
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.5	0.5	1.9	5.6	5.1	4.2	16.7	22.4	20.0	1.1	1.0	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.5	1.7	1.6	2.3	5.0	7.3	9.1	20.2	20.7	28.2	1.5	1.4	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	1.4	1.6	0.6	1.6	5.0	8.4	20.8	22.1	12.7	1.1	1.0	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	1.3	0.5	1.2	3.8	8.7	11.0	6.2	7.2	24.4	1.0	1.1	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	1.7	1.1	1.2	8.1	7.5	8.1	8.2	0.0	28.9	1.2	1.2	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.8	0.4	2.5	0.6	6.4	8.3	15.6	10.1	20.9	55.9	1.6	1.6	0.2
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.2	1.7	1.4	2.4	6.9	8.2	17.6	20.1	44.7	54.9	2.1	1.9	0.2
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.9	1.3	1.5	1.2	5.4	7.6	12.3	13.1	19.1	36.3	1.4	1.4	0.2
1974	0.0	0.3	0.0	0.0	0.3	0.0	0.0	0.4	0.4	1.3	3.2	3.0	8.0	8.0	14.6	27.7	24.0	32.4	2.0	1.9	0.2
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	0.0	0.9	1.2	3.2	13.8	9.5	21.0	44.5	42.1	1.7	1.5	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.8	1.8	2.8	8.3	13.3	17.4	27.6	30.3	20.6	2.0	1.9	0.2
1977	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.8	1.3	0.8	2.2	3.7	7.6	7.9	19.3	15.9	26.5	25.4	1.9	1.8	0.2
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	1.7	3.5	5.5	6.3	9.2	19.8	44.1	45.5	30.0	2.5	2.3	0.2
1974-78	0.0	0.1	0.0	0.0	0.1	0.1	0.1	0.3	0.5	0.9	2.3	3.3	6.7	10.4	16.2	27.5	34.2	30.0	2.0	1.9	0.2
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.2	2.5	1.7	1.9	6.4	14.9	16.0	21.2	25.2	43.9	2.0	2.0	0.2
1980	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	1.6	0.4	3.0	2.8	7.4	15.9	10.3	27.0	45.3	67.6	2.3	2.2	0.2
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	1.2	3.0	2.1	3.7	8.9	15.6	23.8	21.6	32.3	51.9	2.5	2.5	0.3
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	2.5	4.6	7.9	10.5	12.4	28.6	22.4	32.4	1.9	1.8	0.2
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.9	1.5	1.3	1.7	9.0	7.5	12.6	18.3	29.2	29.1	22.4	2.4	2.3	0.3
1979-83	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.4	1.1	1.6	2.2	4.5	7.6	13.9	16.2	25.6	30.7	43.3	2.2	2.2	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE DIGESTIVE ORGANS AND PERITONEUM (ICD9 159)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	2	3	2	2	7	4	11	11	9	9	0	60
1965	0	0	0	0	0	0	0	0	1	3	3	2	3	3	4	8	3	15	0	45
1966	0	0	0	0	0	0	0	0	0	2	3	3	5	5	5	9	6	7	0	45
1967	0	0	0	0	0	0	0	2	0	0	3	1	3	11	8	4	4	8	0	44
1968	0	0	0	0	0	1	0	3	0	0	3	2	4	2	7	10	13	6	0	51
1964-68	0	0	0	0	0	1	0	5	3	8	14	10	22	25	35	42	35	45	0	245
1969	0	0	0	0	0	0	0	0	0	1	2	4	5	8	5	5	6	3	0	39
1970	0	0	0	0	0	0	1	3	0	0	3	6	1	5	11	10	8	4	0	52
1971	0	0	0	0	0	0	0	0	1	0	2	3	4	8	5	8	8	9	0	48
1972	0	0	0	0	0	0	0	0	0	1	1	2	6	12	11	10	15	14	0	72
1973	0	0	0	0	0	0	0	0	0	0	3	6	1	7	7	9	12	20	1	66
1969-73	0	0	0	0	0	0	1	3	1	2	11	21	17	40	39	42	49	50	1	277
1974	0	0	0	0	0	0	0	0	0	4	4	5	7	12	12	20	12	16	0	92
1975	0	0	0	0	0	0	0	0	0	0	4	5	5	9	11	24	9	14	0	81
1976	0	0	0	0	0	1	1	0	2	2	3	5	3	15	17	18	21	23	0	109
1977	0	0	0	0	0	0	1	0	1	6	4	7	4	7	13	16	6	17	0	82
1978	0	0	0	0	0	2	0	1	0	5	5	8	8	7	17	17	12	25	0	107
1974-78	0	0	0	0	0	3	2	1	3	17	20	30	27	48	70	95	60	95	0	471
1979	0	0	0	0	0	0	1	1	2	1	2	4	6	12	11	8	10	20	0	78
1980	0	0	1	0	0	0	0	0	0	1	4	6	4	12	16	9	11	16	0	80
1981	0	0	0	0	0	0	0	0	2	0	2	7	6	10	10	12	16	12	0	77
1982	0	0	0	0	0	0	0	0	1	3	3	9	3	10	7	16	12	23	0	87
1983	0	0	0	0	0	0	0	2	2	2	2	6	6	6	13	8	11	22	1	81
1979-83	0	0	1	0	0	0	1	3	7	7	13	32	25	50	57	53	60	93	1	403

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE DIGESTIVE ORGANS AND PERITONEUM (ICD9 159)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.6	1.2	1.4	5.9	3.9	13.3	18.8	26.0	40.9	1.8	1.4	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.6	1.8	1.4	2.5	2.9	4.7	13.2	8.4	65.2	1.4	1.0	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	1.7	2.1	4.0	4.7	5.6	14.5	16.1	29.5	1.3	1.0	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	1.7	0.7	2.3	10.1	8.9	6.2	10.3	31.9	1.2	1.0	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.3	0.0	0.0	1.7	1.3	3.0	1.8	7.7	14.9	31.9	22.6	1.4	1.0	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.4	0.3	0.8	1.6	1.4	3.5	4.7	8.0	13.4	18.7	37.4	1.4	1.1	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.1	2.5	3.7	7.0	5.4	7.3	14.2	10.6	1.0	0.8	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	0.0	0.0	1.6	3.6	0.7	4.2	11.6	14.2	18.1	13.2	1.3	1.0	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.0	1.7	2.8	6.6	5.1	11.1	17.3	27.7	1.2	0.9	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	1.1	4.0	9.6	11.1	13.5	31.6	40.8	1.7	1.3	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	3.4	0.6	5.4	6.9	11.8	24.5	55.7	1.5	1.1	0.1
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.2	1.1	2.5	2.4	6.6	8.0	11.6	21.4	31.0	1.3	1.0	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	1.8	2.8	4.4	9.1	11.4	25.7	24.0	42.2	2.0	1.6	0.2
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	2.8	3.0	6.7	10.2	30.0	17.4	35.5	1.7	1.3	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.9	0.9	1.3	2.6	1.8	9.3	15.3	21.8	39.4	55.6	2.3	1.7	0.2
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	2.6	1.7	3.5	2.3	4.8	11.3	18.9	10.8	39.3	1.7	1.3	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.4	0.0	2.1	2.1	3.7	4.6	4.7	14.3	19.5	21.0	55.7	2.1	1.6	0.2
1974-78	0.0	0.0	0.0	0.0	0.0	0.2	0.1	0.1	0.3	1.5	1.7	3.1	3.2	6.8	12.6	23.0	22.4	45.9	2.0	1.5	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.8	0.4	0.9	1.8	3.5	7.7	9.0	8.9	16.8	42.5	1.5	1.2	0.1
1980	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	2.6	2.2	7.4	12.7	9.6	17.8	32.6	1.5	1.2	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.9	3.0	3.2	6.1	7.7	12.5	25.0	23.3	1.4	1.0	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	1.3	3.9	1.5	6.0	5.2	16.0	18.3	42.6	1.5	1.1	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.8	0.9	0.9	2.6	2.9	3.6	9.3	7.7	16.2	38.7	1.3	1.0	0.1
1979-83	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.2	0.6	0.6	1.1	2.8	2.6	6.1	8.7	11.0	18.8	36.0	1.4	1.1	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF NASAL CAVITIES, MIDDLE EAR AND ACCESSORY SINUSES (ICD9 160)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	0	0	0	0	0	0	3	2	3	2	4	6	7	3	6	0	1	0	38
1965	0	0	0	0	0	0	0	1	2	1	1	5	2	2	6	3	1	2	0	26
1966	0	0	0	0	0	1	0	1	2	2	5	5	6	3	3	3	3	0	0	34
1967	0	0	0	0	0	0	0	3	0	1	2	3	2	4	2	4	0	2	0	23
1968	0	0	0	0	1	0	0	1	0	3	1	4	7	5	0	4	1	0	0	27
1964-68	1	0	0	0	1	1	0	9	6	10	11	21	23	21	14	20	5	5	0	148
1969	0	0	0	0	0	1	1	0	0	5	6	5	3	10	2	2	4	6	0	45
1970	1	0	0	0	0	1	0	0	2	2	4	4	4	5	5	2	4	3	0	37
1971	0	0	0	1	0	0	1	1	2	0	6	3	2	6	3	3	0	3	0	31
1972	0	0	0	0	0	0	0	1	2	1	3	4	5	4	4	1	1	2	0	28
1973	0	0	0	0	0	1	1	2	1	4	7	8	5	8	2	5	6	0	1	51
1969-73	1	0	0	1	0	3	3	4	7	12	26	24	19	33	16	13	15	14	1	192
1974	0	0	2	0	0	0	1	2	2	7	3	6	8	2	9	8	4	5	0	59
1975	0	0	0	0	2	0	1	1	3	5	4	4	7	5	9	4	3	1	0	49
1976	1	0	0	0	0	2	0	0	4	11	6	6	12	10	7	8	6	2	0	75
1977	0	0	0	2	0	0	2	1	2	5	6	5	6	6	11	5	3	2	0	56
1978	1	0	0	0	2	0	2	0	1	1	2	9	9	9	11	6	7	2	1	63
1974-78	2	0	2	2	4	2	6	4	12	29	21	30	42	32	47	31	23	12	1	302
1979	0	3	0	0	0	0	1	1	3	1	5	6	3	4	5	6	0	1	0	39
1980	1	1	0	0	0	0	0	1	0	3	2	8	6	2	8	4	4	1	0	41
1981	0	1	0	0	0	2	4	4	2	3	5	8	4	2	4	3	5	1	0	48
1982	0	0	0	0	0	0	0	2	1	6	5	3	6	3	6	1	2	1	0	36
1983	0	0	0	0	0	1	1	2	0	2	4	3	6	4	7	5	1	2	1	39
1979-83	1	5	0	0	0	3	6	10	6	15	21	28	25	15	30	19	12	6	1	203

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF NASAL CAVITIES, MIDDLE EAR AND ACCESSORY SINUSES (ICD9 160)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.9	1.6	1.2	2.9	5.3	7.8	4.4	13.2	0.0	7.4	1.1	1.1	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.5	0.6	3.5	1.7	2.2	8.6	6.5	3.9	14.4	0.8	0.7	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.4	0.9	1.0	2.8	3.4	5.0	3.2	4.3	6.4	11.5	0.0	1.0	1.0	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.0	0.5	1.1	2.0	1.6	4.2	2.8	8.4	0.0	13.8	0.7	0.6	0.1
1968	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.4	0.0	1.4	0.5	2.5	5.6	5.1	0.0	8.4	3.7	0.0	0.7	0.7	0.1
1964-68	0.1	0.0	0.0	0.0	0.1	0.1	0.0	0.8	0.5	1.0	1.2	2.8	3.8	4.5	4.0	8.6	3.8	7.0	0.9	0.8	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.0	2.3	3.3	3.1	2.3	10.0	2.8	4.2	14.7	38.2	1.2	1.2	0.1
1970	0.3	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.8	0.9	2.2	2.4	3.0	4.9	6.9	4.1	14.4	18.3	1.0	1.0	0.1
1971	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	0.8	0.0	3.2	1.8	1.5	5.7	4.0	6.1	0.0	17.3	0.8	0.8	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	0.4	1.5	2.3	3.5	3.7	5.2	2.0	3.5	11.2	0.7	0.7	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.8	0.4	1.7	3.4	4.7	3.4	7.3	2.5	10.0	20.6	0.0	1.3	1.2	0.1
1969-73	0.1	0.0	0.0	0.1	0.0	0.2	0.2	0.3	0.6	1.0	2.7	2.9	2.8	6.3	4.3	5.3	10.6	16.4	1.0	1.0	0.1
1974	0.0	0.0	0.5	0.0	0.0	0.0	0.4	0.8	0.8	3.0	1.4	3.6	5.3	1.8	10.9	15.8	13.7	27.0	1.5	1.4	0.1
1975	0.0	0.0	0.0	0.0	0.6	0.0	0.3	0.4	1.2	2.1	1.8	2.3	4.5	4.3	10.7	7.6	10.3	5.3	1.2	1.1	0.1
1976	0.3	0.0	0.0	0.0	0.0	0.6	0.0	0.0	1.7	4.6	2.6	3.4	7.6	8.3	8.1	14.7	20.2	10.3	1.8	1.7	0.2
1977	0.0	0.0	0.0	0.5	0.0	0.0	0.6	0.4	0.8	2.1	2.6	2.6	3.8	4.8	12.5	8.8	9.9	10.2	1.3	1.3	0.2
1978	0.3	0.0	0.0	0.0	0.5	0.0	0.6	0.0	0.4	0.4	0.9	4.5	5.7	6.9	12.1	10.2	22.7	10.0	1.4	1.3	0.2
1974-78	0.1	0.0	0.1	0.1	0.2	0.1	0.4	0.3	1.0	2.4	1.9	3.3	5.4	5.3	10.9	11.4	15.4	12.4	1.4	1.4	0.2
1979	0.0	0.9	0.0	0.0	0.0	0.0	0.3	0.4	1.2	0.4	2.2	2.9	1.9	3.0	5.3	9.8	0.0	4.9	0.9	0.8	0.1
1980	0.3	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.3	0.9	3.8	3.7	1.4	8.2	6.4	12.1	4.8	0.9	0.9	0.1
1981	0.0	0.3	0.0	0.0	0.0	0.6	1.1	1.4	0.8	1.3	2.1	3.7	2.4	1.4	4.0	4.6	14.7	4.7	1.0	1.0	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.4	2.5	2.1	1.4	3.4	2.1	5.7	1.5	5.6	4.6	0.7	0.8	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.6	0.0	0.8	1.7	1.4	3.2	2.8	6.4	7.3	2.6	9.0	0.8	0.7	0.1
1979-83	0.1	0.3	0.0	0.0	0.0	0.2	0.3	0.7	0.5	1.3	1.8	2.6	2.9	2.1	5.9	5.9	7.0	5.6	0.8	0.8	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF NASAL CAVITIES, MIDDLE EAR AND ACCESSORY SINUSES (ICD9 160)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	0	0	1	2	5	2	4	4	2	1	2	0	0	23
1965	0	0	0	0	0	0	0	1	0	0	0	1	0	1	2	0	0	0	0	5
1966	0	0	0	0	0	0	0	0	1	4	0	3	0	1	4	4	1	1	0	19
1967	0	0	0	0	0	0	1	2	0	1	2	1	5	2	6	6	0	1	0	27
1968	0	0	0	0	0	1	0	1	2	2	4	3	2	4	4	2	6	1	0	32
1964-68	0	0	0	0	0	1	1	4	4	9	11	10	11	12	18	13	9	3	0	106
1969	0	0	0	0	1	0	0	1	1	1	4	1	3	3	0	3	1	3	0	22
1970	0	0	0	1	0	0	0	0	0	2	5	1	2	1	2	0	4	3	0	21
1971	0	0	0	0	0	0	0	0	1	1	1	1	3	0	4	1	2	2	0	16
1972	0	0	0	0	0	0	0	0	0	4	1	3	5	3	3	4	2	0	1	26
1973	0	0	0	0	0	0	0	3	2	3	2	2	6	0	8	4	3	2	1	36
1969-73	0	0	0	1	1	0	0	4	4	11	13	8	19	7	17	12	12	10	2	121
1974	0	0	0	0	0	1	1	0	4	3	4	4	8	1	7	5	5	5	0	48
1975	0	0	0	0	0	0	0	1	1	7	4	6	3	5	1	5	6	3	0	42
1976	0	0	0	0	0	1	0	3	6	4	4	4	7	8	10	12	6	11	0	76
1977	1	0	0	0	0	1	3	2	6	5	5	7	8	5	7	13	3	10	0	76
1978	0	0	0	0	0	0	0	2	4	0	2	4	4	7	1	6	8	5	0	43
1974-78	1	0	0	0	0	3	4	8	21	19	19	25	30	26	26	41	28	34	0	285
1979	0	0	0	1	0	1	0	0	1	4	2	4	4	2	3	0	5	2	0	30
1980	1	0	0	0	0	0	0	2	0	0	0	2	2	4	4	4	1	0	0	20
1981	0	0	1	0	0	1	1	1	0	2	6	2	1	3	3	5	2	2	0	30
1982	0	0	0	0	0	0	0	2	0	3	3	0	8	6	3	5	4	3	1	38
1983	1	0	0	0	1	0	1	0	1	1	0	4	0	2	3	5	0	2	0	21
1979-83	2	0	1	1	1	2	3	5	2	10	11	12	15	17	16	19	12	9	1	139

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF NASAL CAVITIES, MIDDLE EAR AND ACCESSORY SINUSES (ICD9 160)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.1	3.0	1.4	3.4	3.9	2.4	1.7	5.8	0.0	0.7	0.6	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.7	0.0	1.0	2.3	0.0	0.0	0.0	0.1	0.1	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.0	0.0	2.1	0.0	0.9	4.5	6.5	2.7	4.2	0.5	0.4	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.0	0.5	1.1	0.7	3.9	1.8	6.7	9.3	0.0	4.0	0.7	0.6	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.8	0.9	2.2	1.9	1.5	3.6	4.4	3.0	14.7	3.8	0.9	0.7	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.3	0.3	0.9	1.3	1.4	1.8	2.3	4.1	4.2	4.8	2.5	0.6	0.5	0.1
1969	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	0.5	2.2	0.6	2.2	2.6	0.0	4.4	2.4	10.6	0.6	0.5	0.0
1970	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.9	2.7	0.6	1.4	0.8	2.1	0.0	9.0	9.9	0.5	0.5	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.6	2.1	0.0	4.1	1.4	4.3	6.2	0.4	0.3	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	0.5	1.7	3.4	2.4	3.0	5.4	4.2	0.0	0.6	0.5	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.9	1.3	0.9	1.1	3.9	0.0	7.9	5.3	6.1	5.6	0.8	0.7	0.1
1969-73	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.3	0.3	0.9	1.3	0.9	2.6	1.2	3.5	3.3	5.2	6.2	0.6	0.5	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.0	1.7	1.3	1.8	2.3	5.0	0.8	6.7	6.4	10.0	13.2	1.1	0.9	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	3.0	1.7	3.3	1.8	3.7	0.9	6.3	11.6	7.6	0.9	0.8	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.3	0.0	1.2	2.6	1.7	1.7	2.1	4.1	5.7	9.0	14.5	11.3	26.6	1.6	1.4	0.1
1977	0.3	0.0	0.0	0.0	0.0	0.3	1.0	0.8	2.6	2.1	2.1	3.5	4.7	3.4	6.1	15.3	5.4	23.1	1.6	1.4	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.7	0.0	0.8	1.9	2.3	4.7	0.8	6.9	14.0	11.1	0.9	0.7	0.1
1974-78	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.7	1.8	1.6	1.6	2.6	3.6	3.7	4.7	9.9	10.5	16.4	1.2	1.0	0.1
1979	0.0	0.0	0.0	0.2	0.0	0.3	0.3	0.0	0.4	1.7	0.9	1.8	2.3	1.3	2.4	0.0	8.4	4.2	0.6	0.5	0.1
1980	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0	0.0	0.9	1.1	2.5	3.2	4.3	1.6	0.0	0.4	0.4	0.0
1981	0.0	0.0	0.3	0.0	0.0	0.3	0.3	0.3	0.0	0.9	2.6	0.9	0.5	1.8	2.3	5.2	3.1	3.9	0.6	0.5	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	1.3	1.3	0.0	4.0	3.6	2.2	5.0	6.1	5.6	0.7	0.6	0.1
1983	0.3	0.0	0.0	0.0	0.2	0.0	0.3	0.0	0.4	0.4	0.0	1.7	0.0	1.2	2.1	4.8	0.0	3.5	0.4	0.3	0.0
1979-83	0.1	0.0	0.1	0.1	0.1	0.1	0.2	0.3	0.2	0.9	0.9	1.0	1.6	2.1	2.4	3.9	3.8	3.5	0.5	0.5	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF LARYNX (ICD9 161)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	1	1	9	13	12	33	28	19	18	6	7	3	0	150
1965	0	0	0	0	0	0	0	2	4	12	24	20	26	23	22	13	11	1	0	158
1966	0	0	0	0	0	0	0	2	10	10	25	25	18	24	27	12	7	3	0	163
1967	0	0	0	0	0	0	0	4	7	6	16	25	36	28	19	17	2	7	0	167
1968	0	1	0	0	0	0	0	2	9	13	19	24	39	29	24	14	6	2	0	182
1964-68	0	1	0	0	0	0	1	11	39	54	96	127	147	123	110	62	33	16	0	820
1969	0	0	0	0	0	0	1	1	4	6	19	24	31	33	25	14	8	2	0	168
1970	0	0	0	0	0	1	1	2	10	15	28	40	28	38	25	19	11	9	0	227
1971	0	0	0	0	0	1	1	1	9	19	21	36	42	21	22	18	10	2	0	203
1972	1	0	0	0	0	1	1	3	9	16	26	31	36	43	24	14	13	3	0	221
1973	0	1	0	0	0	2	0	0	11	12	27	33	44	31	27	15	13	4	0	220
1969-73	1	1	0	0	0	5	4	7	43	68	121	164	181	166	123	80	55	20	0	1039
1974	0	0	0	0	0	0	2	1	15	29	35	33	41	32	29	14	8	9	0	248
1975	0	0	0	0	0	1	1	2	14	15	31	33	47	38	20	16	11	4	0	233
1976	0	0	0	0	1	3	3	4	10	25	46	45	47	44	33	13	5	7	5	291
1977	0	1	0	0	0	1	2	3	7	26	39	35	52	41	28	15	8	4	5	267
1978	0	0	0	0	0	0	1	0	4	13	33	36	50	52	34	14	12	4	1	254
1974-78	0	1	0	0	1	5	9	10	50	108	184	182	237	207	144	72	44	28	11	1293
1979	0	0	0	0	0	1	0	4	5	26	44	51	51	47	35	30	9	5	2	310
1980	0	0	0	0	0	0	1	3	6	14	30	55	63	63	33	23	17	5	0	313
1981	0	0	0	0	0	1	3	3	8	14	43	41	51	58	48	15	7	5	3	300
1982	0	0	0	1	0	0	0	1	3	20	28	52	60	57	53	26	6	5	0	312
1983	0	0	0	0	0	0	0	2	11	24	37	53	61	54	57	21	16	8	0	344
1979-83	0	0	0	1	0	2	4	13	33	98	182	252	286	279	226	115	55	28	5	1579

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF LARYNX (ICD9 161)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	4.1	6.9	7.1	23.6	24.6	21.1	26.4	13.2	27.6	22.1	4.5	4.5	0.6
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.8	6.3	13.7	14.0	22.3	25.1	31.7	28.3	42.5	7.2	4.6	4.6	0.6
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	4.3	5.1	14.0	17.0	15.1	25.8	38.4	25.8	26.9	21.4	4.7	4.6	0.6
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	2.9	3.0	8.8	16.4	29.3	29.5	26.8	35.8	7.6	48.3	4.7	4.8	0.6
1968	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.8	3.7	6.2	10.4	15.3	31.0	29.8	33.8	29.3	22.4	13.3	5.0	5.1	0.7
1964-68	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.9	3.4	5.5	10.8	17.2	24.5	26.3	31.4	26.6	25.3	22.5	4.7	4.7	0.6
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.7	2.8	10.4	14.8	24.0	33.0	34.8	29.1	29.4	12.7	4.6	4.6	0.6
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.8	4.1	6.6	15.1	23.9	21.1	36.9	34.3	39.2	39.6	54.9	6.0	6.0	0.7
1971	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.4	3.7	8.2	11.1	21.1	30.9	19.8	29.6	36.8	35.5	11.6	5.3	5.2	0.6
1972	0.3	0.0	0.0	0.0	0.0	0.3	0.4	1.3	3.7	6.9	13.1	18.1	25.5	39.8	31.3	28.3	45.3	16.8	5.6	5.6	0.7
1973	0.0	0.3	0.0	0.0	0.0	0.6	0.0	0.0	4.5	5.1	12.9	19.5	30.2	28.2	33.9	30.1	44.7	22.0	5.5	5.4	0.7
1969-73	0.1	0.1	0.0	0.0	0.0	0.3	0.3	0.6	3.5	5.9	12.5	19.5	26.5	31.5	32.8	32.7	39.0	23.4	5.4	5.4	0.7
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.4	6.1	12.3	16.0	19.6	27.3	28.3	35.2	27.7	27.4	48.6	6.0	6.1	0.7
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.8	5.8	6.3	13.9	19.2	30.3	32.8	23.7	30.5	37.7	21.1	5.6	5.5	0.7
1976	0.0	0.0	0.0	0.0	0.3	0.8	1.0	1.6	4.2	10.5	20.3	25.1	29.8	36.5	38.2	23.9	16.8	36.1	6.6	6.7	0.8
1977	0.0	0.3	0.0	0.0	0.0	0.3	0.6	1.2	2.9	10.8	17.1	18.5	33.0	32.6	31.7	26.5	26.5	20.3	5.9	6.0	0.7
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	1.7	5.4	14.4	18.1	31.7	40.0	37.5	23.8	39.0	20.0	5.6	5.6	0.7
1974-78	0.0	0.1	0.0	0.0	0.1	0.3	0.6	0.8	4.1	9.0	16.3	20.1	30.5	34.2	33.3	26.4	29.5	29.0	6.0	6.0	0.7
1979	0.0	0.0	0.0	0.0	0.0	0.3	0.0	1.5	2.1	10.8	19.1	24.5	32.5	34.9	37.4	49.0	28.3	24.4	6.7	6.7	0.8
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.3	1.1	2.5	5.9	12.9	25.8	39.1	45.4	34.0	36.6	51.4	24.2	6.6	6.6	0.8
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.8	1.0	3.2	5.9	18.3	19.0	30.2	41.1	47.7	23.2	20.5	23.6	6.1	6.2	0.8
1982	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.3	1.2	8.5	11.8	23.8	33.7	39.9	50.7	39.1	16.8	23.1	6.3	6.3	0.8
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	4.2	10.2	15.6	24.0	32.5	37.9	52.2	30.7	42.3	35.9	6.8	6.8	0.9
1979-83	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.9	2.6	8.3	15.5	23.4	33.6	39.9	44.7	35.5	31.9	26.3	6.5	6.5	0.8

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF LARYNX (ICD9 161)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	2	3	0	2	5	1	1	2	0	1	0	0	17
1965	0	0	0	0	0	0	0	2	1	1	1	2	3	4	2	0	1	1	0	18
1966	0	0	0	0	0	0	2	0	4	1	5	0	1	1	3	3	1	0	0	21
1967	0	0	0	0	0	1	0	0	3	1	0	1	4	0	3	2	2	0	0	17
1968	0	0	0	0	0	0	1	0	2	1	5	1	1	2	2	1	2	0	0	18
1964-68	0	0	0	0	0	1	3	4	13	4	13	9	10	8	12	6	7	1	0	91
1969	0	0	0	0	0	0	1	0	1	1	5	3	6	4	3	1	1	3	0	29
1970	0	0	0	0	0	1	0	2	2	4	1	2	3	3	2	1	0	0	0	21
1971	0	0	0	0	0	0	0	0	0	3	3	6	4	2	2	0	0	1	0	21
1972	0	0	0	1	1	0	2	0	4	4	6	8	8	3	2	2	1	0	0	42
1973	0	0	0	1	0	0	0	1	3	3	4	7	6	4	2	1	3	0	0	35
1969-73	0	0	0	2	1	1	3	3	10	15	19	26	27	16	11	5	5	4	0	148
1974	0	0	0	0	0	2	0	2	1	3	9	6	5	6	6	4	2	1	0	47
1975	0	0	0	0	1	0	0	1	4	4	5	3	5	3	5	3	1	0	0	35
1976	0	0	0	0	1	2	1	0	2	1	12	9	8	5	3	3	1	2	0	50
1977	0	0	0	1	0	1	1	0	2	6	9	7	8	3	3	1	3	1	3	49
1978	0	0	0	0	0	0	1	1	2	6	6	12	12	4	0	0	2	3	0	49
1974-78	0	0	0	1	2	5	3	4	11	20	41	37	38	21	17	11	9	7	3	230
1979	0	0	0	0	0	0	1	1	6	2	8	8	3	7	5	2	1	0	0	44
1980	0	0	0	0	0	1	3	1	3	6	6	7	8	9	3	2	0	0	0	49
1981	0	0	0	0	1	1	0	0	0	3	7	11	12	7	4	1	1	1	0	49
1982	0	0	1	0	0	0	0	0	2	4	3	9	8	9	8	3	5	2	0	54
1983	0	0	0	0	0	0	1	2	1	3	6	5	12	7	3	3	3	1	1	48
1979-83	0	0	1	0	1	2	5	4	12	18	30	40	43	39	23	11	10	4	1	244

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF LARYNX (ICD9 161)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.4	0.0	1.2	3.6	0.8	1.0	2.4	0.0	2.9	0.0	0.5	0.5	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	0.5	0.6	1.4	2.5	3.9	2.3	0.0	2.8	4.3	0.5	0.5	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	1.7	0.5	2.9	0.0	0.8	0.9	3.4	4.8	2.7	0.0	0.6	0.5	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.2	0.5	0.0	0.7	3.1	0.0	3.3	3.1	5.1	0.0	0.5	0.4	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.8	0.5	2.8	0.6	0.8	1.8	2.2	1.5	4.9	0.0	0.5	0.4	0.0
1964-68	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.3	1.1	0.4	1.5	1.2	1.6	1.5	2.7	1.9	3.7	0.8	0.5	0.5	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.5	2.7	1.8	4.4	3.5	3.2	1.5	2.4	10.6	0.8	0.7	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	0.9	1.7	0.5	1.2	2.1	2.5	2.1	1.4	0.0	0.0	0.5	0.5	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	1.6	3.5	2.8	1.6	2.1	0.0	0.0	3.1	0.5	0.5	0.1
1972	0.0	0.0	0.0	0.3	0.3	0.0	0.8	0.0	1.7	1.7	2.9	4.6	5.4	2.4	2.0	2.7	2.1	0.0	1.0	1.0	0.1
1973	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	1.3	1.3	1.9	4.0	3.9	3.1	2.0	1.3	6.1	0.0	0.8	0.8	0.1
1969-73	0.0	0.0	0.0	0.1	0.1	0.1	0.3	0.3	0.9	1.3	1.9	3.0	3.7	2.6	2.3	1.4	2.2	2.5	0.7	0.7	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.9	0.4	1.3	4.0	3.4	3.1	4.5	5.7	5.1	4.0	2.6	1.1	1.0	0.1
1975	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	1.7	1.7	2.1	1.7	3.0	2.2	4.6	3.7	1.9	0.0	0.8	0.8	0.1
1976	0.0	0.0	0.0	0.0	0.3	0.6	0.3	0.0	0.9	0.4	5.0	4.7	4.7	3.6	2.7	3.6	1.9	4.8	1.1	1.0	0.1
1977	0.0	0.0	0.0	0.3	0.0	0.3	0.3	0.0	0.9	2.6	3.8	3.5	4.7	2.1	2.6	1.2	5.4	2.3	1.0	1.0	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.9	2.6	2.5	5.6	7.0	2.7	0.0	0.0	3.5	6.7	1.0	1.0	0.1
1974-78	0.0	0.0	0.0	0.1	0.1	0.3	0.2	0.3	0.9	1.7	3.5	3.8	4.5	3.0	3.0	2.7	3.4	3.4	1.0	0.9	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	2.5	0.9	3.4	3.6	1.7	4.5	4.1	2.2	1.7	0.0	0.9	0.9	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.3	0.8	0.4	1.3	2.6	2.6	3.0	4.5	5.6	2.4	2.1	0.0	0.0	1.0	1.0	0.1
1981	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.0	1.3	3.0	4.7	6.4	4.3	3.1	1.0	1.6	1.9	1.0	0.9	0.1
1982	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.8	1.7	1.3	3.9	4.0	5.4	6.0	3.0	7.6	3.7	1.0	0.9	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.6	0.4	1.3	2.6	2.1	5.7	4.2	2.1	2.9	4.4	1.8	0.9	0.8	0.1
1979-83	0.0	0.0	0.1	0.0	0.1	0.1	0.3	0.3	1.0	1.6	2.6	3.5	4.5	4.8	3.5	2.3	3.1	1.5	1.0	0.9	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF TRACHEA, BRONCHUS AND LUNG (ICD9 162)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	2	1	2	15	51	67	131	241	299	294	231	135	58	24	0	1551
1965	0	0	1	0	0	1	3	13	28	65	139	205	237	262	208	130	53	24	0	1369
1966	0	0	0	1	0	0	5	17	53	72	151	241	250	314	237	138	71	30	0	1580
1967	0	0	0	1	1	0	4	15	49	84	142	224	305	311	277	144	73	25	0	1655
1968	0	0	1	1	1	4	6	20	44	113	152	236	281	312	247	156	76	29	0	1679
1964-68	0	0	2	3	4	6	20	80	225	401	715	1147	1372	1493	1200	703	331	132	0	7834
1969	0	0	0	0	0	2	3	18	43	104	175	259	340	342	273	150	79	44	1	1833
1970	0	0	1	1	1	4	6	21	45	104	150	273	340	309	303	168	99	31	1	1857
1971	1	1	0	0	1	3	7	21	54	99	166	259	351	411	339	213	114	51	0	2091
1972	0	0	1	0	1	5	9	22	62	127	209	307	392	406	314	203	127	58	0	2243
1973	0	0	1	3	0	3	3	22	53	110	222	263	408	411	397	228	118	58	3	2303
1969-73	1	1	3	4	3	17	28	104	257	544	922	1361	1831	1879	1626	962	537	242	5	10327
1974	2	0	2	0	1	2	7	13	50	128	216	329	416	462	398	246	132	57	8	2469
1975	0	0	0	0	1	5	5	13	53	107	214	314	482	451	442	292	143	69	3	2594
1976	1	0	0	1	4	3	5	12	46	143	249	349	455	500	461	308	171	62	5	2775
1977	0	0	0	1	3	6	11	13	48	122	254	345	427	511	426	302	132	61	1	2663
1978	0	0	0	2	2	4	8	15	42	118	287	373	464	546	460	346	172	77	1	2917
1974-78	3	0	2	4	11	20	36	66	239	618	1220	1710	2244	2470	2187	1494	750	326	18	13418
1979	0	0	0	1	1	0	9	18	41	116	255	394	463	525	455	367	151	77	5	2878
1980	0	0	0	0	0	2	8	8	51	119	267	435	441	574	510	373	181	91	7	3067
1981	0	0	0	1	1	2	8	16	40	102	263	413	525	570	520	374	184	90	0	3109
1982	0	0	0	0	0	0	6	24	42	111	234	389	549	629	552	372	196	113	7	3224
1983	0	0	0	1	2	0	11	13	48	105	235	443	575	609	564	405	222	104	5	3342
1979-83	0	0	0	3	4	4	42	79	222	553	1254	2074	2553	2907	2601	1891	934	475	24	15620

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF TRACHEA, BRONCHUS AND LUNG (ICD9 162)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.9	0.5	0.9	6.4	23.4	35.5	77.1	172.5	262.5	326.7	338.2	297.4	228.3	176.5	46.8	46.9	6.0
1965	0.0	0.0	0.3	0.0	0.0	0.5	1.3	5.5	12.4	34.0	79.5	143.2	203.1	286.3	299.7	283.2	204.6	172.7	40.5	40.4	5.2
1966	0.0	0.0	0.0	0.3	0.0	0.0	2.2	7.1	22.7	36.8	84.4	163.8	209.0	337.3	336.6	295.1	273.1	214.3	45.8	45.5	5.8
1967	0.0	0.0	0.0	0.3	0.4	0.0	1.8	6.2	20.6	41.7	78.1	147.4	248.2	327.4	390.7	303.2	276.5	172.4	46.9	46.9	6.1
1968	0.0	0.0	0.3	0.3	0.4	1.7	2.6	8.4	18.3	54.1	83.2	150.1	223.2	321.0	347.4	326.4	283.6	193.3	46.7	46.5	5.9
1964-68	0.0	0.0	0.1	0.2	0.3	0.5	1.8	6.7	19.5	40.7	80.5	155.2	229.0	319.8	342.8	301.5	253.6	185.9	45.3	45.3	5.8
1969	0.0	0.0	0.0	0.0	0.0	0.8	1.3	7.6	17.8	47.8	95.6	159.5	263.8	342.0	380.2	311.9	290.4	280.3	49.9	50.1	6.4
1970	0.0	0.0	0.3	0.3	0.3	1.5	2.5	8.8	18.6	46.0	81.1	163.4	256.8	300.0	415.6	346.4	356.1	189.0	49.4	49.1	6.3
1971	0.3	0.2	0.0	0.0	0.3	1.0	2.9	8.8	22.3	42.6	87.9	152.0	257.9	387.4	455.6	435.6	404.3	294.8	54.5	54.1	6.9
1972	0.0	0.0	0.2	0.0	0.3	1.6	3.6	9.3	25.5	54.4	105.2	179.7	278.0	375.9	409.4	410.9	442.5	324.0	57.2	56.7	7.0
1973	0.0	0.0	0.2	0.8	0.0	0.9	1.2	9.3	21.8	46.8	106.3	155.1	280.4	373.3	498.7	457.8	405.5	318.7	57.7	57.0	7.2
1969-73	0.1	0.1	0.1	0.2	0.2	1.2	2.3	8.7	21.2	47.5	95.6	162.0	267.7	356.4	433.1	393.1	380.9	283.0	53.8	53.5	6.7
1974	0.6	0.0	0.5	0.0	0.3	0.6	2.5	5.4	20.5	54.1	98.7	195.1	276.8	409.2	483.6	486.2	452.1	308.1	60.6	59.6	7.4
1975	0.0	0.0	0.0	0.0	0.3	1.4	1.7	5.3	21.8	45.0	95.7	182.6	310.6	388.8	523.7	557.3	489.7	363.2	62.4	61.1	7.6
1976	0.3	0.0	0.0	0.2	1.1	0.8	1.7	4.9	19.1	59.8	109.6	195.0	288.9	414.9	533.6	566.2	575.8	319.6	65.0	63.4	7.8
1977	0.0	0.0	0.0	0.2	0.8	1.7	3.5	5.1	20.0	50.7	111.6	182.7	270.6	405.9	482.4	534.5	437.1	309.6	60.7	59.6	7.4
1978	0.0	0.0	0.0	0.5	0.5	1.1	2.4	5.7	17.6	48.8	125.3	187.4	294.6	419.7	507.2	587.4	558.4	385.0	64.8	63.5	7.7
1974-78	0.2	0.0	0.1	0.2	0.6	1.1	2.4	5.3	19.8	51.7	108.3	188.5	288.3	408.0	506.1	547.7	503.0	337.5	62.7	61.5	7.6
1979	0.0	0.0	0.0	0.2	0.3	0.0	2.6	6.6	17.0	48.1	110.8	189.0	295.5	390.3	485.6	599.7	474.8	375.6	62.2	61.1	7.4
1980	0.0	0.0	0.0	0.0	0.0	0.6	2.3	2.8	21.0	49.9	114.6	204.4	273.6	413.5	525.8	593.0	546.8	439.6	64.6	63.2	7.7
1981	0.0	0.0	0.0	0.2	0.3	0.6	2.2	5.5	16.2	43.1	111.7	191.4	311.0	404.0	516.4	578.1	539.6	424.5	64.1	62.8	7.7
1982	0.0	0.0	0.0	0.0	0.0	0.0	1.7	7.7	16.5	47.2	98.8	178.3	308.8	440.8	527.7	559.4	549.0	523.1	64.7	63.5	7.8
1983	0.0	0.0	0.0	0.3	0.5	0.0	3.1	4.0	18.2	44.6	99.1	200.5	306.3	428.0	517.0	591.2	587.3	466.4	65.5	63.8	7.8
1979-83	0.0	0.0	0.0	0.1	0.2	0.2	2.4	5.3	17.8	46.6	106.9	192.7	299.6	415.6	514.9	584.0	541.4	446.8	64.3	63.0	7.7

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF TRACHEA, BRONCHUS AND LUNG (ICD9 162)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	8	12	6	15	28	25	32	31	32	17	22	8	0	236
1965	0	0	0	0	1	0	4	10	15	18	34	23	23	35	36	15	12	10	0	236
1966	0	0	0	0	0	1	4	4	16	25	35	38	22	48	25	20	22	10	0	270
1967	1	0	0	0	0	2	1	5	15	17	34	34	47	42	29	26	30	8	0	291
1968	0	1	0	1	1	2	6	13	23	22	34	34	39	31	31	35	20	18	0	311
1964-68	1	1	0	1	2	5	23	44	75	97	165	154	163	187	153	113	106	54	0	1344
1969	0	0	0	0	2	3	5	9	23	44	50	57	37	63	46	27	21	15	0	402
1970	0	0	0	0	1	0	1	11	24	39	37	64	42	44	44	27	20	11	0	365
1971	0	0	2	1	3	2	3	5	27	43	31	72	62	53	52	42	20	11	0	429
1972	0	0	1	0	0	1	2	11	25	45	48	69	80	58	62	48	31	20	0	501
1973	0	0	0	0	3	2	3	7	21	43	66	60	67	67	60	60	38	18	2	537
1969-73	0	0	3	1	9	8	14	43	120	214	232	342	288	285	264	204	130	75	2	2234
1974	1	0	0	1	3	4	3	13	26	55	68	87	99	76	70	49	36	26	4	621
1975	0	1	0	1	0	3	7	9	29	47	77	74	108	92	79	48	38	28	4	645
1976	0	0	1	0	1	2	9	9	23	37	90	111	123	107	79	58	44	21	0	715
1977	0	0	0	0	1	2	5	16	17	53	86	121	121	134	89	59	47	30	1	782
1978	0	0	0	0	1	1	4	7	35	45	91	141	147	138	108	80	42	32	0	872
1974-78	1	1	1	2	6	12	28	54	130	237	412	534	598	547	425	294	207	137	9	3635
1979	0	1	0	0	1	2	2	11	15	42	98	135	154	176	128	78	54	42	0	939
1980	1	0	0	0	1	1	5	11	41	63	110	141	180	152	119	88	53	39	1	1006
1981	0	0	0	0	3	2	7	10	41	74	124	170	195	191	145	78	68	32	0	1140
1982	1	0	1	0	1	2	5	13	25	75	114	165	180	204	155	101	64	46	1	1153
1983	0	0	0	2	3	2	4	12	42	63	124	195	213	220	214	110	71	64	4	1343
1979-83	2	1	1	2	9	9	23	57	164	317	570	806	922	943	761	455	310	223	6	5581

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF TRACHEA, BRONCHUS AND LUNG (ICD9 162)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	3.7	5.1	2.7	8.1	17.0	18.1	27.1	30.5	38.6	29.0	63.6	36.4	7.1	6.3	0.8
1965	0.0	0.0	0.0	0.0	0.4	0.0	1.8	4.3	6.5	9.5	20.0	16.2	19.0	33.8	42.1	24.8	33.4	43.5	6.9	6.3	0.8
1966	0.0	0.0	0.0	0.0	0.0	0.5	1.8	1.7	6.8	12.8	20.1	26.0	17.7	45.2	28.2	32.3	59.1	42.2	7.7	6.9	0.8
1967	0.3	0.0	0.0	0.0	0.0	0.9	0.4	2.2	6.2	8.4	19.0	22.5	36.6	38.7	32.2	40.2	76.9	31.9	8.1	7.2	0.8
1968	0.0	0.3	0.0	0.3	0.4	0.8	2.7	5.7	9.6	10.4	18.8	21.6	29.5	27.9	34.0	52.2	49.1	67.9	8.4	7.5	0.8
1964-68	0.1	0.1	0.0	0.1	0.2	0.5	2.1	3.8	6.4	9.8	19.0	21.0	26.1	35.2	34.9	36.1	56.6	44.9	7.7	6.9	0.8
1969	0.0	0.0	0.0	0.0	0.7	1.2	2.2	4.0	9.7	19.8	27.3	35.0	27.3	55.0	49.7	39.2	49.5	53.0	10.6	9.7	1.2
1970	0.0	0.0	0.0	0.0	0.3	0.0	0.4	4.8	10.2	16.9	19.8	38.0	30.1	37.3	46.5	38.3	45.2	36.3	9.3	8.5	1.0
1971	0.0	0.0	0.5	0.3	0.9	0.7	1.3	2.2	11.6	18.1	16.1	41.6	43.1	43.5	53.6	58.3	43.3	33.8	10.7	9.7	1.2
1972	0.0	0.0	0.3	0.0	0.0	0.3	0.8	4.8	10.8	19.0	23.5	39.5	53.7	46.3	62.6	64.6	65.3	58.3	12.2	11.0	1.3
1973	0.0	0.0	0.0	0.0	0.9	0.6	1.2	3.0	9.0	18.2	30.6	45.7	43.5	52.1	58.9	78.9	77.6	50.1	12.7	11.3	1.3
1969-73	0.0	0.0	0.2	0.1	0.6	0.6	1.2	3.8	10.3	18.4	23.6	40.0	39.9	46.9	54.4	56.4	56.7	46.5	11.1	10.1	1.2
1974	0.3	0.0	0.0	0.3	0.8	1.2	1.1	5.5	11.1	23.3	29.9	49.4	62.1	57.4	66.7	62.9	71.9	68.6	14.3	13.0	1.5
1975	0.0	0.3	0.0	0.3	0.0	0.9	2.5	3.8	12.4	20.0	32.9	40.8	65.1	68.0	73.1	60.0	73.6	71.1	14.5	13.2	1.6
1976	0.0	0.0	0.3	0.0	0.3	0.6	3.1	3.7	9.8	15.9	37.8	58.3	72.9	76.2	71.2	70.1	82.6	50.7	15.7	14.3	1.7
1977	0.0	0.0	0.0	0.0	0.3	0.6	1.6	6.4	7.3	22.8	36.4	59.8	70.7	92.0	77.5	69.7	84.8	69.3	16.7	15.2	1.9
1978	0.0	0.0	0.0	0.0	0.3	0.3	1.2	2.7	15.0	19.3	38.6	65.7	85.3	91.9	91.0	91.5	73.4	71.3	18.1	16.5	2.0
1974-78	0.1	0.1	0.1	0.1	0.3	0.7	1.9	4.4	11.1	20.3	35.2	55.4	71.4	77.7	76.2	71.2	77.4	66.2	15.9	14.5	1.8
1979	0.0	0.3	0.0	0.0	0.3	0.5	0.6	4.1	6.4	18.0	41.9	60.0	88.9	113.6	104.1	86.5	90.6	89.2	18.9	17.1	2.2
1980	0.3	0.0	0.0	0.0	0.3	0.3	1.4	3.9	17.1	27.1	47.1	61.3	100.7	94.4	94.3	94.3	85.8	79.4	19.9	18.4	2.2
1981	0.0	0.0	0.0	0.0	0.8	0.5	1.9	3.5	16.8	32.0	53.1	72.9	103.9	116.2	111.3	81.0	106.4	62.1	22.1	20.4	2.5
1982	0.3	0.0	0.3	0.0	0.2	0.5	1.4	4.2	9.9	32.4	48.8	70.7	90.8	122.0	115.3	100.9	97.9	85.2	21.7	19.8	2.5
1983	0.0	0.0	0.0	0.5	0.7	0.5	1.1	3.7	16.0	27.1	53.1	83.1	101.3	131.7	153.3	106.1	104.3	112.5	24.5	22.2	2.8
1979-83	0.1	0.1	0.1	0.1	0.5	0.5	1.3	3.8	13.3	27.3	48.8	69.7	97.2	115.8	116.5	94.1	97.2	86.2	21.5	19.6	2.4

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF PLEURA (ICD9 163)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1
1965	0	0	0	0	0	0	1	0	0	0	1	2	3	0	0	0	0	0	0	7
1966	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1
1967	0	0	0	0	0	0	0	0	0	0	0	0	3	2	0	0	1	0	0	6
1968	0	0	0	0	0	0	0	1	0	0	1	2	1	0	1	0	0	0	0	6
1964-68	0	0	0	0	0	0	1	1	0	0	2	4	7	2	3	0	1	0	0	21
1969	0	0	0	0	0	0	0	0	1	0	1	2	0	1	2	0	0	0	0	7
1970	0	0	0	0	0	0	0	0	0	0	1	2	1	0	1	0	1	1	0	7
1971	0	0	0	0	0	0	0	0	2	1	0	1	1	0	2	0	0	0	0	7
1972	0	0	0	0	0	0	1	1	2	1	1	0	0	2	1	0	0	0	0	9
1973	0	0	0	0	0	1	0	0	0	0	2	0	0	1	0	0	0	0	0	4
1969-73	0	0	0	0	0	1	1	1	5	2	5	5	2	4	6	0	1	1	0	34
1974	0	0	0	0	0	0	0	0	2	0	1	3	2	3	0	1	0	0	0	12
1975	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	3
1976	0	0	0	0	0	0	0	0	1	1	1	3	3	1	1	0	0	0	0	11
1977	0	0	0	0	0	0	0	0	1	0	0	1	1	2	2	2	3	1	0	13
1978	0	0	0	0	0	0	0	0	0	1	2	2	1	3	1	5	1	0	0	16
1974-78	0	0	0	0	0	0	0	0	4	2	4	9	10	9	4	8	4	1	0	55
1979	0	0	0	1	0	0	1	1	2	2	1	1	5	3	3	2	0	0	1	23
1980	0	0	0	0	0	0	0	0	2	1	2	4	5	1	6	2	1	0	0	24
1981	0	0	0	0	0	0	0	0	0	1	1	5	2	6	1	3	0	0	0	19
1982	0	0	0	0	0	1	0	0	4	1	3	6	4	3	3	1	0	0	0	26
1983	0	0	0	0	0	0	0	0	1	2	4	7	9	3	5	3	1	1	0	36
1979-83	0	0	0	1	0	1	1	1	9	7	11	23	25	16	18	11	2	1	1	128

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF PLEURA (ICD9 163)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.5	0.0	0.0	0.0	0.0	0.0	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.6	1.4	2.6	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	0.0	0.0	0.0	0.0	0.0	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.4	2.1	0.0	0.0	3.8	0.0	0.2	0.2	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	1.3	0.8	0.0	1.4	0.0	0.0	0.0	0.2	0.2	0.0
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.2	0.5	1.2	0.4	0.9	0.0	0.8	0.0	0.1	0.1	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	1.2	0.0	1.0	2.8	0.0	0.0	0.0	0.2	0.2	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.2	0.8	0.0	1.4	0.0	3.6	6.1	0.2	0.2	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	0.0	0.6	0.7	0.0	2.7	0.0	0.0	0.0	0.2	0.2	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.8	0.4	0.5	0.0	0.0	1.9	1.3	0.0	0.0	0.0	0.2	0.2	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	1.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.4	0.2	0.5	0.6	0.3	0.8	1.6	0.0	0.7	1.2	0.1	0.1	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.5	1.8	1.3	2.7	0.0	2.0	0.0	0.0	0.2	0.2	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.9	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	1.7	1.9	0.8	1.2	0.0	0.0	0.0	0.1	0.1	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	0.6	1.6	2.3	3.5	9.9	5.1	0.3	0.3	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	1.0	0.6	2.3	1.1	8.5	3.2	0.0	0.3	0.3	0.0
1974-78	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.2	0.4	1.0	1.3	1.5	0.9	2.9	2.7	1.0	0.4	0.3	0.0
1979	0.0	0.0	0.0	0.2	0.0	0.0	0.3	0.4	0.8	0.8	0.4	0.5	3.2	2.2	3.2	3.3	0.0	0.0	0.3	0.2	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	0.9	1.9	3.1	0.7	6.2	3.2	3.0	0.0	0.5	0.5	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.3	1.2	4.3	1.0	4.6	0.0	0.0	0.5	0.5	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	1.6	0.4	1.3	2.7	2.2	2.1	2.9	1.5	0.0	0.0	0.4	0.4	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.7	3.2	4.8	2.1	4.6	4.4	2.6	4.5	0.5	0.5	0.1
1979-83	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.7	0.6	0.9	2.1	2.9	2.3	3.6	3.4	1.2	0.9	0.7	0.7	0.1
																			0.5	0.5	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF PLEURA (ICD9 163)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1
1965	0	0	0	0	0	1	0	0	0	0	1	0	1	0	2	0	0	0	0	5
1966	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	0	0	3
1967	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	2
1968	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1
1964-68	0	0	0	0	0	1	0	1	0	0	2	0	2	1	2	2	1	0	0	12
1969	0	0	0	0	0	0	0	0	0	0	0	1	0	0	2	1	1	1	0	6
1970	0	0	0	0	0	1	0	0	0	1	1	0	3	0	0	0	1	0	0	7
1971	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1972	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	1	0	0	4
1973	0	0	0	0	0	0	0	1	0	0	0	1	0	0	1	0	0	1	0	4
1969-73	0	0	0	0	0	1	0	1	0	2	1	2	3	1	4	1	3	2	0	21
1974	0	0	0	0	0	0	0	0	0	1	1	0	0	2	1	0	0	1	0	6
1975	0	0	0	0	0	0	0	0	0	1	1	1	1	0	1	0	1	0	0	6
1976	0	0	0	0	0	1	0	0	1	0	1	1	1	0	0	1	0	0	0	6
1977	0	0	0	0	0	0	0	0	0	2	2	1	0	0	0	2	0	0	0	7
1978	0	0	0	0	0	0	1	0	0	0	0	0	1	1	1	0	0	0	0	4
1974-78	0	0	0	0	0	1	1	0	1	4	5	3	3	3	3	3	1	1	0	29
1979	0	0	0	0	0	0	0	0	0	1	0	2	2	1	2	0	1	0	0	9
1980	0	0	0	0	0	0	0	0	0	0	2	0	0	1	3	2	0	0	0	8
1981	0	0	0	0	0	0	0	0	2	0	2	2	2	2	0	1	3	1	0	15
1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	1	1	0	4
1983	0	0	0	0	0	0	0	0	0	0	0	1	2	2	4	1	0	2	0	12
1979-83	0	0	0	0	0	0	0	0	2	1	4	5	6	6	11	4	5	4	0	48

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF PLEURA (ICD9 163)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	0.0	0.0	0.0	0.0	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.6	0.0	0.8	0.0	2.3	0.0	0.0	0.0	0.1	0.1	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.6	0.0	0.0	0.0	0.0	1.6	0.0	0.0	0.1	0.1	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.9	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.5	0.0	0.0	0.0	0.0
1964-68	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.2	0.0	0.3	0.2	0.5	0.6	0.5	0.0	0.1	0.1	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	2.2	1.5	2.4	3.5	0.2	0.1	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.4	0.5	0.0	2.1	0.0	0.0	0.0	2.3	0.0	0.2	0.2	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.8	1.0	0.0	2.1	0.0	0.1	0.1	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.6	0.0	0.0	1.0	0.0	2.8	0.1	0.1	0.0	
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.2	0.1	0.2	0.4	0.2	0.8	0.3	1.3	1.2	0.1	0.1	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.0	1.5	1.0	0.0	0.0	2.6	0.1	0.1	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.6	0.6	0.0	0.9	0.0	1.9	0.0	0.1	0.1	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	0.4	0.5	0.6	0.0	0.0	1.2	0.0	0.0	0.1	0.1	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.8	0.5	0.0	0.0	0.0	2.4	0.0	0.0	0.2	0.1	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.6	0.7	0.8	0.0	0.0	0.0	0.1	0.1	0.0
1974-78	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.3	0.4	0.3	0.4	0.4	0.5	0.7	0.4	0.5	0.1	0.1	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	1.2	0.6	1.6	0.0	1.7	0.0	0.2	0.2	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.6	2.4	2.1	0.0	0.0	0.2	0.1	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.9	0.9	1.1	1.2	0.0	1.0	4.7	1.9	0.3	0.2	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.5	0.0	1.5	1.9	0.1	0.0	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.0	1.2	2.9	1.0	0.0	3.5	0.2	0.2	0.0
1979-83	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.1	0.3	0.4	0.6	0.7	1.7	0.8	1.6	1.5	0.2	0.2	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF THYMUS, HEART AND MEDIASTINUM (ICD9 164)

YEAR OF DIAGNOSIS	AGE (YEARS)																			ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	1	0	0	0	0	1	0	3	0	1	2	1	1	0	0	0	0	0	0	10
1965	1	0	0	3	0	1	0	0	2	0	1	0	1	1	0	0	1	0	0	11
1966	1	0	0	2	0	0	1	1	0	1	1	2	1	1	0	0	1	0	0	12
1967	0	0	0	0	0	2	0	0	2	0	0	0	1	0	0	1	1	0	0	7
1968	0	0	0	1	2	0	0	0	1	3	1	2	1	0	0	1	0	0	0	12
1964-68	3	0	0	6	2	4	1	4	5	5	5	5	5	2	0	2	3	0	0	52
1969	1	1	0	1	1	0	0	1	1	1	2	0	0	1	1	2	0	1	0	14
1970	1	1	0	0	0	2	2	0	3	1	1	2	2	1	1	0	0	0	0	17
1971	0	0	0	0	2	0	0	0	0	4	0	1	3	1	0	1	0	0	0	12
1972	0	0	1	1	2	0	0	2	1	0	2	0	0	1	1	0	0	0	0	11
1973	2	0	0	1	1	2	1	0	1	1	1	1	2	1	1	0	0	1	0	16
1969-73	4	2	1	3	6	4	3	3	6	7	6	4	7	5	4	3	0	2	0	70
1974	3	0	2	1	2	0	1	0	1	1	1	2	1	3	1	1	0	0	0	20
1975	3	0	0	1	1	1	3	1	1	3	2	0	3	2	2	3	0	0	0	26
1976	2	0	0	0	0	2	1	0	0	1	0	3	3	1	1	0	0	0	0	14
1977	0	2	0	1	0	1	1	1	1	0	0	1	0	1	0	0	1	0	0	10
1978	0	0	0	0	4	0	4	0	0	0	2	3	1	2	2	1	0	0	0	19
1974-78	8	2	2	3	7	4	10	2	3	5	5	9	8	9	6	5	1	0	0	89
1979	4	0	3	2	1	0	0	1	2	1	3	0	2	2	2	0	2	0	0	25
1980	1	0	0	1	1	1	0	1	0	2	0	0	1	2	3	0	1	0	0	14
1981	1	0	0	1	2	0	0	1	0	3	3	2	0	3	0	3	0	0	0	19
1982	0	0	0	1	2	2	1	3	1	1	1	2	2	1	1	1	0	0	0	19
1983	2	0	0	0	5	2	0	2	1	0	3	2	4	2	4	1	0	1	0	29
1979-83	8	0	3	5	11	5	1	8	4	7	10	6	9	10	10	5	3	1	0	106

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF THYMUS, HEART AND MEDIASTINUM (ICD9 164)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.0	0.0	0.0	0.0	0.5	0.0	1.3	0.0	0.5	1.2	0.7	0.9	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1965	0.3	0.0	0.0	1.0	0.0	0.5	0.0	0.0	0.9	0.0	0.6	0.0	0.9	1.1	0.0	0.0	3.9	0.0	0.3	0.3	0.0
1966	0.3	0.0	0.0	0.7	0.0	0.0	0.4	0.4	0.0	0.5	0.6	1.4	0.8	1.1	0.0	0.0	3.8	0.0	0.3	0.3	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.8	0.0	0.0	0.0	0.8	0.0	0.0	2.1	3.8	0.0	0.2	0.2	0.0
1968	0.0	0.0	0.0	0.3	0.7	0.0	0.0	0.0	0.4	1.4	0.5	1.3	0.8	0.0	0.0	2.1	0.0	0.0	0.3	0.3	0.0
1964-68	0.2	0.0	0.0	0.4	0.2	0.4	0.1	0.3	0.4	0.5	0.6	0.7	0.8	0.4	0.0	0.9	2.3	0.0	0.3	0.3	0.0
1969	0.3	0.2	0.0	0.3	0.3	0.0	0.0	0.4	0.4	0.5	1.1	0.0	0.0	1.0	1.4	4.2	0.0	6.4	0.4	0.4	0.0
1970	0.3	0.2	0.0	0.0	0.0	0.7	0.8	0.0	1.2	0.4	0.5	1.2	1.5	1.0	1.4	0.0	0.0	0.0	0.4	0.5	0.0
1971	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	1.7	0.0	0.6	2.2	0.9	0.0	2.0	0.0	0.0	0.3	0.3	0.0
1972	0.0	0.0	0.2	0.3	0.6	0.0	0.0	0.8	0.4	0.0	1.0	0.0	0.0	0.9	1.3	0.0	0.0	0.0	0.3	0.3	0.0
1973	0.6	0.0	0.0	0.3	0.3	0.6	0.4	0.0	0.4	0.4	0.5	0.6	1.4	0.9	1.3	0.0	0.0	5.5	0.4	0.4	0.0
1969-73	0.2	0.1	0.0	0.2	0.4	0.3	0.2	0.3	0.5	0.6	0.6	0.5	1.0	0.9	1.1	1.2	0.0	2.3	0.4	0.4	0.0
1974	0.9	0.0	0.5	0.3	0.6	0.0	0.4	0.0	0.4	0.4	0.5	1.2	0.7	2.7	1.2	2.0	0.0	0.0	0.5	0.5	0.0
1975	0.9	0.0	0.0	0.2	0.3	0.3	1.0	0.4	0.4	1.3	0.9	0.0	1.9	1.7	2.4	5.7	0.0	0.0	0.6	0.6	0.1
1976	0.6	0.0	0.0	0.0	0.0	0.6	0.3	0.0	0.0	0.4	0.0	1.7	1.9	0.8	1.2	0.0	0.0	0.0	0.3	0.4	0.0
1977	0.0	0.6	0.0	0.2	0.0	0.3	0.3	0.4	0.4	0.0	0.0	0.5	0.0	0.8	0.0	0.0	3.3	0.0	0.2	0.2	0.0
1978	0.0	0.0	0.0	0.0	1.0	0.0	1.2	0.0	0.0	0.0	0.9	1.5	0.6	1.5	2.2	1.7	0.0	0.0	0.4	0.4	0.0
1974-78	0.5	0.1	0.1	0.1	0.4	0.2	0.7	0.2	0.2	0.4	0.4	1.0	1.0	1.5	1.4	1.8	0.7	0.0	0.4	0.4	0.0
1979	1.3	0.0	0.8	0.5	0.3	0.0	0.0	0.4	0.8	0.4	1.3	0.0	1.3	1.5	2.1	0.0	6.3	0.0	0.6	0.6	0.1
1980	0.3	0.0	0.0	0.2	0.3	0.3	0.0	0.4	0.0	0.8	0.0	0.0	0.6	1.4	3.1	0.0	3.0	0.0	0.3	0.3	0.0
1981	0.3	0.0	0.0	0.2	0.5	0.0	0.0	0.3	0.0	1.3	1.3	0.9	0.0	2.1	0.0	4.6	0.0	0.0	0.4	0.4	0.0
1982	0.0	0.0	0.0	0.2	0.5	0.5	0.3	1.0	0.4	0.4	0.4	0.9	1.1	0.7	1.0	1.5	0.0	0.0	0.4	0.4	0.0
1983	0.6	0.0	0.0	0.0	1.2	0.5	0.0	0.6	0.4	0.0	1.3	0.9	2.1	1.4	3.7	1.5	0.0	4.5	0.6	0.6	0.1
1979-83	0.5	0.0	0.2	0.2	0.6	0.3	0.1	0.5	0.3	0.6	0.9	0.6	1.1	1.4	2.0	1.5	1.7	0.9	0.5	0.5	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF THYMUS, HEART AND MEDIASTINUM (ICD9 164)

YEAR OF DIAGNOSIS	AGE (YEARS)																			ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	1	1	0	5
1965	0	0	0	0	0	1	0	2	0	1	1	0	1	0	2	0	0	0	0	8
1966	1	1	0	0	0	0	0	0	1	1	2	2	1	0	0	1	0	1	0	11
1967	0	0	0	0	0	1	0	1	0	0	2	1	0	0	1	0	0	0	0	6
1968	1	0	0	1	0	1	1	1	2	0	0	3	2	0	1	0	0	0	0	13
1964-68	2	1	0	1	0	3	2	5	3	2	5	7	4	0	4	1	1	2	0	43
1969	1	0	0	1	0	1	0	0	0	0	0	0	1	0	1	0	0	2	0	7
1970	1	0	0	0	0	0	2	1	0	0	0	1	1	4	0	1	0	0	0	11
1971	0	0	0	0	1	0	0	2	1	1	0	0	1	0	0	1	0	1	0	8
1972	0	0	0	0	1	1	1	0	1	1	1	1	1	0	1	0	0	0	0	9
1973	0	0	0	0	1	1	0	0	0	0	1	3	0	0	1	0	0	0	0	7
1969-73	2	0	0	1	3	3	3	3	2	2	2	5	4	4	3	2	0	3	0	42
1974	1	0	0	1	0	2	0	0	0	0	2	1	2	2	0	0	1	0	0	12
1975	0	0	0	0	0	0	1	0	1	1	3	2	2	3	2	1	0	1	0	17
1976	0	0	0	1	0	0	0	0	0	2	1	0	1	0	1	1	0	0	0	7
1977	1	0	0	0	0	1	0	0	1	1	1	0	1	1	0	0	0	0	0	7
1978	1	0	0	1	0	0	1	1	0	0	1	2	3	0	2	1	0	0	0	13
1974-78	3	0	0	3	0	3	2	1	2	4	8	5	9	6	5	3	1	1	0	56
1979	3	0	0	0	0	0	1	2	1	2	0	1	2	0	0	0	1	0	0	13
1980	1	0	0	0	1	1	1	1	1	1	2	2	2	0	1	2	0	1	0	17
1981	0	1	1	1	0	2	0	0	2	0	3	2	2	0	2	1	1	0	0	18
1982	2	0	0	1	1	0	1	1	1	2	3	3	3	1	1	2	1	2	1	26
1983	1	1	0	0	0	1	1	1	0	3	0	1	2	2	3	3	0	0	0	19
1979-83	7	2	1	2	2	4	4	5	5	8	8	9	11	3	7	8	3	3	1	93

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF THYMUS, HEART AND MEDIASTINUM (ICD9 164)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.4	0.0	0.0	0.0	0.7	0.0	0.0	0.0	0.0	2.9	4.5	0.2	0.1	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.9	0.0	0.5	0.6	0.0	0.8	0.0	2.3	0.0	0.0	0.0	0.2	0.2	0.0
1966	0.3	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	1.1	1.4	0.8	0.0	0.0	1.6	0.0	4.2	0.3	0.3	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.0	0.0	1.1	0.7	0.0	0.0	1.1	0.0	0.0	0.0	0.2	0.2	0.0
1968	0.3	0.0	0.0	0.3	0.0	0.4	0.4	0.4	0.8	0.0	0.0	1.9	1.5	0.0	1.1	0.0	0.0	0.0	0.4	0.4	0.0
1964-68	0.1	0.1	0.0	0.1	0.0	0.3	0.2	0.4	0.3	0.2	0.6	1.0	0.6	0.0	0.9	0.3	0.5	1.7	0.2	0.2	0.0
1969	0.3	0.0	0.0	0.3	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.0	1.1	0.0	0.0	7.1	0.2	0.2	0.0
1970	0.3	0.0	0.0	0.0	0.0	0.0	0.9	0.4	0.0	0.0	0.0	0.6	0.7	3.4	0.0	1.4	0.0	0.0	0.3	0.3	0.0
1971	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.9	0.4	0.4	0.0	0.0	0.7	0.0	0.0	1.4	0.0	3.1	0.2	0.2	0.0
1972	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.0	0.4	0.4	0.5	0.6	0.7	0.0	1.0	0.0	0.0	0.0	0.2	0.2	0.0
1973	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.0	0.0	0.5	1.7	0.0	0.0	1.0	0.0	0.0	0.0	0.2	0.2	0.0
1969-73	0.1	0.0	0.0	0.1	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.6	0.6	0.7	0.6	0.6	0.0	1.9	0.2	0.2	0.0
1974	0.3	0.0	0.0	0.3	0.0	0.6	0.0	0.0	0.0	0.0	0.9	0.6	1.3	1.5	0.0	0.0	2.0	0.0	0.3	0.3	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.4	1.3	1.1	1.2	2.2	1.9	1.2	0.0	2.5	0.4	0.4	0.0
1976	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.9	0.4	0.0	0.6	0.0	0.9	1.2	0.0	0.0	0.2	0.1	0.0
1977	0.3	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	0.4	0.0	0.6	0.7	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1978	0.3	0.0	0.0	0.2	0.0	0.0	0.3	0.4	0.0	0.0	0.4	0.9	1.7	0.0	1.7	1.1	0.0	0.0	0.3	0.3	0.0
1974-78	0.2	0.0	0.0	0.2	0.0	0.2	0.1	0.1	0.2	0.3	0.7	0.5	1.1	0.9	0.9	0.7	0.4	0.5	0.3	0.2	0.0
1979	1.0	0.0	0.0	0.0	0.0	0.0	0.3	0.7	0.4	0.9	0.0	0.4	1.2	0.0	0.0	0.0	1.7	0.0	0.3	0.3	0.0
1980	0.3	0.0	0.0	0.0	0.3	0.3	0.3	0.4	0.4	0.4	0.9	0.9	1.1	0.0	0.8	2.1	0.0	2.0	0.4	0.3	0.0
1981	0.0	0.3	0.3	0.3	0.0	0.5	0.0	0.0	0.8	0.0	1.3	0.9	1.1	0.0	1.5	1.0	1.6	0.0	0.4	0.4	0.0
1982	0.7	0.0	0.0	0.3	0.2	0.0	0.3	0.3	0.4	0.9	1.3	1.3	1.5	0.6	0.7	2.0	1.5	3.7	0.5	0.5	0.0
1983	0.3	0.3	0.0	0.0	0.0	0.3	0.3	0.3	0.0	1.3	0.0	0.4	1.0	1.2	2.1	2.9	0.0	0.0	0.4	0.4	0.0
1979-83	0.5	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.4	0.7	0.7	0.8	1.2	0.4	1.1	1.7	0.9	1.2	0.4	0.4	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE RESPIRATORY SYSTEM AND INTRATHORACIC ORGANS (ICD9 165)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	1	0	2	1	0	1	0	0	0	0	5
1965	0	0	0	0	0	0	0	0	1	1	1	1	2	4	1	1	0	0	0	12
1966	0	0	0	0	0	0	0	0	0	0	1	1	2	3	0	0	0	0	0	7
1967	0	1	0	0	0	1	0	1	1	0	2	1	3	1	1	2	1	0	0	15
1968	0	0	0	0	0	0	0	2	0	1	0	3	1	3	1	0	0	0	0	11
1964-68	0	1	0	0	0	1	0	3	2	3	4	8	9	11	4	3	1	0	0	50
1969	0	0	0	0	0	0	0	0	2	1	1	1	1	0	0	0	0	0	0	6
1970	0	0	0	0	0	0	0	0	0	1	1	3	1	1	0	3	0	1	0	11
1971	0	0	0	0	0	0	1	0	0	0	0	4	1	4	1	0	0	0	0	11
1972	0	0	0	0	0	0	0	0	0	0	2	2	3	6	3	0	0	0	0	16
1973	0	0	0	0	0	0	0	0	0	1	3	2	0	1	1	1	1	0	0	10
1969-73	0	0	0	0	0	0	1	0	2	3	7	12	6	12	5	4	1	1	0	54
1974	0	0	0	0	0	0	0	0	1	1	5	3	0	1	1	1	1	0	0	14
1975	0	0	0	0	0	0	0	0	1	0	1	2	4	6	1	1	0	1	0	17
1976	0	0	0	0	0	0	0	0	0	0	0	4	4	5	2	2	0	0	0	17
1977	0	0	0	0	0	0	0	0	0	1	3	2	1	3	2	1	0	0	0	13
1978	0	0	0	0	0	0	0	0	1	0	2	2	4	2	3	2	0	0	0	16
1974-78	0	0	0	0	0	0	0	0	3	2	11	13	13	17	9	7	1	1	0	77
1979	0	0	0	0	0	0	0	0	1	0	0	4	6	2	0	1	1	1	0	16
1980	0	0	0	0	0	0	1	0	0	0	3	4	2	4	4	0	0	0	0	18
1981	0	0	0	0	0	0	1	0	0	1	3	0	3	2	0	2	1	0	0	13
1982	0	0	0	0	0	0	0	2	1	0	0	3	2	1	2	1	0	0	0	12
1983	0	0	0	0	0	0	0	0	0	1	2	2	3	8	3	1	2	0	0	22
1979-83	0	0	0	0	0	0	2	2	2	2	8	13	16	17	9	5	4	1	0	81

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE RESPIRATORY SYSTEM AND INTRATHORACIC ORGANS (ICD9 165)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	1.4	0.9	0.0	1.5	0.0	0.0	0.0	0.2	0.2	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.6	0.7	1.7	4.4	1.4	2.2	0.0	0.0	0.4	0.4	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	1.7	3.2	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1967	0.0	0.3	0.0	0.0	0.0	0.4	0.0	0.4	0.4	0.0	1.1	0.7	2.4	1.1	1.4	4.2	3.8	0.0	0.4	0.4	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.5	0.0	1.9	0.8	3.1	1.4	0.0	0.0	0.0	0.3	0.3	0.0
1964-68	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.3	0.2	0.3	0.5	1.1	1.5	2.4	1.1	1.3	0.8	0.0	0.3	0.3	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.5	0.5	0.6	0.8	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	1.8	0.8	1.0	0.0	6.2	0.0	6.1	0.3	0.3	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	2.3	0.7	3.8	1.3	0.0	0.0	0.0	0.3	0.3	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	1.2	2.1	5.6	3.9	0.0	0.0	0.0	0.4	0.4	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.4	1.2	0.0	0.9	1.3	2.0	3.4	0.0	0.2	0.2	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.3	0.7	1.4	0.9	2.3	1.3	1.6	0.7	1.2	0.3	0.3	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.3	1.8	0.0	0.9	1.2	2.0	3.4	0.0	0.3	0.3	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	1.2	2.6	5.2	1.2	1.9	0.0	5.3	0.4	0.4	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.2	2.5	4.1	2.3	3.7	0.0	0.0	0.4	0.4	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	1.1	0.6	2.4	2.3	1.8	0.0	0.0	0.3	0.3	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	1.0	2.5	1.5	3.3	3.4	0.0	0.0	0.4	0.4	0.0
1974-78	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	1.0	1.4	1.7	2.8	2.1	2.6	0.7	1.0	0.4	0.4	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.9	3.8	1.5	0.0	1.6	3.1	4.9	0.3	0.4	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	1.3	1.9	1.2	2.9	4.1	0.0	0.0	0.0	0.4	0.4	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	1.3	0.0	1.8	1.4	0.0	3.1	2.9	0.0	0.3	0.3	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.4	0.0	0.0	1.4	1.1	0.7	1.9	1.5	0.0	0.0	0.2	0.2	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	0.9	1.6	5.6	2.7	1.5	5.3	0.0	0.4	0.4	0.1
1979-83	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.2	0.7	1.2	1.9	2.4	1.8	1.5	2.3	0.9	0.3	0.3	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE RESPIRATORY SYSTEM AND INTRATHORACIC ORGANS (ICD9 165)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1
1965	0	0	0	0	0	0	0	1	0	1	0	0	0	0	1	0	0	0	0	3
1966	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	2
1967	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	2
1968	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	2
1964-68	0	0	0	0	0	0	0	2	0	3	0	2	0	0	2	0	1	0	0	10
1969	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	1	0	3
1970	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1
1971	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1
1972	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1
1973	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	2
1969-73	0	0	0	0	0	1	0	0	0	2	1	0	1	0	2	0	0	1	0	8
1974	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	0	0	3
1975	0	0	0	0	0	0	0	0	0	0	1	1	1	0	0	1	0	0	0	4
1976	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	4
1977	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	2
1978	0	0	0	0	0	1	0	0	0	0	0	0	1	1	0	0	0	1	0	4
1974-78	0	0	0	1	0	1	0	0	0	2	1	2	4	2	0	2	1	1	0	17
1979	0	0	0	0	0	0	0	0	0	0	0	1	1	1	0	0	1	0	0	4
1980	0	0	0	0	0	0	0	0	0	0	2	0	3	0	0	0	0	1	0	6
1981	0	0	0	0	0	0	0	0	1	0	0	0	0	3	0	0	0	0	0	4
1982	0	0	0	0	0	0	1	0	0	0	1	2	0	0	0	0	0	0	0	4
1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1
1979-83	0	0	0	0	0	0	1	0	1	0	3	3	4	4	0	1	1	1	0	19

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES WITHIN THE RESPIRATORY SYSTEM AND INTRATHORACIC ORGANS (ICD9 165)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.9	0.0	0.0	0.0	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	0.0	0.0	0.0	0.0	1.2	0.0	0.0	0.0	0.1	0.1	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0	0.0	0.1	0.0	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.3	0.0	0.3	0.0	0.0	0.5	0.0	0.5	0.0	0.1	0.1	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	1.1	0.0	0.0	3.5	0.1	0.1	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.2	0.1	0.0	0.1	0.0	0.4	0.0	0.0	0.6	0.0	0.0	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.6	0.6	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.6	0.6	0.0	0.0	1.2	0.0	0.0	0.1	0.1	0.0
1976	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.0	1.2	1.9	0.0	0.1	0.1	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	0.0	0.0	0.0	2.2	0.1	0.1	0.0
1974-78	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.2	0.1	0.2	0.5	0.3	0.0	0.5	0.4	0.5	0.1	0.1	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.6	0.6	0.0	0.0	1.7	0.0	0.1	0.1	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	1.7	0.0	0.0	0.0	0.0	2.0	0.1	0.1	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	1.8	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0
1979-83	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.3	0.3	0.4	0.5	0.0	0.2	0.3	0.4	0.1	0.1	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF BONE AND ARTICULAR CARTILAGE (ICD9 170)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	6	8	4	1	3	0	2	4	4	4	3	4	5	1	1	0	0	50
1965	0	2	3	6	2	1	0	0	3	2	2	1	3	2	2	2	2	2	0	35
1966	1	6	4	4	2	0	0	0	4	1	3	3	1	2	1	5	3	0	0	40
1967	1	4	4	3	1	1	2	4	0	3	1	3	0	2	2	1	0	2	0	34
1968	0	3	4	5	3	1	0	1	1	1	0	3	3	5	3	2	3	1	0	39
1964-68	2	15	21	26	12	4	5	5	10	11	10	14	10	15	13	11	9	5	0	198
1969	0	4	5	4	4	1	2	2	6	1	6	4	1	1	2	6	2	1	0	52
1970	0	0	4	5	1	3	1	1	1	8	2	3	3	5	6	5	1	0	0	49
1971	3	1	5	7	2	1	0	1	2	2	4	2	4	6	5	6	3	1	0	55
1972	0	0	1	9	4	2	1	1	2	3	3	3	3	3	3	0	1	3	0	42
1973	1	3	1	14	5	1	4	1	3	1	2	0	4	10	3	2	3	1	0	59
1969-73	4	8	16	39	16	8	8	6	14	15	17	12	15	25	19	19	10	6	0	257
1974	0	1	5	10	12	6	1	5	2	3	7	5	5	5	4	4	8	3	1	87
1975	0	1	11	6	7	2	2	3	2	4	5	4	7	5	5	2	3	1	0	70
1976	0	3	8	14	6	9	5	3	5	1	6	5	11	7	7	4	3	5	1	103
1977	0	2	5	7	6	6	1	1	0	1	4	5	7	6	3	3	3	2	1	63
1978	1	5	4	11	9	3	2	2	0	1	7	5	5	3	2	3	3	2	0	68
1974-78	1	12	33	48	40	26	11	14	9	10	29	24	35	26	21	16	20	13	3	391
1979	0	3	5	10	5	3	1	2	5	6	4	1	2	3	4	1	3	4	0	62
1980	1	2	6	12	9	4	2	2	1	2	0	5	2	3	5	3	2	1	0	62
1981	0	2	3	6	10	4	1	2	4	3	2	9	4	9	5	3	2	4	0	73
1982	0	2	4	7	3	2	3	3	3	4	4	2	2	3	4	6	0	0	0	52
1983	0	2	7	9	6	5	3	0	3	7	2	5	4	5	2	2	1	0	0	63
1979-83	1	11	25	44	33	18	10	9	16	22	12	22	14	23	20	15	8	9	0	312

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF BONE AND ARTICULAR CARTILAGE (ICD9 170)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	1.8	2.9	1.9	0.5	1.3	0.0	0.9	2.1	2.4	2.9	2.6	4.4	7.3	2.2	3.9	0.0	1.5	1.5	0.2
1965	0.0	0.5	0.9	2.1	0.9	0.5	0.0	0.0	1.3	1.0	1.1	0.7	2.6	2.2	2.9	4.4	7.7	14.4	1.0	1.0	0.1
1966	0.3	1.5	1.1	1.3	0.8	0.0	0.0	0.0	1.7	0.5	1.7	2.0	0.8	2.1	1.4	10.7	11.5	0.0	1.1	1.1	0.1
1967	0.3	1.0	1.1	0.9	0.4	0.4	0.9	1.7	0.0	1.5	0.5	2.0	0.0	2.1	2.8	2.1	0.0	13.8	1.0	0.9	0.1
1968	0.0	0.7	1.2	1.5	1.1	0.4	0.0	0.4	0.4	0.5	0.0	1.9	2.4	5.1	4.2	4.2	11.2	6.7	1.1	1.0	0.1
1964-68	0.1	0.8	1.2	1.7	1.0	0.4	0.4	0.4	0.9	1.1	1.1	1.9	1.7	3.2	3.7	4.7	6.9	7.0	1.1	1.1	0.1
1969	0.0	1.0	1.3	1.2	1.3	0.4	0.9	0.8	2.5	0.5	3.3	2.5	0.8	1.0	2.8	12.5	7.4	6.4	1.4	1.3	0.1
1970	0.0	0.0	1.0	1.4	0.3	1.1	0.4	0.4	0.4	3.5	1.1	1.8	2.3	4.9	8.2	10.3	3.6	0.0	1.3	1.3	0.1
1971	0.9	0.2	1.2	1.9	0.6	0.3	0.0	0.4	0.8	0.9	2.1	1.2	2.9	5.7	6.7	12.3	10.6	5.8	1.4	1.4	0.1
1972	0.0	0.0	0.2	2.4	1.2	0.7	0.4	0.4	0.8	1.3	1.5	1.8	2.1	2.8	3.9	0.0	3.5	16.8	1.1	1.1	0.1
1973	0.3	0.8	0.2	3.7	1.5	0.3	1.5	0.4	1.2	0.4	1.0	0.0	2.7	9.1	3.8	4.0	10.3	5.5	1.5	1.5	0.1
1969-73	0.2	0.4	0.8	2.2	1.0	0.6	0.7	0.5	1.2	1.3	1.8	1.4	2.2	4.7	5.1	7.8	7.1	7.0	1.4	1.3	0.1
1974	0.0	0.3	1.2	2.5	3.4	1.8	0.4	2.1	0.8	1.3	3.2	3.0	3.3	4.4	4.9	7.9	27.4	16.2	2.2	2.0	0.2
1975	0.0	0.3	2.6	1.5	1.9	0.6	0.7	1.2	0.8	1.7	2.2	2.3	4.5	4.3	5.9	3.8	10.3	5.3	1.7	1.6	0.2
1976	0.0	0.9	2.0	3.4	1.6	2.5	1.7	1.2	2.1	0.4	2.6	2.8	7.0	5.8	8.1	7.4	10.1	25.8	2.5	2.3	0.2
1977	0.0	0.6	1.3	1.7	1.6	1.7	0.3	0.4	0.0	0.4	1.8	2.6	4.4	4.8	3.4	5.3	9.9	10.2	1.5	1.4	0.1
1978	0.3	1.5	1.0	2.6	2.4	0.9	0.6	0.8	0.0	0.4	3.1	2.5	3.2	2.3	2.2	5.1	9.7	10.0	1.6	1.5	0.1
1974-78	0.1	0.7	1.6	2.4	2.2	1.5	0.7	1.1	0.7	0.8	2.6	2.6	4.5	4.3	4.9	5.9	13.4	13.5	1.9	1.8	0.2
1979	0.0	0.9	1.4	2.4	1.3	0.8	0.3	0.7	2.1	2.5	1.7	0.5	1.3	2.2	4.3	1.6	9.4	19.5	1.5	1.4	0.1
1980	0.3	0.6	1.7	2.9	2.3	1.1	0.6	0.7	0.4	0.8	0.0	2.3	1.2	2.2	5.2	4.8	6.0	4.8	1.5	1.3	0.1
1981	0.0	0.6	0.9	1.5	2.5	1.1	0.3	0.7	1.6	1.3	0.8	4.2	2.4	6.4	5.0	4.6	5.9	18.9	1.7	1.6	0.1
1982	0.0	0.6	1.2	1.7	0.7	0.5	0.9	1.0	1.2	1.7	1.7	0.9	1.1	2.1	3.8	9.0	0.0	0.0	1.2	1.1	0.1
1983	0.0	0.6	2.1	2.3	1.5	1.3	0.9	0.0	1.1	3.0	0.8	2.3	2.1	3.5	1.8	2.9	2.6	0.0	1.4	1.4	0.1
1979-83	0.1	0.7	1.4	2.2	1.7	1.0	0.6	0.6	1.3	1.9	1.0	2.0	1.6	3.3	4.0	4.6	4.6	8.5	1.4	1.4	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF BONE AND ARTICULAR CARTILAGE (ICD9 170)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	1	3	4	1	3	0	1	0	1	2	3	5	2	1	1	0	1	0	29
1965	2	2	1	6	0	0	1	1	0	2	1	0	1	2	2	1	2	3	0	27
1966	1	1	2	1	0	0	0	0	1	2	0	0	1	0	1	1	2	1	0	14
1967	0	3	3	0	0	2	0	0	2	0	0	2	1	2	1	1	2	2	0	21
1968	0	1	6	3	1	0	2	2	3	2	0	0	2	1	4	1	1	2	0	31
1964-68	3	8	15	14	2	5	3	4	6	7	3	5	10	7	9	5	7	9	0	122
1969	2	5	2	1	2	1	1	1	2	4	2	4	1	1	0	3	2	2	0	36
1970	1	0	13	2	0	1	0	0	0	2	4	1	4	4	2	1	3	2	0	40
1971	1	3	0	3	4	0	4	0	3	0	1	3	4	1	1	3	2	3	0	36
1972	2	5	8	6	2	4	1	0	0	0	4	0	3	5	5	2	3	2	0	52
1973	1	4	7	4	2	0	0	2	1	4	3	5	4	4	4	1	0	3	0	49
1969-73	7	17	30	16	10	6	6	3	6	10	14	13	16	15	12	10	10	12	0	213
1974	0	1	4	5	4	4	2	2	1	1	2	2	3	8	10	7	3	3	0	62
1975	0	3	7	9	3	6	3	3	1	2	1	2	5	3	1	3	2	5	1	60
1976	0	2	2	9	3	4	5	2	5	5	2	3	7	8	2	5	4	2	1	71
1977	0	1	3	8	8	5	2	1	0	2	1	2	1	2	7	7	2	4	0	56
1978	0	3	3	8	6	1	0	1	2	0	2	1	2	3	2	5	4	3	0	46
1974-78	0	10	19	39	24	20	12	9	9	10	8	10	18	24	22	27	15	17	2	295
1979	1	0	3	6	1	6	3	2	2	4	2	3	2	5	2	2	1	3	0	48
1980	1	3	2	4	4	2	2	1	1	0	1	1	1	2	3	2	3	2	0	35
1981	0	1	7	11	4	2	5	3	1	4	4	3	4	4	3	3	3	3	0	65
1982	0	0	6	6	6	1	3	0	0	2	0	3	2	6	5	3	6	0	1	50
1983	0	2	6	5	3	3	2	2	0	1	1	0	2	6	4	3	0	4	0	44
1979-83	2	6	24	32	18	14	15	8	4	11	8	10	11	23	17	13	13	12	1	242

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF BONE AND ARTICULAR CARTILAGE (ICD9 170)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.3	0.9	1.5	0.5	1.4	0.0	0.4	0.0	0.5	1.2	2.2	4.2	2.0	1.2	1.7	0.0	4.5	0.9	0.9	0.1
1965	0.5	0.5	0.3	2.1	0.0	0.0	0.5	0.4	0.0	1.1	0.6	0.0	0.8	1.9	2.3	1.7	5.6	13.0	0.8	0.7	0.1
1966	0.3	0.3	0.6	0.3	0.0	0.0	0.0	0.0	0.4	1.0	0.0	0.0	0.8	0.0	1.1	1.6	5.4	4.2	0.4	0.3	0.0
1967	0.0	0.8	0.9	0.0	0.0	0.9	0.0	0.0	0.8	0.0	0.0	1.3	0.8	1.8	1.1	1.5	5.1	8.0	0.6	0.5	0.0
1968	0.0	0.3	1.7	0.9	0.4	0.0	0.9	0.9	1.2	0.9	0.0	0.0	1.5	0.9	4.4	1.5	2.5	7.5	0.8	0.8	0.1
1964-68	0.2	0.4	0.9	0.9	0.2	0.5	0.3	0.3	0.5	0.7	0.3	0.7	1.6	1.3	2.1	1.6	3.7	7.5	0.7	0.6	0.1
1969	0.6	1.3	0.5	0.3	0.7	0.4	0.4	0.4	0.8	1.8	1.1	2.5	0.7	0.9	0.0	4.4	4.7	7.1	1.0	0.9	0.1
1970	0.3	0.0	3.4	0.6	0.0	0.4	0.0	0.0	0.0	0.9	2.1	0.6	2.9	3.4	2.1	1.4	6.8	6.6	1.0	1.0	0.1
1971	0.3	0.8	0.0	0.9	1.2	0.0	1.7	0.0	1.3	0.0	0.5	1.7	2.8	0.8	1.0	4.2	4.3	9.2	0.9	0.8	0.1
1972	0.6	1.4	2.1	1.7	0.6	1.3	0.4	0.0	0.0	0.0	2.0	0.0	2.0	4.0	5.0	2.7	6.3	5.8	1.3	1.2	0.1
1973	0.3	1.1	1.8	1.1	0.6	0.0	0.0	0.9	0.4	1.7	1.4	2.9	2.6	3.1	3.9	1.3	0.0	8.4	1.2	1.2	0.1
1969-73	0.4	0.9	1.6	0.9	0.6	0.4	0.5	0.3	0.5	0.9	1.4	1.5	2.2	2.5	2.5	2.8	4.4	7.4	1.1	1.0	0.1
1974	0.0	0.3	1.0	1.3	1.1	1.2	0.7	0.9	0.4	0.4	0.9	1.1	1.9	6.0	9.5	9.0	6.0	7.9	1.5	1.3	0.1
1975	0.0	0.9	1.8	2.3	0.8	1.7	1.1	1.3	0.4	0.9	0.4	1.1	3.0	2.2	0.9	3.7	3.9	12.7	1.4	1.3	0.1
1976	0.0	0.6	0.5	2.3	0.8	1.1	1.7	0.8	2.1	2.1	0.8	1.6	4.1	5.7	1.8	6.0	7.5	4.8	1.6	1.5	0.1
1977	0.0	0.3	0.8	2.0	2.1	1.4	0.6	0.4	0.0	0.9	0.4	1.0	0.6	1.4	6.1	8.3	3.6	9.2	1.3	1.1	0.1
1978	0.0	0.9	0.8	2.0	1.5	0.3	0.0	0.4	0.9	0.0	0.8	0.5	1.2	2.0	1.7	5.7	7.0	6.7	1.1	0.9	0.1
1974-78	0.0	0.6	1.0	2.0	1.3	1.1	0.8	0.7	0.8	0.9	0.7	1.0	2.1	3.4	3.9	6.5	5.6	8.2	1.3	1.2	0.1
1979	0.3	0.0	0.9	1.5	0.3	1.6	0.9	0.7	0.8	1.7	0.9	1.3	1.2	3.2	1.6	2.2	1.7	6.4	1.0	1.0	0.1
1980	0.3	1.0	0.6	1.0	1.0	0.5	0.6	0.4	0.4	0.0	0.4	0.4	0.6	1.2	2.4	2.1	4.9	4.1	0.8	0.7	0.1
1981	0.0	0.3	2.1	2.8	1.0	0.5	1.4	1.0	0.4	1.7	1.7	1.3	2.1	2.4	2.3	3.1	4.7	5.8	1.4	1.3	0.1
1982	0.0	0.0	1.8	1.6	1.5	0.3	0.8	0.0	0.0	0.9	0.0	1.3	1.0	3.6	3.7	3.0	9.2	0.0	1.1	0.9	0.1
1983	0.0	0.7	1.9	1.4	0.7	0.8	0.5	0.6	0.0	0.4	0.4	0.0	1.0	3.6	2.9	2.9	0.0	7.0	0.9	0.9	0.1
1979-83	0.1	0.4	1.4	1.6	0.9	0.7	0.8	0.5	0.3	0.9	0.7	0.9	1.2	2.8	2.6	2.7	4.1	4.6	1.0	0.9	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF CONNECTIVE AND OTHER SOFT TISSUE (ICD9 171)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	7	2	2	3	2	1	2	6	4	8	5	3	4	12	3	2	1	1	0	68
1965	5	0	1	2	3	2	6	7	3	10	2	2	5	8	8	8	3	3	0	78
1966	4	3	1	0	3	3	2	7	2	4	7	8	7	3	1	4	4	2	0	65
1967	9	2	2	2	1	4	1	4	6	9	5	5	5	5	4	6	6	2	0	78
1968	8	5	3	4	2	2	3	0	2	7	7	4	11	5	7	2	2	4	0	78
1964-68	33	12	9	11	11	12	14	24	17	38	26	22	32	33	23	22	16	12	0	367
1969	10	3	4	4	4	3	3	6	6	7	4	4	7	15	6	8	5	1	0	100
1970	7	2	5	6	10	0	4	1	3	6	5	5	5	7	6	2	3	5	0	82
1971	11	4	4	3	1	4	4	7	2	7	3	7	4	7	7	5	2	0	0	82
1972	7	5	0	3	4	4	4	3	8	6	8	10	10	7	11	8	8	1	0	107
1973	8	3	5	5	5	3	4	2	2	8	13	7	6	11	13	1	6	3	0	105
1969-73	43	17	18	21	24	14	19	19	21	34	33	33	32	47	43	24	24	10	0	476
1974	4	1	4	2	8	10	5	8	10	9	11	9	11	11	11	4	6	5	3	132
1975	12	6	4	4	1	6	3	6	5	11	6	8	7	7	9	7	7	4	0	113
1976	7	3	4	7	4	9	10	6	5	7	9	12	12	16	12	9	4	3	0	139
1977	9	3	6	9	2	4	8	5	8	6	7	9	9	10	9	8	4	3	1	120
1978	7	1	1	3	3	6	6	4	3	14	8	5	9	14	13	13	8	5	1	124
1974-78	39	14	19	25	18	35	32	29	31	47	41	43	48	58	54	41	29	20	5	628
1979	2	2	2	2	6	5	10	9	7	11	9	12	8	12	6	4	3	7	0	117
1980	1	1	2	4	4	3	6	7	5	8	11	13	12	9	9	6	3	6	0	110
1981	1	2	5	5	7	3	4	4	7	6	3	15	13	14	8	10	4	2	1	114
1982	3	1	3	6	4	9	4	8	7	8	7	8	10	13	9	13	5	2	0	120
1983	5	1	1	5	9	9	4	10	4	5	11	15	15	17	16	13	8	4	0	152
1979-83	12	7	13	22	30	29	28	38	30	38	41	63	58	65	48	46	23	21	1	613

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF CONNECTIVE AND OTHER SOFT TISSUE (ICD9 171)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	1.8	0.5	0.6	1.1	0.9	0.5	0.9	2.5	1.8	4.2	2.9	2.1	3.5	13.3	4.4	4.4	3.9	7.4	2.0	2.1	0.2
1965	1.3	0.0	0.3	0.7	1.3	1.0	2.7	2.9	1.3	5.2	1.1	1.4	4.3	8.7	11.5	17.4	11.6	21.6	2.3	2.3	0.2
1966	1.0	0.8	0.3	0.0	1.2	1.4	0.9	2.9	0.9	2.0	3.9	5.4	5.9	3.2	1.4	8.6	15.4	14.3	1.9	1.8	0.2
1967	2.4	0.5	0.6	0.6	0.4	1.8	0.4	1.7	2.5	4.5	2.7	3.3	4.1	5.3	5.6	12.6	22.7	13.8	2.2	2.2	0.2
1968	2.2	1.2	0.8	1.2	0.7	0.8	1.3	0.0	0.8	3.4	3.8	2.5	8.7	5.1	9.8	4.2	7.5	26.7	2.2	2.2	0.2
1964-68	1.7	0.6	0.5	0.7	0.9	1.1	1.2	2.0	1.5	3.9	2.9	3.0	5.3	7.1	6.6	9.4	12.3	16.9	2.1	2.1	0.2
1969	2.9	0.7	1.0	1.2	1.3	1.2	1.3	2.5	2.5	3.2	2.2	2.5	5.4	15.0	8.4	16.6	18.4	6.4	2.7	2.7	0.3
1970	2.1	0.5	1.3	1.7	3.1	0.0	1.7	0.4	1.2	2.7	2.7	3.0	3.8	6.8	8.2	4.1	10.8	30.5	2.2	2.2	0.2
1971	3.4	1.0	1.0	0.8	0.3	1.4	1.7	2.9	0.8	3.0	1.6	4.1	2.9	6.6	9.4	10.2	7.1	0.0	2.1	2.2	0.2
1972	2.2	1.3	0.0	0.8	1.2	1.3	1.6	1.3	3.3	2.6	4.0	5.9	7.1	6.5	14.3	16.2	27.9	5.6	2.7	2.7	0.3
1973	2.5	0.8	1.2	1.3	1.5	0.9	1.5	0.8	0.8	3.4	6.2	4.1	4.1	10.0	16.3	2.0	20.6	16.5	2.6	2.7	0.3
1969-73	2.6	0.9	0.9	1.2	1.5	1.0	1.6	1.6	1.7	3.0	3.4	3.9	4.7	8.9	11.5	9.8	17.0	11.7	2.5	2.5	0.2
1974	1.2	0.3	1.0	0.5	2.3	3.0	1.8	3.3	4.1	3.8	5.0	5.3	7.3	9.7	13.4	7.9	20.5	27.0	3.2	3.1	0.3
1975	3.7	1.7	1.0	1.0	0.3	1.7	1.0	2.5	2.1	4.6	2.7	4.7	4.5	6.0	10.7	13.4	24.0	21.1	2.8	2.8	0.2
1976	2.2	0.9	1.0	1.7	1.1	2.5	3.4	2.4	2.1	2.9	4.0	6.7	7.6	13.3	13.9	16.5	13.5	15.5	3.3	3.3	0.3
1977	2.9	0.9	1.5	2.2	0.5	1.1	2.5	2.0	3.3	2.5	3.1	4.8	5.7	7.9	10.2	14.2	13.2	15.2	2.8	2.8	0.3
1978	2.3	0.3	0.3	0.7	0.8	1.7	1.8	1.5	1.3	5.8	3.5	2.5	5.7	10.8	14.3	22.1	26.0	25.0	2.8	2.8	0.3
1974-78	2.5	0.8	0.9	1.2	1.0	2.0	2.1	2.3	2.6	3.9	3.6	4.7	6.2	9.6	12.5	15.0	19.5	20.7	3.0	3.0	0.3
1979	0.6	0.6	0.5	0.5	1.6	1.4	2.9	3.3	2.9	4.6	3.9	5.8	5.1	8.9	6.4	6.5	9.4	34.1	2.6	2.6	0.2
1980	0.3	0.3	0.6	1.0	1.0	0.8	1.7	2.5	2.1	3.4	4.7	6.1	7.4	6.5	9.3	9.5	9.1	29.0	2.4	2.4	0.2
1981	0.3	0.6	1.4	1.2	1.8	0.8	1.1	1.4	2.8	2.5	1.3	7.0	7.7	9.9	7.9	15.5	11.7	9.4	2.5	2.4	0.2
1982	1.0	0.3	0.9	1.5	1.0	2.5	1.1	2.6	2.8	3.4	3.0	3.7	5.6	9.1	8.6	19.5	14.0	9.3	2.6	2.5	0.2
1983	1.6	0.3	0.3	1.3	2.2	2.4	1.1	3.0	1.5	2.1	4.6	6.8	8.0	11.9	14.7	19.0	21.2	17.9	3.2	3.1	0.3
1979-83	0.8	0.4	0.7	1.1	1.5	1.6	1.6	2.6	2.4	3.2	3.5	5.9	6.8	9.3	9.5	14.2	13.3	19.8	2.7	2.6	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF CONNECTIVE AND OTHER SOFT TISSUE (ICD9 171)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	10	2	0	2	1	1	1	6	3	7	5	7	6	1	4	2	3	2	0	63
1965	7	2	2	1	0	5	2	3	4	4	4	2	5	3	5	8	6	1	0	64
1966	2	1	1	5	4	2	1	4	3	3	3	4	4	5	5	2	1	1	0	51
1967	7	2	2	4	4	2	2	2	4	2	11	1	11	1	5	5	2	0	0	67
1968	4	0	4	2	1	7	4	2	3	4	4	5	5	8	1	6	4	1	0	65
1964-68	30	7	9	14	10	17	10	17	17	20	27	19	31	18	20	23	16	5	0	310
1969	11	3	4	2	4	4	3	5	1	1	3	7	1	5	6	2	2	2	0	66
1970	4	1	4	2	1	1	4	3	4	3	5	7	3	9	8	4	5	4	0	72
1971	11	1	1	2	2	5	1	4	4	3	7	3	8	4	12	9	1	4	0	82
1972	6	2	4	2	3	4	2	4	2	11	3	8	10	9	6	7	4	2	0	89
1973	7	2	3	6	2	9	2	1	6	8	6	7	11	7	4	6	4	6	0	97
1969-73	39	9	16	14	12	23	12	17	17	26	24	32	33	34	36	28	16	18	0	406
1974	4	1	6	6	1	4	5	6	7	5	9	10	6	12	11	8	11	6	1	119
1975	11	3	1	7	8	2	2	3	4	10	4	7	7	13	7	7	6	7	0	109
1976	6	2	5	7	8	5	3	7	7	9	8	14	14	8	7	8	8	6	0	132
1977	3	3	0	0	4	7	4	5	6	11	9	7	4	8	16	12	7	6	0	112
1978	10	2	0	4	0	3	6	5	4	5	16	5	8	11	7	9	8	9	1	113
1974-78	34	11	12	24	21	21	20	26	28	40	46	43	39	52	48	44	40	34	2	585
1979	3	4	2	2	4	6	7	4	6	4	3	9	12	6	5	8	11	8	0	104
1980	2	0	2	1	6	1	5	6	5	6	4	10	13	4	13	4	7	2	0	91
1981	1	1	1	6	4	4	7	8	8	5	8	5	13	11	10	5	7	2	0	106
1982	4	1	3	2	5	2	9	7	3	2	7	8	9	14	2	5	7	8	2	100
1983	1	1	4	5	3	5	7	8	2	12	7	8	9	10	10	12	10	13	2	129
1979-83	11	7	12	16	22	18	35	33	24	29	29	40	56	45	40	34	42	33	4	530

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF CONNECTIVE AND OTHER SOFT TISSUE (ICD9 171)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	2.7	0.6	0.0	0.8	0.5	0.5	0.5	2.6	1.4	3.8	3.0	5.1	5.1	1.0	4.8	3.4	8.7	9.1	1.9	1.8	0.2
1965	1.9	0.5	0.6	0.4	0.0	2.4	0.9	1.3	1.7	2.1	2.4	1.4	4.1	2.9	5.8	13.2	16.7	4.3	1.9	1.7	0.1
1966	0.6	0.3	0.3	1.7	1.6	0.9	0.5	1.7	1.3	1.5	1.7	2.7	3.2	4.7	5.6	3.2	2.7	4.2	1.5	1.4	0.1
1967	2.0	0.5	0.6	1.3	1.5	0.9	0.9	0.9	1.7	1.0	6.2	0.7	8.6	0.9	5.5	7.7	5.1	0.0	1.9	1.8	0.2
1968	1.2	0.0	1.1	0.6	0.4	2.9	1.8	0.9	1.2	1.9	2.2	3.2	3.8	7.2	1.1	9.0	9.8	3.8	1.8	1.7	0.1
1964-68	1.7	0.4	0.5	0.9	0.8	1.5	0.9	1.5	1.5	2.0	3.1	2.6	5.0	3.4	4.6	7.4	8.5	4.2	1.8	1.7	0.2
1969	3.4	0.8	1.1	0.6	1.3	1.6	1.3	2.2	0.4	0.5	1.6	4.3	0.7	4.4	6.5	2.9	4.7	7.1	1.8	1.8	0.2
1970	1.3	0.3	1.1	0.6	0.3	0.4	1.7	1.3	1.7	1.3	2.7	4.2	2.1	7.6	8.4	5.7	11.3	13.2	1.9	1.7	0.2
1971	3.5	0.3	0.3	0.6	0.6	1.8	0.4	1.7	1.7	1.3	3.6	1.7	5.6	3.3	12.4	12.5	2.2	12.3	2.1	2.0	0.2
1972	1.9	0.5	1.0	0.6	0.9	1.3	0.8	1.7	0.9	4.6	1.5	4.6	6.7	7.2	6.1	9.4	8.4	5.8	2.2	2.1	0.2
1973	2.3	0.6	0.8	1.6	0.6	2.8	0.8	0.4	2.6	3.4	2.8	4.0	7.1	5.4	3.9	7.9	8.2	16.7	2.4	2.3	0.2
1969-73	2.5	0.5	0.8	0.8	0.7	1.6	1.0	1.5	1.5	2.2	2.4	3.7	4.6	5.6	7.4	7.7	7.0	11.2	2.1	2.0	0.2
1974	1.3	0.3	1.5	1.6	0.3	1.2	1.9	2.6	3.0	2.1	4.0	5.7	3.8	9.1	10.5	10.3	22.0	15.8	2.8	2.5	0.2
1975	3.6	0.9	0.3	1.8	2.2	0.6	0.7	1.3	1.7	4.3	1.7	3.9	4.2	9.6	6.5	8.7	11.6	17.8	2.6	2.5	0.2
1976	2.0	0.6	1.3	1.8	2.1	1.4	1.0	2.9	3.0	3.9	3.4	7.4	8.3	5.7	6.3	9.7	15.0	14.5	3.0	2.8	0.3
1977	1.0	0.9	0.0	0.0	1.0	2.0	1.3	2.0	2.6	4.7	3.8	3.5	2.3	5.5	13.9	14.2	12.6	13.9	2.4	2.2	0.2
1978	3.4	0.6	0.0	1.0	0.0	0.8	1.8	1.9	1.7	2.1	6.8	2.3	4.6	7.3	5.9	10.3	14.0	20.0	2.4	2.3	0.2
1974-78	2.3	0.7	0.6	1.2	1.1	1.2	1.3	2.1	2.4	3.4	3.9	4.5	4.7	7.4	8.6	10.7	14.9	16.4	2.6	2.5	0.2
1979	1.0	1.3	0.6	0.5	1.0	1.6	2.1	1.5	2.5	1.7	1.3	4.0	6.9	3.9	4.1	8.9	18.5	17.0	2.2	2.0	0.2
1980	0.7	0.0	0.6	0.2	1.5	0.3	1.4	2.1	2.1	2.6	1.7	4.3	7.3	2.5	10.3	4.3	11.3	4.1	1.9	1.7	0.2
1981	0.3	0.3	0.3	1.5	1.0	1.1	1.9	2.8	3.3	2.2	3.4	2.1	6.9	6.7	7.7	5.2	11.0	3.9	2.2	2.0	0.2
1982	1.4	0.3	0.9	0.5	1.2	0.5	2.5	2.2	1.2	0.9	3.0	3.4	4.5	8.4	1.5	5.0	10.7	14.8	1.9	1.8	0.2
1983	0.3	0.3	1.2	1.4	0.7	1.3	1.9	2.4	0.8	5.2	3.0	3.4	4.3	6.0	7.2	11.6	14.7	22.8	2.4	2.2	0.2
1979-83	0.8	0.5	0.7	0.8	1.1	1.0	2.0	2.2	1.9	2.5	2.5	3.5	5.9	5.5	6.1	7.0	13.2	12.8	2.1	1.9	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT MELANOMA OF SKIN (ICD9 172)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	1	2	1	6	12	19	11	19	9	9	9	8	9	5	1	1	0	122
1965	0	1	2	1	4	10	8	10	12	7	12	10	7	5	6	3	3	3	0	104
1966	0	0	0	4	6	6	10	16	9	12	11	14	10	6	8	6	2	0	0	120
1967	0	0	1	3	2	5	3	8	13	3	12	5	5	2	5	1	2	3	1	74
1968	0	0	2	3	7	3	6	14	14	15	9	10	6	8	3	2	8	2	2	114
1964-68	0	1	6	13	20	30	39	67	59	56	53	48	37	29	31	17	16	9	3	534
1969	0	0	0	1	7	13	12	9	14	9	8	12	9	5	9	0	8	1	1	118
1970	0	0	2	3	4	5	7	8	19	13	19	13	21	13	6	6	4	4	1	148
1971	0	0	1	5	5	9	3	13	18	15	15	19	14	15	7	3	4	2	0	148
1972	1	2	0	4	6	14	11	8	11	10	19	9	12	12	9	10	1	4	0	143
1973	1	1	2	3	10	6	16	14	15	15	17	22	18	9	9	5	10	2	1	176
1969-73	2	3	5	16	32	47	49	52	77	62	78	75	74	54	40	24	27	13	3	733
1974	1	1	2	4	10	11	15	12	15	17	26	20	13	16	6	11	11	9	0	200
1975	1	0	0	1	7	21	23	14	20	25	21	14	19	20	14	9	7	4	1	221
1976	2	0	0	4	5	9	18	16	20	34	27	19	18	27	19	9	7	3	2	239
1977	1	0	0	4	9	17	19	14	30	42	40	28	25	22	15	7	4	9	2	288
1978	1	0	0	5	10	13	13	21	23	24	43	23	33	24	21	18	12	8	4	296
1974-78	6	1	2	18	41	71	88	77	108	142	157	104	108	109	75	54	41	33	9	1244
1979	0	0	0	1	11	20	16	29	27	28	40	34	31	31	23	16	8	9	4	328
1980	0	0	0	3	6	20	31	29	29	27	32	33	35	30	33	19	7	3	1	338
1981	0	0	0	4	15	17	28	29	31	40	39	42	38	33	22	16	11	2	5	372
1982	0	0	1	3	11	15	35	22	26	30	47	41	46	35	32	16	17	14	1	392
1983	0	2	0	3	8	15	22	31	26	22	51	47	38	47	34	16	14	11	5	392
1979-83	0	2	1	14	51	87	132	140	139	147	209	197	188	176	144	83	57	39	16	1822

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT MELANOMA OF SKIN (ICD9 172)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.3	0.7	0.5	2.9	5.3	8.0	5.0	10.1	5.3	6.4	7.9	8.9	13.2	11.0	3.9	7.4	3.6	3.6	0.4
1965	0.0	0.3	0.6	0.3	1.8	4.8	3.6	4.2	5.3	3.7	6.9	7.0	6.0	5.5	8.6	6.5	11.6	21.6	3.0	3.1	0.3
1966	0.0	0.0	0.0	1.3	2.5	2.8	4.4	6.7	3.9	6.1	6.1	9.5	8.4	6.4	11.4	12.9	7.7	0.0	3.4	3.4	0.3
1967	0.0	0.0	0.3	0.9	0.8	2.2	1.3	3.3	5.5	1.5	6.6	3.3	4.1	2.1	7.1	2.1	7.6	20.7	2.0	2.0	0.2
1968	0.0	0.0	0.5	0.9	2.5	1.3	2.6	5.8	5.8	7.2	4.9	6.4	4.8	8.2	4.2	4.2	29.9	13.3	3.0	3.0	0.3
1964-68	0.0	0.1	0.3	0.9	1.6	2.7	3.4	5.6	5.1	5.7	6.0	6.5	6.2	6.2	8.9	7.3	12.3	12.7	3.0	3.0	0.3
1969	0.0	0.0	0.0	0.3	2.4	5.2	5.2	3.8	5.8	4.1	4.4	7.4	7.0	5.0	12.5	0.0	29.4	6.4	3.2	3.1	0.3
1970	0.0	0.0	0.5	0.9	1.3	1.9	3.0	3.4	7.9	5.7	10.3	7.8	15.9	12.6	8.2	12.4	14.4	24.4	3.9	3.9	0.4
1971	0.0	0.0	0.2	1.4	1.5	3.1	1.2	5.4	7.4	6.5	7.9	11.2	10.3	14.1	9.4	6.1	14.2	11.6	3.8	3.8	0.4
1972	0.3	0.5	0.0	1.1	1.8	4.6	4.4	3.4	4.5	4.3	9.6	5.3	8.5	11.1	11.7	20.2	3.5	22.3	3.6	3.6	0.4
1973	0.3	0.3	0.5	0.8	3.0	1.9	6.1	5.9	6.2	6.4	8.1	13.0	12.4	8.2	11.3	10.0	34.4	11.0	4.3	4.3	0.4
1969-73	0.1	0.2	0.2	0.9	2.0	3.3	4.0	4.4	6.4	5.4	8.1	8.9	10.8	10.2	10.7	9.8	19.1	15.2	3.8	3.7	0.4
1974	0.3	0.3	0.5	1.0	2.9	3.3	5.5	5.0	6.1	7.2	11.9	11.9	8.6	14.2	7.3	21.7	37.7	48.6	4.9	4.8	0.4
1975	0.3	0.0	0.0	0.2	1.9	6.0	8.0	5.7	8.2	10.5	9.4	8.1	12.2	17.2	16.6	17.2	24.0	21.1	5.2	5.2	0.5
1976	0.6	0.0	0.0	1.0	1.4	2.5	6.1	6.5	8.3	14.2	11.9	10.6	11.4	22.4	22.0	16.5	23.6	15.5	5.5	5.5	0.6
1977	0.3	0.0	0.0	1.0	2.4	4.8	6.0	5.5	12.5	17.5	17.6	14.8	15.8	17.5	17.0	12.4	13.2	45.7	6.5	6.6	0.7
1978	0.3	0.0	0.0	1.2	2.6	3.7	3.9	7.9	9.6	9.9	18.8	11.6	21.0	18.4	23.2	30.6	39.0	40.0	6.6	6.5	0.7
1974-78	0.4	0.1	0.1	0.9	2.2	4.1	5.9	6.2	9.0	11.9	13.9	11.5	13.9	18.0	17.4	19.8	27.5	34.2	5.7	5.7	0.6
1979	0.0	0.0	0.0	0.2	2.8	5.6	4.7	10.6	11.2	11.6	17.4	16.3	19.8	23.0	24.5	26.1	25.2	43.9	7.1	7.1	0.7
1980	0.0	0.0	0.0	0.7	1.5	5.6	8.9	10.3	11.9	11.3	13.7	15.5	21.7	21.6	34.0	30.2	21.1	14.5	7.2	7.2	0.8
1981	0.0	0.0	0.0	1.0	3.8	4.7	7.8	9.9	12.5	16.9	16.6	19.5	22.5	23.4	21.8	24.7	32.3	9.4	7.8	7.7	0.8
1982	0.0	0.0	0.3	0.7	2.7	4.1	9.9	7.0	10.2	12.7	19.8	18.8	25.9	24.5	30.6	24.1	47.6	64.8	8.1	8.0	0.8
1983	0.0	0.6	0.0	0.8	1.9	4.0	6.2	9.5	9.8	9.3	21.5	21.3	20.2	33.0	31.2	23.4	37.0	49.3	7.8	7.7	0.8
1979-83	0.0	0.1	0.1	0.7	2.6	4.8	7.5	9.4	11.1	12.4	17.8	18.3	22.1	25.2	28.5	25.6	33.0	36.7	7.6	7.5	0.8

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT MELANOMA OF SKIN (ICD9 172)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	4	6	3	8	12	16	25	12	16	10	13	13	9	4	0	6	0	157
1965	0	0	0	5	10	12	15	19	17	16	12	3	5	11	4	6	3	3	1	142
1966	0	0	1	7	3	10	19	10	16	9	27	9	10	7	7	7	5	5	0	152
1967	0	0	0	0	9	8	10	15	13	15	20	6	12	7	8	7	6	5	1	142
1968	0	0	2	2	11	9	13	11	17	15	12	14	10	5	7	7	5	4	1	145
1964-68	0	0	7	20	36	47	69	71	88	67	87	42	50	43	35	31	19	23	3	738
1969	0	0	1	3	9	14	7	17	29	25	15	19	10	19	13	9	4	8	1	203
1970	0	0	3	5	7	16	10	15	15	15	15	14	13	8	17	8	3	2	2	168
1971	0	0	3	2	11	9	12	11	21	19	20	8	10	14	11	7	5	8	0	171
1972	0	1	3	5	9	10	17	15	23	12	15	6	11	11	14	9	4	6	0	171
1973	0	1	1	5	19	20	22	17	22	27	22	14	16	14	17	13	8	4	3	245
1969-73	0	2	11	20	55	69	68	75	110	98	87	61	60	66	72	46	24	28	6	958
1974	0	0	0	10	8	23	14	23	14	27	29	22	12	13	14	12	7	13	2	243
1975	0	0	1	9	10	21	21	15	18	27	32	20	14	22	10	22	6	11	5	264
1976	1	0	0	9	13	22	24	25	22	32	31	28	31	18	9	12	15	13	3	308
1977	0	0	1	3	17	18	15	19	26	34	29	22	27	14	14	17	11	12	3	282
1978	0	0	3	4	16	28	36	24	38	39	41	41	22	23	30	15	13	14	8	395
1974-78	1	0	5	35	64	112	110	106	118	159	162	133	106	90	77	78	52	63	21	1492
1979	0	1	0	4	24	23	35	24	34	41	43	31	34	23	18	16	13	11	1	376
1980	0	0	0	8	15	42	37	29	32	45	30	39	37	26	23	15	14	14	4	410
1981	0	0	1	3	22	36	39	35	31	35	33	35	27	28	27	16	15	8	3	394
1982	0	0	1	4	25	33	47	32	35	38	40	35	39	25	29	16	14	10	7	430
1983	1	0	1	8	6	29	38	51	29	40	45	47	35	30	23	17	13	15	8	436
1979-83	1	1	3	27	92	163	196	171	161	199	191	187	172	132	120	80	69	58	23	2046

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT MELANOMA OF SKIN (ICD9 172)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	1.3	2.3	1.4	3.9	5.5	6.8	11.3	6.5	9.7	7.2	11.0	12.8	10.8	6.8	0.0	27.3	4.6	4.6	0.5
1965	0.0	0.0	0.0	1.8	4.4	5.7	6.9	8.2	7.4	8.5	7.1	2.1	4.1	10.6	4.7	9.9	8.4	13.0	4.1	4.0	0.4
1966	0.0	0.0	0.3	2.4	1.2	4.6	8.6	4.3	6.8	4.6	15.5	6.2	8.0	6.6	7.9	11.3	13.4	21.1	4.3	4.1	0.4
1967	0.0	0.0	0.0	0.0	3.4	3.5	4.5	6.5	5.4	7.4	11.2	4.0	9.3	6.5	8.9	10.8	15.4	19.9	3.9	3.7	0.4
1968	0.0	0.0	0.6	0.6	3.9	3.8	5.8	4.8	7.1	7.1	6.6	8.9	7.6	4.5	7.7	10.4	12.3	15.1	3.9	3.7	0.3
1964-68	0.0	0.0	0.4	1.4	2.9	4.3	6.3	6.1	7.5	6.8	10.0	5.7	8.0	8.1	8.0	9.9	10.1	19.1	4.2	4.0	0.4
1969	0.0	0.0	0.3	0.9	3.0	5.6	3.1	7.5	12.2	11.3	8.2	11.7	7.4	16.6	14.0	13.1	9.4	28.3	5.3	5.1	0.5
1970	0.0	0.0	0.8	1.5	2.2	6.0	4.3	6.6	6.4	6.5	8.0	8.3	9.3	6.8	18.0	11.3	6.8	6.6	4.3	4.1	0.4
1971	0.0	0.0	0.8	0.6	3.2	3.2	5.1	4.8	9.0	8.0	10.4	4.6	6.9	11.5	11.3	9.7	10.8	24.6	4.3	4.1	0.4
1972	0.0	0.3	0.8	1.4	2.7	3.3	7.0	6.5	9.9	5.1	7.4	3.4	7.4	8.8	14.1	12.1	8.4	17.5	4.2	4.0	0.4
1973	0.0	0.3	0.3	1.4	5.6	6.3	8.6	7.4	9.5	11.4	10.2	8.0	10.4	10.9	16.7	17.1	16.3	11.1	5.8	5.5	0.5
1969-73	0.0	0.1	0.6	1.1	3.4	4.9	5.7	6.6	9.4	8.4	8.9	7.1	8.3	10.9	14.8	12.7	10.5	17.4	4.8	4.6	0.4
1974	0.0	0.0	0.0	2.6	2.3	6.9	5.2	9.8	6.0	11.5	12.7	12.5	7.5	9.8	13.3	15.4	14.0	34.3	5.6	5.3	0.5
1975	0.0	0.0	0.3	2.3	2.7	6.0	7.5	6.3	7.7	11.5	13.7	11.0	8.4	16.3	9.3	27.5	11.6	27.9	5.9	5.5	0.5
1976	0.3	0.0	0.0	2.3	3.5	6.1	8.3	10.3	9.4	13.7	13.0	14.7	18.4	12.8	8.1	14.5	28.1	31.4	6.8	6.5	0.6
1977	0.0	0.0	0.3	0.8	4.4	5.0	4.8	7.6	11.1	14.6	12.3	10.9	15.8	9.6	12.2	20.1	19.9	27.7	6.1	5.8	0.5
1978	0.0	0.0	0.8	1.0	4.1	7.7	10.9	9.2	16.3	16.7	17.4	19.1	12.8	15.3	25.3	17.2	22.7	31.2	8.3	7.8	0.8
1974-78	0.1	0.0	0.3	1.8	3.4	6.3	7.4	8.6	10.1	13.6	13.8	13.8	12.7	12.8	13.8	18.9	19.4	30.4	6.6	6.2	0.6
1979	0.0	0.3	0.0	1.0	6.1	6.3	10.3	8.8	14.4	17.6	18.4	13.8	19.6	14.8	14.6	17.7	21.8	23.4	8.0	7.6	0.7
1980	0.0	0.0	0.0	2.0	3.8	11.2	10.5	10.4	13.3	19.4	12.8	16.9	20.7	16.1	18.2	16.1	22.7	28.5	8.3	8.0	0.8
1981	0.0	0.0	0.3	0.8	5.5	9.6	10.7	12.1	12.7	15.1	14.1	15.0	14.4	17.0	20.7	16.6	23.5	15.5	7.9	7.5	0.7
1982	0.0	0.0	0.3	1.0	6.2	8.6	13.0	10.3	13.9	16.4	17.1	15.0	19.7	15.0	21.6	16.0	21.4	18.5	8.4	8.0	0.8
1983	0.3	0.0	0.3	2.2	1.5	7.4	10.4	15.5	11.0	17.2	19.3	20.0	16.6	18.0	16.5	16.4	19.1	26.4	8.3	7.9	0.8
1979-83	0.1	0.1	0.2	1.4	4.6	8.6	11.0	11.5	13.0	17.1	16.3	16.2	18.1	16.2	18.4	16.5	21.6	22.4	8.2	7.8	0.8

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF FEMALE/MALE BREAST (ICD9 174/5)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	1	0	0	0	0	1	4	1	5	7	0	3	1	0	0	23
1965	0	0	0	0	0	1	0	0	1	1	3	5	3	5	3	4	2	0	0	28
1966	0	0	0	0	0	0	0	0	1	1	3	2	2	1	3	1	1	2	0	17
1967	0	0	0	0	0	0	0	0	0	1	0	3	3	4	3	3	0	2	0	19
1968	0	0	0	0	0	0	0	0	0	0	2	3	3	1	6	2	3	0	0	20
1964-68	0	0	0	0	1	1	0	0	2	4	12	14	16	18	15	13	7	4	0	107
1969	0	0	0	0	1	0	0	0	0	2	4	2	1	4	4	0	1	3	0	22
1970	0	0	0	0	1	1	0	1	0	2	2	1	4	3	3	1	0	2	0	21
1971	0	0	0	0	0	0	0	0	0	2	0	1	2	3	2	2	2	1	0	15
1972	0	0	0	0	0	1	0	0	3	1	2	3	6	3	3	5	1	1	0	29
1973	0	0	0	0	0	0	0	0	4	2	4	1	9	3	7	7	1	2	0	40
1969-73	0	0	0	0	2	2	0	1	7	9	12	8	22	16	19	15	5	9	0	127
1974	0	0	0	0	0	0	1	3	1	2	1	5	4	3	4	5	2	2	0	33
1975	0	0	0	1	0	0	1	0	0	3	2	4	5	2	4	5	2	0	0	29
1976	0	0	0	2	0	0	0	1	0	4	3	5	5	3	4	3	2	3	0	35
1977	0	0	0	0	0	1	0	0	0	2	1	1	9	2	6	5	1	2	1	31
1978	0	0	0	0	0	0	2	0	2	1	7	3	6	10	7	1	1	0	1	41
1974-78	0	0	0	3	0	1	4	4	3	12	14	18	29	20	25	19	8	7	2	169
1979	0	0	0	0	0	0	0	0	1	0	1	2	2	1	4	4	1	2	0	18
1980	0	0	0	0	0	0	0	0	0	2	3	1	2	5	5	5	3	2	0	28
1981	0	0	0	0	0	1	1	1	1	1	3	4	4	5	3	6	1	3	0	34
1982	0	0	0	0	0	0	0	0	1	1	2	3	4	5	2	7	1	1	0	27
1983	0	0	0	0	1	0	0	0	0	1	2	7	7	1	5	9	5	1	0	39
1979-83	0	0	0	0	1	1	1	1	3	5	11	17	19	17	19	31	11	9	0	146

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF FEMALE/MALE BREAST (ICD9 174/5)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.5	2.4	0.7	4.4	7.8	0.0	6.6	3.9	0.0	0.7	0.7	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.4	0.5	1.7	3.5	2.6	5.5	4.3	8.7	7.7	0.0	0.8	0.8	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	1.7	1.4	1.7	1.1	4.3	2.1	3.8	14.3	0.5	0.5	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	2.0	2.4	4.2	4.2	6.3	0.0	13.8	0.5	0.5	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	1.9	2.4	1.0	8.4	4.2	11.2	0.0	0.6	0.5	0.1
1964-68	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.2	0.4	1.4	1.9	2.7	3.9	4.3	5.6	5.4	5.6	0.6	0.6	0.1
1969	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.9	2.2	1.2	0.8	4.0	5.6	0.0	3.7	19.1	0.6	0.6	0.1
1970	0.0	0.0	0.0	0.0	0.3	0.4	0.0	0.4	0.0	0.9	1.1	0.6	3.0	2.9	4.1	2.1	0.0	12.2	0.6	0.6	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.6	1.5	2.8	2.7	4.1	7.1	5.8	0.4	0.4	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	1.2	0.4	1.0	1.8	4.3	2.8	3.9	10.1	3.5	5.6	0.7	0.7	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	0.9	1.9	0.6	6.2	2.7	8.8	14.1	3.4	11.0	1.0	1.0	0.1
1969-73	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.6	0.8	1.2	1.0	3.2	3.0	5.1	6.1	3.5	10.5	0.7	0.7	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.2	0.4	0.8	0.5	3.0	2.7	2.7	4.9	9.9	6.8	10.8	0.8	0.8	0.1
1975	0.0	0.0	0.0	0.2	0.0	0.0	0.3	0.0	0.0	1.3	0.9	2.3	3.2	1.7	4.7	9.5	6.8	0.0	0.7	0.7	0.1
1976	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.4	0.0	1.7	1.3	2.8	3.2	2.5	4.6	5.5	6.7	15.5	0.8	0.8	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.8	0.4	0.5	5.7	1.6	6.8	8.8	3.3	10.2	0.7	0.7	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.8	0.4	3.1	1.5	3.8	7.7	7.7	1.7	3.2	0.0	0.9	0.9	0.1
1974-78	0.0	0.0	0.0	0.1	0.0	0.1	0.3	0.3	0.2	1.0	1.2	2.0	3.7	3.3	5.8	7.0	5.4	7.2	0.8	0.8	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	1.0	1.3	0.7	4.3	6.5	3.1	9.8	0.4	0.4	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.3	0.5	1.2	3.6	5.2	7.9	9.1	9.7	0.6	0.6	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.3	0.4	0.4	1.3	1.9	2.4	3.5	3.0	9.3	2.9	14.2	0.7	0.7	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.8	1.4	2.2	3.5	1.9	10.5	2.8	4.6	0.5	0.5	0.1
1983	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.4	0.8	3.2	3.7	0.7	4.6	13.1	13.2	4.5	0.8	0.7	0.1
1979-83	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.4	0.9	1.6	2.2	2.4	3.8	9.6	6.4	8.5	0.6	0.6	0.1

SEE TEXT SECTION 'INTERPPETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF FEMALE/MALE BREAST (ICD9 174/5)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	2	6	8	58	120	267	291	303	260	254	250	234	191	129	72	0	2445
1965	0	0	0	1	4	18	70	134	225	270	266	241	231	189	243	185	126	84	0	2287
1966	0	0	0	0	5	21	54	117	264	296	249	303	257	237	202	176	129	104	0	2414
1967	0	0	0	0	4	24	48	131	225	333	277	269	269	238	241	190	106	81	0	2436
1968	0	0	0	1	2	21	67	136	257	331	284	262	260	239	218	167	114	83	0	2442
1964-68	0	0	0	4	21	92	297	638	1238	1521	1379	1335	1271	1153	1138	909	604	424	0	12024
1969	0	0	0	0	4	17	62	143	272	349	323	274	287	243	237	195	136	100	1	2643
1970	0	0	0	0	12	18	55	112	248	346	299	274	300	234	232	207	128	98	0	2563
1971	0	0	0	1	6	17	73	136	237	402	334	362	307	292	247	210	161	109	0	2894
1972	0	0	0	2	2	26	78	116	253	408	332	331	309	306	248	216	176	118	0	2922
1973	1	0	0	1	3	30	70	140	254	386	401	322	315	320	243	234	176	155	10	3060
1969-73	0	0	0	4	27	108	338	647	1264	1891	1689	1563	1518	1395	1207	1062	777	580	11	14082
1974	0	0	1	2	8	26	59	151	279	432	442	355	384	332	292	263	214	162	14	3416
1975	0	0	0	1	5	26	68	133	250	366	417	406	408	363	307	226	179	160	6	3321
1976	0	0	0	10	3	36	74	135	273	456	440	410	399	399	314	228	169	150	7	3503
1977	0	0	1	5	6	32	81	119	232	358	369	400	356	375	304	234	181	145	5	3203
1978	0	0	0	1	4	30	71	146	239	386	369	437	432	405	315	236	163	189	4	3427
1974-78	0	0	2	19	26	150	353	684	1273	1998	2037	2008	1979	1874	1532	1187	906	806	36	16870
1979	0	0	1	2	1	36	79	138	237	365	420	472	378	403	358	289	179	171	4	3533
1980	0	0	1	2	6	22	96	150	235	337	357	479	414	426	321	283	212	163	2	3506
1981	0	0	0	2	3	22	89	173	250	337	374	446	427	430	355	296	249	198	3	3654
1982	0	0	0	0	7	27	77	165	217	354	394	432	485	438	361	310	235	206	8	3716
1983	0	0	0	0	4	19	72	164	251	335	399	501	521	445	445	307	250	222	13	3948
1979-83	0	0	2	6	21	126	413	790	1190	1728	1944	2330	2225	2142	1840	1485	1125	960	30	18357

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF FEMALE/MALE BREAST (ICD9 174/5)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.8	2.8	3.9	26.5	51.3	120.5	157.2	184.2	188.4	215.3	246.1	281.9	325.9	372.8	327.3	73.0	67.1	7.1
1965	0.0	0.0	0.0	0.4	1.8	8.5	32.0	57.6	97.8	142.6	156.7	169.7	190.8	182.4	283.9	306.3	351.0	365.2	66.6	60.7	6.4
1966	0.0	0.0	0.0	0.0	2.0	9.6	24.5	50.2	111.7	151.4	142.8	207.3	206.4	223.4	228.0	284.3	346.8	438.8	68.6	62.9	6.6
1967	0.0	0.0	0.0	0.0	1.5	10.5	21.6	56.7	93.7	164.1	155.0	177.7	209.5	219.4	267.5	293.7	271.8	322.7	67.2	62.2	6.7
1968	0.0	0.0	0.0	0.3	0.7	8.8	30.0	59.4	107.1	156.1	156.6	166.8	196.8	214.7	239.0	249.3	280.1	313.2	65.6	61.0	6.5
1964-68	0.0	0.0	0.0	0.3	1.7	8.4	26.9	55.0	106.0	154.4	158.8	181.7	203.7	217.1	259.5	290.8	322.3	352.5	68.1	62.7	6.6
1969	0.0	0.0	0.0	0.0	1.3	6.8	27.5	62.8	114.7	157.2	176.1	168.3	211.5	212.0	255.9	283.4	320.8	353.4	69.2	64.1	6.7
1970	0.0	0.0	0.0	0.0	3.8	6.8	23.9	49.0	105.8	149.7	160.0	162.5	214.9	198.1	245.0	293.6	289.6	323.4	65.5	60.5	6.4
1971	0.0	0.0	0.0	0.3	1.8	6.1	30.9	59.3	102.1	169.4	173.7	209.0	213.3	239.9	254.4	291.3	348.5	335.4	72.0	66.6	7.0
1972	0.3	0.0	0.0	0.6	0.6	8.5	32.0	50.6	109.0	172.2	162.8	189.5	207.4	244.4	250.3	290.7	370.5	344.0	71.0	65.5	6.9
1973	0.0	0.0	0.0	0.3	0.9	9.4	27.4	60.9	109.4	162.9	185.7	184.0	204.5	248.8	238.7	307.9	359.2	431.8	72.3	66.6	6.9
1969-73	0.1	0.0	0.0	0.2	1.7	7.6	28.4	56.5	108.2	162.4	171.9	183.0	210.2	229.4	248.7	293.6	338.9	359.6	70.1	64.7	6.8
1974	0.0	0.0	0.3	0.5	2.3	7.8	22.0	64.4	119.4	183.4	194.1	201.5	240.8	250.8	278.4	337.6	427.1	427.4	78.8	72.4	7.5
1975	0.0	0.0	0.0	0.3	1.4	7.5	24.2	55.8	107.1	156.0	178.4	223.8	245.8	268.5	284.0	282.5	346.9	406.1	75.1	69.3	7.5
1976	0.0	0.0	0.0	2.5	0.8	10.0	25.4	55.6	116.8	195.7	185.0	215.3	236.4	284.2	283.1	275.7	317.1	362.3	77.4	72.4	7.7
1977	0.0	0.0	0.3	1.3	1.6	8.9	25.9	47.5	99.4	153.7	156.1	197.8	207.9	257.6	264.8	276.3	326.7	334.9	68.9	63.7	6.9
1978	0.0	0.0	0.0	0.2	1.0	8.3	21.6	55.8	102.5	165.4	156.6	203.7	250.7	269.6	265.4	270.0	285.0	420.9	72.0	67.1	7.2
1974-78	0.0	0.0	0.1	1.0	1.4	8.5	23.8	55.7	109.0	170.8	173.9	208.1	236.2	266.3	274.8	287.6	338.6	389.6	74.4	68.9	7.4
1979	0.0	0.0	0.3	0.5	0.3	9.8	23.2	50.8	100.3	156.5	179.7	209.7	218.1	260.2	291.3	320.4	300.3	363.1	72.5	67.0	7.2
1980	0.0	0.0	0.3	0.5	1.5	5.9	27.2	53.6	98.0	145.2	152.8	208.1	231.5	264.4	254.4	303.3	343.0	332.0	70.2	64.7	7.0
1981	0.0	0.0	0.0	0.5	0.8	5.9	24.4	59.7	102.2	145.6	160.2	191.3	227.5	261.6	272.4	307.4	389.7	384.5	71.4	65.5	7.0
1982	0.0	0.0	0.0	0.0	1.7	7.1	21.3	52.9	86.0	153.0	168.5	185.0	244.6	262.0	268.6	309.7	359.3	381.5	70.8	65.1	7.0
1983	0.0	0.0	0.0	0.0	1.0	4.9	19.7	49.9	95.5	144.2	170.8	213.5	247.7	266.5	318.8	296.0	367.1	390.2	73.1	67.1	7.4
1979-83	0.0	0.0	0.1	0.3	1.1	6.7	23.1	53.3	96.3	148.9	166.4	201.5	234.6	263.0	281.6	307.1	352.9	371.2	71.6	65.9	7.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF UTERUS, PART UNSPECIFIED (ICD9 179)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	1	0	0	0	1	1	3	6	5	7	6	9	6	9	1	6	4	0	65
1965	0	0	0	0	0	1	1	4	8	6	10	8	9	5	5	2	6	1	0	66
1966	0	0	0	0	0	0	2	1	5	6	2	4	4	7	5	2	1	3	0	42
1967	0	0	0	0	0	0	0	4	8	2	7	8	1	6	3	6	1	1	0	47
1968	0	0	0	0	0	0	0	3	4	2	10	10	5	10	8	2	4	4	0	62
1964-68	0	1	0	0	0	2	4	15	31	21	36	36	28	34	30	13	18	13	0	282
1969	0	0	0	0	1	0	0	1	4	4	4	2	5	2	2	6	1	2	0	35
1970	0	0	0	0	1	3	1	2	3	9	9	7	9	6	4	3	4	2	0	63
1971	0	0	0	0	1	6	1	2	5	8	12	7	7	10	3	7	2	3	0	74
1972	0	0	1	0	0	0	0	3	3	10	8	6	7	5	5	3	3	5	0	59
1973	0	0	0	0	1	1	2	2	3	6	8	12	5	10	3	5	3	3	1	65
1969-73	0	0	1	0	4	10	4	10	18	37	41	34	33	33	17	24	13	15	1	295
1974	0	0	0	0	2	1	3	3	1	7	6	12	11	3	7	4	2	0	1	63
1975	0	0	0	0	0	0	1	2	4	6	6	8	6	1	3	6	4	2	1	50
1976	0	0	0	0	0	1	3	4	13	3	7	5	6	10	4	7	2	3	0	68
1977	0	0	0	0	1	1	4	6	5	2	11	9	7	9	5	7	5	3	0	73
1978	0	0	0	0	0	0	0	0	7	1	4	3	5	1	1	5	4	6	0	37
1974-78	0	0	0	0	3	3	11	15	30	19	34	37	35	24	18	29	17	14	2	291
1979	0	0	0	0	1	1	0	1	4	5	2	4	9	6	1	7	4	2	0	47
1980	0	0	0	0	0	0	1	6	3	3	2	2	3	2	3	0	0	7	0	32
1981	0	0	0	0	0	1	4	3	5	4	2	4	1	4	5	1	2	5	1	42
1982	0	0	0	0	0	1	0	0	0	2	1	1	1	4	2	4	2	2	0	20
1983	0	0	0	0	0	0	4	0	5	2	7	4	4	1	1	4	2	4	0	38
1979-83	0	0	0	0	1	3	9	10	17	16	14	15	18	17	12	16	10	20	1	179

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF UTERUS, PART UNSPECIFIED (ICD9 179)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.3	0.0	0.0	0.0	0.5	0.5	1.3	2.7	2.7	4.3	4.3	7.6	5.9	10.8	1.7	17.3	18.2	2.0	1.8	0.2
1965	0.0	0.0	0.0	0.0	0.0	0.5	0.5	1.7	3.5	3.2	5.9	5.6	7.4	4.8	5.8	3.3	16.7	4.3	1.9	1.8	0.2
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	2.1	3.1	1.1	2.7	3.2	6.6	5.6	3.2	2.7	12.7	1.2	1.1	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	3.3	1.0	3.9	5.3	0.8	5.5	3.3	9.3	2.6	4.0	1.3	1.2	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	1.7	0.9	5.5	6.4	3.8	9.0	8.8	3.0	9.8	15.1	1.7	1.5	0.2
1964-68	0.0	0.1	0.0	0.0	0.0	0.2	0.4	1.3	2.7	2.1	4.1	4.9	4.5	6.4	6.8	4.2	9.6	10.8	1.6	1.5	0.2
1969	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	1.7	1.8	2.2	1.2	3.7	1.7	2.2	8.7	2.4	7.1	0.9	0.8	0.1
1970	0.0	0.0	0.0	0.0	0.3	1.1	0.4	0.9	1.3	3.9	4.8	4.2	6.4	5.1	4.2	4.3	9.0	6.6	1.6	1.5	0.2
1971	0.0	0.0	0.0	0.0	0.3	2.1	0.4	0.9	2.2	3.4	6.2	4.0	4.9	8.2	3.1	9.7	4.3	9.2	1.8	1.7	0.2
1972	0.0	0.0	0.3	0.0	0.0	0.0	0.0	1.3	1.3	4.2	3.9	3.4	4.7	4.0	5.0	4.0	6.3	14.6	1.4	1.3	0.1
1973	0.0	0.0	0.0	0.0	0.3	0.3	0.8	0.9	1.3	2.5	3.7	6.9	3.2	7.8	2.9	6.6	6.1	8.4	1.5	1.4	0.2
1969-73	0.0	0.0	0.1	0.0	0.2	0.7	0.3	0.9	1.5	3.2	4.2	4.0	4.6	5.4	3.5	6.6	5.7	9.3	1.5	1.4	0.1
1974	0.0	0.0	0.0	0.0	0.6	0.3	1.1	1.3	0.4	3.0	2.6	6.8	6.9	2.3	6.7	5.1	4.0	0.0	1.5	1.4	0.2
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.7	2.6	2.6	4.4	3.6	0.7	2.8	7.5	7.8	5.1	1.1	1.0	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.3	1.0	1.6	5.6	1.3	2.9	2.6	3.6	7.1	3.6	8.5	3.8	7.2	1.5	1.4	0.1
1977	0.0	0.0	0.0	0.0	0.3	0.3	1.3	2.4	2.1	0.9	4.7	4.5	4.1	6.2	2.6	8.3	9.0	6.9	1.6	1.4	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.0	0.4	1.7	1.4	2.9	0.7	0.8	5.7	7.0	13.4	0.8	0.7	0.1
1974-78	0.0	0.0	0.0	0.0	0.2	0.2	0.7	1.2	2.6	1.6	2.9	3.8	4.2	3.4	3.2	7.0	6.4	6.8	1.3	1.2	0.1
1979	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.4	1.7	2.1	0.9	1.8	5.2	3.9	0.8	7.8	6.7	4.2	1.0	0.9	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.3	2.1	1.3	1.3	0.9	0.9	1.7	1.2	2.4	0.0	0.0	14.3	0.6	0.6	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.3	1.1	1.0	2.0	1.7	0.9	1.7	0.5	2.4	3.8	1.0	3.1	9.7	0.8	0.7	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.9	0.4	0.4	0.5	2.4	1.5	4.0	3.1	3.7	0.4	0.3	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	1.9	0.9	3.0	1.7	1.9	0.6	0.7	3.9	2.9	7.0	0.7	0.6	0.1
1979-83	0.0	0.0	0.0	0.0	0.1	0.2	0.5	0.7	1.4	1.4	1.2	1.3	1.9	2.1	1.8	3.3	3.1	7.7	0.7	0.6	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF CERVIX UTERI (ICD9 180)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	3	29	46	107	117	103	88	75	69	46	47	36	16	10	1	793
1965	1	0	0	0	4	29	66	89	124	111	112	84	50	44	41	33	18	5	0	811
1966	0	0	0	0	12	28	52	103	143	92	87	61	75	43	43	24	11	12	0	786
1967	0	0	0	2	8	34	56	89	147	88	104	80	56	46	37	29	21	6	0	803
1968	0	0	0	0	10	41	73	90	108	116	74	81	63	51	36	29	12	10	0	794
1964-68	1	0	0	2	37	161	293	478	639	510	465	381	313	230	204	151	78	43	1	3987
1969	0	0	0	1	7	35	64	81	78	107	69	77	60	37	32	19	10	6	0	683
1970	0	0	0	0	6	28	50	42	74	87	85	77	74	45	33	28	16	13	0	658
1971	0	0	0	1	17	45	47	78	101	95	87	83	63	46	31	26	12	12	0	744
1972	0	0	1	0	11	29	59	59	93	91	77	57	71	49	41	25	10	7	0	680
1973	0	0	0	3	8	28	54	56	73	81	65	61	54	44	28	32	16	9	0	612
1969-73	0	0	1	5	49	165	274	316	419	461	383	355	322	221	165	130	64	47	0	3377
1974	0	0	0	0	13	42	42	49	55	74	73	72	53	58	43	23	15	15	4	631
1975	0	0	0	2	7	28	39	52	68	64	62	75	55	46	41	29	15	17	4	604
1976	0	0	0	1	21	43	62	51	53	59	66	76	45	48	35	21	13	13	3	610
1977	0	0	0	1	11	27	54	60	39	57	56	75	61	40	24	27	24	10	0	566
1978	0	0	0	1	7	34	58	51	56	39	60	58	40	55	34	25	11	15	1	545
1974-78	0	0	0	5	59	174	255	263	271	293	317	356	254	247	177	125	78	70	12	2956
1979	0	0	0	0	11	37	48	54	50	41	59	68	54	42	25	21	27	21	1	559
1980	0	0	0	1	11	37	60	42	54	50	47	67	44	38	32	30	16	12	1	542
1981	0	0	0	1	12	43	58	65	62	48	55	42	48	51	34	17	16	15	0	567
1982	0	0	0	3	9	44	61	53	52	46	47	52	56	46	36	21	9	12	1	548
1983	0	0	0	1	16	41	72	72	43	56	34	43	43	59	29	27	14	14	1	565
1979-83	0	0	0	6	59	202	299	286	261	241	242	272	245	236	156	116	82	74	4	2781

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF CERVIX UTERI (ICD9 180)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	1.4	14.0	21.1	45.7	52.8	55.6	53.5	54.3	58.5	45.3	56.6	61.4	46.2	45.5	23.2	22.5	2.3
1965	0.3	0.0	0.0	0.0	1.8	13.8	30.2	38.2	53.9	58.6	66.0	59.2	41.3	42.5	47.9	54.6	50.1	21.7	23.3	22.6	2.2
1966	0.0	0.0	0.0	0.0	4.9	12.8	23.6	44.2	60.5	47.1	49.9	41.7	60.2	40.5	48.5	38.8	29.6	50.6	21.8	21.5	2.1
1967	0.0	0.0	0.0	0.6	3.0	14.9	25.2	38.5	61.2	43.4	58.2	52.8	43.6	42.4	41.1	44.8	53.8	23.9	21.9	21.3	2.1
1968	0.0	0.0	0.0	0.0	3.6	17.3	32.6	39.3	45.0	54.7	40.8	51.6	47.7	45.8	39.5	43.3	29.5	37.7	21.3	20.9	2.1
1964-68	0.1	0.0	0.0	0.1	3.0	14.6	26.5	41.2	54.7	51.8	53.5	51.9	50.2	43.3	46.5	48.3	41.6	35.7	22.3	21.7	2.2
1969	0.0	0.0	0.0	0.3	2.4	14.1	28.4	35.6	32.9	48.2	37.6	47.3	44.2	32.3	34.6	27.6	23.6	21.2	18.0	17.7	1.8
1970	0.0	0.0	0.0	0.0	1.9	10.6	21.7	18.4	31.6	37.6	45.5	45.7	53.0	38.1	34.8	39.7	36.2	42.9	16.9	16.4	1.7
1971	0.0	0.0	0.0	0.3	5.0	16.0	19.9	34.0	43.5	40.0	45.2	47.9	43.8	37.8	31.9	36.1	26.0	36.9	18.7	18.3	1.8
1972	0.0	0.0	0.3	0.0	3.3	9.5	24.2	25.7	40.1	38.4	37.8	32.6	47.7	39.1	41.4	33.6	21.1	20.4	16.7	16.4	1.7
1973	0.0	0.0	0.0	0.8	2.3	8.8	21.2	24.4	31.5	34.2	30.1	34.9	35.1	34.2	27.5	42.1	32.7	25.1	14.7	14.2	1.4
1969-73	0.0	0.0	0.1	0.3	3.0	11.6	23.0	27.6	35.9	39.6	39.0	41.6	44.6	36.3	34.0	35.9	27.9	29.1	16.9	16.6	1.7
1974	0.0	0.0	0.0	0.0	3.7	12.5	15.6	20.9	23.5	31.4	32.1	40.9	33.2	43.8	41.0	29.5	29.9	39.6	14.6	14.1	1.5
1975	0.0	0.0	0.0	0.5	1.9	8.0	13.9	21.8	29.1	27.3	26.5	41.3	33.1	34.0	37.9	36.2	29.1	43.1	13.7	13.2	1.4
1976	0.0	0.0	0.0	0.3	5.6	12.0	21.3	21.0	22.7	25.3	27.7	39.9	26.7	34.2	31.6	25.4	24.4	31.4	13.5	13.1	1.3
1977	0.0	0.0	0.0	0.3	2.9	7.5	17.3	24.0	16.7	24.5	23.7	37.1	35.6	27.5	20.9	31.9	43.3	23.1	12.4	11.8	1.2
1978	0.0	0.0	0.0	0.2	1.8	9.4	17.6	19.5	24.0	16.7	25.5	27.0	23.2	36.6	28.6	28.6	19.2	33.4	11.5	11.1	1.1
1974-78	0.0	0.0	0.0	0.3	3.2	9.9	17.2	21.4	23.2	25.1	27.1	36.9	30.3	35.1	31.8	30.3	29.1	33.8	13.1	12.6	1.3
1979	0.0	0.0	0.0	0.0	2.8	10.1	14.1	19.9	21.2	17.6	25.2	30.2	31.2	27.1	20.3	23.3	45.3	44.6	11.6	11.0	1.1
1980	0.0	0.0	0.0	0.2	2.8	9.9	17.0	15.0	22.5	21.5	20.1	29.1	24.6	23.6	25.4	32.2	25.9	24.4	11.0	10.5	1.1
1981	0.0	0.0	0.0	0.3	3.0	11.4	15.9	22.4	25.3	20.7	23.6	18.0	25.6	31.0	26.1	17.7	25.0	29.1	11.4	11.1	1.1
1982	0.0	0.0	0.0	0.8	2.2	11.5	16.8	17.0	20.6	19.9	20.1	22.3	28.2	27.5	26.8	21.0	13.8	22.2	10.7	10.4	1.1
1983	0.0	0.0	0.0	0.3	3.9	10.5	19.7	21.9	16.4	24.1	14.6	18.3	20.4	35.3	20.8	26.0	20.6	24.6	10.7	10.3	1.0
1979-83	0.0	0.0	0.0	0.3	3.0	10.7	16.7	19.3	21.1	20.8	20.7	23.5	25.8	29.0	23.9	24.0	25.7	28.6	11.1	10.7	1.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF PLACENTA (ICD9 181)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	1	4	0	0	0	0	0	0	0	0	1	0	0	0	0	0	6
1965	0	0	0	0	2	0	0	0	1	0	0	0	0	0	0	0	0	0	0	3
1966	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	3
1967	0	0	0	1	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	6
1968	0	0	0	0	2	2	1	1	0	0	0	0	0	0	0	0	0	0	0	5
1964-68	0	0	0	0	3	2	0	0	0	0	0	0	0	0	0	0	0	0	0	23
1969	0	0	0	2	12	4	1	1	2	0	0	0	0	1	0	0	0	0	0	4
1970	0	0	0	0	2	0	0	0	0	1	0	0	0	1	0	0	0	0	0	4
1971	0	0	0	1	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4
1972	0	0	0	1	1	2	0	0	0	0	0	0	0	0	0	0	0	0	0	7
1973	0	0	0	1	2	3	0	0	0	1	0	0	0	0	0	0	0	0	0	5
1969-73	0	0	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	24
1974	0	0	0	4	9	6	0	0	0	3	0	0	0	1	0	0	0	0	0	11
1975	0	0	0	1	2	3	2	1	1	0	0	1	0	0	0	0	0	0	1	17
1976	0	0	0	3	4	3	2	2	1	0	1	0	0	0	0	0	0	0	0	17
1977	0	0	0	1	5	7	1	0	1	2	0	0	0	0	0	0	0	0	0	14
1978	0	0	0	1	2	5	5	0	1	0	0	0	0	0	0	0	0	0	0	15
1974-78	0	0	0	2	3	4	3	2	1	0	0	0	0	0	0	0	0	0	1	74
1979	0	0	0	8	16	22	13	5	5	2	1	1	0	0	0	0	0	0	0	10
1980	0	0	0	1	3	0	3	2	1	0	0	0	0	0	0	0	0	0	0	4
1981	0	0	0	2	0	0	2	0	0	0	0	0	0	0	0	0	1	0	0	5
1982	0	0	0	1	0	0	2	0	1	0	0	0	0	0	0	0	1	0	0	11
1983	0	0	0	1	4	2	1	1	1	0	0	0	0	0	0	0	0	0	0	3
1979-83	0	0	0	6	7	2	8	4	3	1	0	0	0	0	0	0	2	0	0	33

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF PLACENTA (ICD9 181)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.4	1.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1965	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1966	0.0	0.0	0.0	0.3	0.4	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1967	0.0	0.0	0.0	0.0	0.8	0.9	0.4	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1968	0.0	0.0	0.0	0.0	1.1	0.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.0
1964-68	0.0	0.0	0.0	0.1	1.0	0.4	0.1	0.1	0.2	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1969	0.0	0.0	0.0	0.0	0.7	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1970	0.0	0.0	0.0	0.3	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1971	0.0	0.0	0.0	0.3	0.3	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1972	0.0	0.0	0.0	0.3	0.6	1.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1973	0.0	0.0	0.3	0.3	0.3	0.3	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1969-73	0.0	0.0	0.1	0.2	0.6	0.4	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1974	0.0	0.0	0.0	0.3	0.6	0.9	0.7	0.4	0.4	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1975	0.0	0.0	0.0	0.8	1.1	0.9	0.7	0.8	0.4	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0
1976	0.0	0.0	0.0	0.3	1.3	1.9	0.3	0.0	0.4	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0
1977	0.0	0.0	0.0	0.3	0.5	1.4	1.6	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1978	0.0	0.0	0.0	0.5	0.8	1.1	0.9	0.8	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1974-78	0.0	0.0	0.0	0.4	0.9	1.2	0.9	0.4	0.4	0.2	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1979	0.0	0.0	0.0	0.2	0.8	0.0	0.9	0.7	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1980	0.0	0.0	0.0	0.5	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1981	0.0	0.0	0.0	0.3	0.0	0.0	0.5	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	0.0	0.1	0.1	0.0
1982	0.0	0.0	0.0	0.3	1.0	0.5	0.3	0.3	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.5	0.0	0.3	0.2	0.0
1983	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.3	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1979-83	0.0	0.0	0.0	0.3	0.4	0.1	0.4	0.3	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.1	0.1	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF BODY OF UTERUS (ICD9 182)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	1	1	0	2	7	28	38	75	83	75	76	58	34	15	10	0	503
1965	0	0	0	1	0	1	3	5	16	43	76	68	81	85	49	30	17	7	0	482
1966	0	0	0	1	2	2	1	12	21	55	67	71	53	73	51	41	28	7	0	485
1967	0	0	0	0	1	2	4	11	23	52	68	80	69	78	54	34	18	16	0	510
1968	0	0	0	0	1	2	2	16	23	43	74	84	59	71	47	31	15	15	0	483
1964-68	0	0	0	3	5	7	12	51	111	231	360	386	337	383	259	170	93	55	0	2463
1969	0	0	0	1	0	1	5	12	18	46	83	90	85	69	62	41	29	10	0	552
1970	0	0	0	1	4	1	2	9	24	50	78	112	87	80	72	28	23	16	0	587
1971	0	0	0	1	0	2	3	11	16	41	83	108	84	80	62	37	15	10	0	553
1972	0	0	0	0	2	2	6	11	24	39	96	108	121	94	56	36	25	24	0	644
1973	0	0	1	0	0	6	10	14	32	60	128	149	137	93	74	68	32	17	2	823
1969-73	0	0	1	3	6	12	26	57	114	236	468	567	514	416	326	210	124	77	2	3159
1974	0	0	1	2	0	2	2	5	23	59	124	158	148	106	73	49	27	16	2	797
1975	0	0	0	0	0	3	8	15	28	55	146	191	147	90	87	50	32	11	2	865
1976	0	0	0	0	2	5	8	17	20	61	143	189	183	120	87	64	30	17	1	947
1977	0	0	0	0	1	7	10	8	21	48	121	149	191	135	94	53	32	24	2	896
1978	0	0	0	1	1	0	11	16	20	65	123	184	177	123	81	56	46	23	0	927
1974-78	0	0	1	3	4	17	39	61	112	288	657	871	846	574	422	272	167	91	7	4432
1979	0	0	0	0	0	3	9	17	26	51	111	191	195	151	99	62	37	26	1	979
1980	0	0	0	1	0	2	12	12	33	52	101	173	192	147	107	69	33	27	1	962
1981	0	0	0	1	0	2	10	14	23	50	117	169	167	130	127	90	41	24	2	967
1982	0	0	0	1	0	5	11	9	22	41	113	140	192	176	117	71	45	31	4	978
1983	0	0	0	0	1	6	7	24	23	54	90	153	190	177	135	67	44	23	2	996
1979-83	0	0	0	3	1	18	49	76	127	248	532	826	936	781	585	359	200	131	10	4882

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF BODY OF UTERUS (ICD9 182)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.4	0.5	0.0	0.9	3.0	12.6	20.5	45.6	60.1	63.6	74.8	69.9	58.0	43.4	45.5	15.2	14.2	1.7
1965	0.0	0.0	0.0	0.4	0.0	0.5	1.4	2.1	7.0	22.7	44.8	47.9	66.9	82.0	57.2	49.7	47.4	30.4	14.3	13.4	1.7
1966	0.0	0.0	0.0	0.3	0.8	0.9	0.5	5.2	8.9	28.1	38.4	48.6	42.6	68.8	57.6	66.2	75.3	29.5	13.9	12.7	1.5
1967	0.0	0.0	0.0	0.0	0.4	0.9	1.8	4.8	9.6	25.6	38.1	52.8	53.7	71.9	59.9	52.6	46.2	63.7	14.2	13.2	1.6
1968	0.0	0.0	0.0	0.0	0.4	0.8	0.9	7.0	9.6	20.3	40.8	53.5	44.7	63.8	51.5	46.3	36.9	56.6	13.1	12.2	1.5
1964-68	0.0	0.0	0.0	0.2	0.4	0.6	1.1	4.4	9.5	23.5	41.4	52.5	54.0	72.1	59.1	54.4	49.6	45.7	14.1	13.1	1.6
1969	0.0	0.0	0.0	0.3	0.0	0.4	2.2	5.3	7.6	20.7	45.3	55.3	62.6	60.2	67.0	59.6	68.4	35.3	14.6	13.4	1.6
1970	0.0	0.0	0.0	0.3	1.3	0.4	0.9	3.9	10.2	21.6	41.7	66.4	62.3	67.7	76.0	39.7	52.0	52.8	15.1	14.1	1.7
1971	0.0	0.0	0.0	0.3	0.0	0.7	1.3	4.8	6.9	17.3	43.2	62.4	58.4	65.7	63.9	51.3	32.5	30.8	13.8	13.0	1.6
1972	0.0	0.0	0.0	0.0	0.6	0.7	2.5	4.8	10.3	16.5	47.1	61.8	81.2	75.1	56.5	48.5	52.6	70.0	15.7	14.7	1.8
1973	0.0	0.0	0.3	0.0	0.0	1.9	3.9	6.1	13.8	25.3	59.3	85.1	89.0	72.3	72.7	89.5	65.3	47.4	19.6	18.1	2.1
1969-73	0.0	0.0	0.1	0.2	0.4	0.8	2.2	5.0	9.8	20.3	47.6	66.4	71.2	68.4	67.2	58.1	54.1	47.7	15.8	14.7	1.8
1974	0.0	0.0	0.3	0.5	0.0	0.6	0.7	2.1	9.8	25.0	54.5	89.7	92.8	80.1	69.6	62.9	53.9	42.2	18.6	17.3	2.1
1975	0.0	0.0	0.0	0.0	0.0	0.9	2.9	6.3	12.0	23.4	62.4	105.3	88.6	66.6	80.5	62.5	62.0	27.9	19.7	18.3	2.2
1976	0.0	0.0	0.0	0.0	0.5	1.4	2.8	7.0	8.6	26.2	60.1	99.3	108.4	85.5	78.4	77.4	56.3	41.1	21.0	19.5	2.4
1977	0.0	0.0	0.0	0.0	0.3	2.0	3.2	3.2	9.0	20.6	51.2	73.7	111.6	92.7	81.9	62.6	57.8	55.4	19.2	17.9	2.2
1978	0.0	0.0	0.0	0.2	0.3	0.0	3.3	6.1	8.6	27.8	52.2	85.8	102.7	81.9	68.2	64.1	80.4	51.2	19.5	18.1	2.2
1974-78	0.0	0.0	0.1	0.2	0.2	1.0	2.6	5.0	9.6	24.6	56.1	90.3	101.0	81.6	75.7	65.9	62.4	44.0	19.6	18.2	2.2
1979	0.0	0.0	0.0	0.0	0.0	0.8	2.6	6.3	11.0	21.9	47.5	84.9	112.5	97.5	80.6	68.7	62.1	55.2	19.9	18.7	2.3
1980	0.0	0.0	0.0	0.2	0.0	0.5	3.4	4.3	13.8	22.4	43.2	75.2	107.4	91.2	84.8	74.0	53.4	55.0	19.1	17.9	2.2
1981	0.0	0.0	0.0	0.3	0.0	0.5	2.7	4.8	9.4	21.6	50.1	72.5	89.0	79.1	97.5	93.5	64.2	46.6	18.7	17.2	2.1
1982	0.0	0.0	0.0	0.3	0.0	1.3	3.0	2.9	8.7	17.7	48.3	60.0	96.8	105.3	87.1	70.9	68.8	57.4	18.4	17.0	2.1
1983	0.0	0.0	0.0	0.0	0.2	1.5	1.9	7.3	8.8	23.2	38.5	65.2	90.3	106.0	96.7	64.6	64.6	40.4	18.3	17.0	2.2
1979-83	0.0	0.0	0.0	0.2	0.1	1.0	2.7	5.1	10.3	21.4	45.5	71.4	98.7	95.9	89.5	74.2	62.7	50.7	18.9	17.5	2.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OVARY AND OTHER UTERINE ADNEXA (ICD9 183)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	1	1	3	1	5	7	24	37	61	58	71	72	51	52	31	17	16	0	508
1965	0	0	3	2	3	2	11	19	28	44	54	57	58	51	44	33	20	7	0	436
1966	0	0	2	1	5	8	18	21	36	46	60	55	64	41	60	30	11	10	0	468
1967	0	0	1	6	4	9	9	22	36	60	70	45	47	53	45	39	16	10	0	472
1968	0	0	2	2	5	4	8	13	27	47	55	54	43	56	46	25	10	8	0	405
1964-68	0	1	9	14	18	28	53	99	164	258	297	282	284	252	247	158	74	51	0	2289
1969	0	2	0	2	4	4	11	17	47	55	59	69	68	55	47	35	14	11	0	500
1970	0	1	2	5	7	8	13	13	41	48	55	67	65	55	43	35	20	12	0	490
1971	0	0	0	3	8	8	12	12	35	52	68	67	55	52	47	31	19	7	0	476
1972	0	1	6	8	4	11	12	23	33	71	71	69	50	60	52	31	30	10	0	542
1973	0	1	2	5	7	9	13	18	29	62	66	75	94	60	35	40	25	23	1	565
1969-73	0	5	10	23	30	40	61	83	185	288	319	347	332	282	224	172	108	63	1	2573
1974	0	1	4	2	8	10	17	20	41	59	72	70	62	63	55	42	23	19	2	570
1975	0	0	2	5	16	11	9	24	36	51	69	65	66	80	58	49	15	17	1	574
1976	0	0	2	3	8	12	13	15	38	58	79	77	87	75	57	58	26	15	2	625
1977	1	1	0	6	13	14	9	20	36	56	78	74	89	75	61	43	24	22	0	622
1978	0	0	5	7	11	15	25	25	27	60	67	90	75	68	69	47	35	27	0	653
1974-78	1	2	13	23	56	62	73	104	178	284	365	376	379	361	300	239	123	100	5	3044
1979	1	2	0	7	10	13	15	24	29	59	74	96	90	71	59	39	30	27	0	646
1980	0	2	3	7	10	13	18	24	34	51	75	82	93	73	62	48	31	28	1	655
1981	1	1	2	5	16	13	13	24	28	52	64	84	91	100	82	50	28	23	2	679
1982	1	0	4	7	11	6	15	23	27	54	64	79	81	80	70	45	45	20	1	633
1983	0	0	1	4	12	16	17	25	33	52	62	80	83	67	79	49	43	24	1	648
1979-83	3	5	10	30	59	61	78	120	151	268	339	421	438	391	352	231	177	122	5	3261

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OVARY AND OTHER UTERINE ADNEXA (ICD9 183)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.3	0.3	1.1	0.5	2.4	3.2	10.3	16.7	33.0	35.3	51.4	61.0	50.2	62.7	52.9	49.1	72.7	15.3	14.3	1.6
1965	0.0	0.0	0.9	0.7	1.3	0.9	5.0	8.2	12.2	23.2	31.8	40.1	47.9	49.2	51.4	54.6	55.7	30.4	12.8	11.8	1.4
1966	0.0	0.0	0.6	0.3	2.0	3.7	8.2	9.0	15.2	23.5	34.4	37.6	51.4	38.6	67.7	48.5	29.6	42.2	13.3	12.5	1.5
1967	0.0	0.0	0.3	1.9	1.5	4.0	4.0	9.5	15.0	29.6	39.2	29.7	36.6	48.8	49.9	60.3	41.0	39.6	13.0	12.2	1.3
1968	0.0	0.0	0.6	0.6	1.8	1.7	3.6	5.7	11.2	22.2	30.3	34.4	32.6	50.3	50.4	37.3	24.6	30.2	10.9	10.3	1.2
1964-68	0.0	0.1	0.5	0.9	1.5	2.5	4.8	8.5	14.0	26.2	34.2	38.4	45.5	47.4	56.3	50.5	39.5	42.4	13.0	12.2	1.4
1969	0.0	0.5	0.0	0.6	1.3	1.6	4.9	7.5	19.8	24.8	32.2	42.4	50.1	48.0	50.8	50.9	33.0	38.9	13.2	12.4	1.4
1970	0.0	0.3	0.5	1.5	2.2	3.0	5.6	5.7	17.5	20.8	29.4	39.7	46.6	46.6	45.4	49.6	45.2	39.6	12.6	11.7	1.3
1971	0.0	0.0	0.0	0.9	2.4	2.9	5.1	5.2	15.1	21.9	35.4	38.7	38.2	42.7	48.4	43.0	41.1	21.5	11.9	11.2	1.3
1972	0.0	0.3	1.5	2.2	1.2	3.6	4.9	10.0	14.2	30.0	34.8	39.5	33.6	47.9	52.5	41.7	63.2	29.2	13.2	12.3	1.4
1973	0.0	0.3	0.5	1.4	2.0	2.8	5.1	7.8	12.5	26.2	30.6	42.9	61.0	46.7	34.4	52.6	51.0	64.1	13.5	12.6	1.4
1969-73	0.0	0.3	0.5	1.3	1.8	2.8	5.1	7.3	15.8	24.7	32.5	40.6	46.0	46.4	46.2	47.6	47.1	39.1	12.9	12.0	1.3
1974	0.0	0.3	1.0	0.5	2.3	3.0	6.3	8.5	17.5	25.0	31.6	39.7	38.9	47.6	52.4	53.9	45.9	50.1	13.2	12.3	1.4
1975	0.0	0.0	0.5	1.3	4.4	3.2	3.2	10.1	15.4	21.7	29.5	35.8	39.8	59.2	53.7	61.2	29.1	43.1	13.1	12.1	1.4
1976	0.0	0.0	0.5	0.8	2.1	3.3	4.5	6.2	16.3	24.9	33.2	40.4	51.5	53.4	51.4	70.1	48.8	36.2	13.8	12.8	1.4
1977	0.3	0.3	0.0	1.5	3.4	3.9	2.9	8.0	15.4	24.0	33.0	36.6	52.0	51.5	53.1	50.8	43.3	50.8	13.5	12.6	1.4
1978	0.0	0.0	1.4	1.7	2.8	4.1	7.6	9.6	11.6	25.7	28.4	42.0	43.5	45.3	58.1	53.8	61.2	60.1	13.8	12.6	1.4
1974-78	0.1	0.1	0.7	1.2	3.0	3.5	4.9	8.5	15.2	24.3	31.2	39.0	45.2	51.3	53.8	57.9	46.0	48.3	13.5	12.5	1.4
1979	0.3	0.6	0.0	1.7	2.6	3.5	4.4	8.8	12.3	25.3	31.7	42.6	51.9	45.8	48.0	43.2	50.3	57.3	13.4	12.5	1.4
1980	0.0	0.7	0.9	1.7	2.5	3.5	5.1	8.6	14.2	22.0	32.1	35.6	52.0	45.3	49.1	51.4	50.2	57.0	13.2	12.3	1.4
1981	0.3	0.3	0.6	1.3	4.0	3.5	3.6	8.3	11.4	22.5	27.4	36.0	48.5	60.8	62.9	51.9	43.8	44.7	13.4	12.4	1.4
1982	0.3	0.0	1.2	1.8	2.7	1.6	4.1	7.4	10.7	23.3	27.4	33.8	40.8	47.8	52.1	45.0	68.8	37.0	12.3	11.2	1.3
1983	0.0	0.0	0.3	1.1	2.9	4.1	4.7	7.6	12.6	22.4	26.5	34.1	39.5	40.1	56.6	47.3	63.1	42.2	12.2	11.1	1.3
1979-83	0.2	0.3	0.6	1.5	3.0	3.2	4.4	8.1	12.2	23.1	29.0	36.4	46.2	48.0	53.9	47.8	55.5	47.2	12.9	11.9	1.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED FEMALE GENITAL ORGANS (ICD9 184)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	1	0	1	0	2	3	7	6	6	5	11	8	15	19	8	7	0	99
1965	0	0	0	0	0	0	1	0	3	3	4	6	14	11	7	12	7	9	0	77
1966	0	0	0	0	2	1	1	2	3	6	7	6	9	10	19	22	11	7	0	106
1967	1	0	0	1	0	0	1	2	3	3	8	4	7	5	13	14	15	8	0	85
1968	1	0	0	0	0	1	0	4	3	3	6	17	6	9	15	10	9	9	0	93
1964-68	2	0	1	1	3	2	5	11	19	21	31	38	47	43	69	77	50	40	0	460
1969	0	0	0	0	0	0	2	1	5	8	4	14	15	12	18	17	11	15	0	122
1970	0	0	0	0	0	2	0	5	2	5	5	8	13	15	20	8	6	10	0	99
1971	0	0	0	1	0	0	1	3	3	8	6	13	12	9	11	13	8	10	0	98
1972	0	0	0	1	0	0	1	1	3	3	9	10	12	10	17	13	9	14	0	103
1973	1	0	0	0	1	2	4	3	6	4	11	14	11	13	21	22	14	11	2	140
1969-73	1	0	0	2	1	4	8	13	19	28	35	59	63	59	87	73	48	60	2	562
1974	0	1	0	0	2	4	1	2	6	7	10	9	18	27	18	15	19	23	0	162
1975	0	0	0	0	5	0	6	6	5	4	10	11	14	11	10	19	14	7	2	124
1976	0	0	0	2	2	0	1	5	0	10	7	8	17	15	14	18	16	8	0	123
1977	2	0	0	0	5	1	4	2	3	8	8	7	20	13	21	18	22	17	0	151
1978	0	0	0	0	0	3	3	9	3	5	13	8	19	13	17	16	22	17	0	148
1974-78	2	1	0	2	14	8	15	24	17	34	48	43	88	79	80	86	93	72	2	708
1979	0	0	0	1	0	6	5	3	4	9	3	14	13	11	20	14	16	14	0	133
1980	0	0	0	0	2	0	4	4	1	2	8	10	11	16	19	15	17	11	1	121
1981	0	0	0	0	1	1	4	8	4	7	11	10	11	15	18	18	21	12	0	141
1982	0	0	0	0	0	1	2	6	2	5	6	13	14	11	11	16	19	18	2	126
1983	1	0	0	0	1	0	0	5	2	6	14	11	12	22	11	25	18	22	0	150
1979-83	1	0	0	1	4	8	15	26	13	29	42	58	61	75	79	88	91	77	3	671

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED FEMALE GENITAL ORGANS (ICD9 184)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.3	0.0	0.5	0.0	0.9	1.3	3.2	3.2	3.6	3.6	9.3	7.9	18.1	32.4	23.1	31.8	3.0	2.5	0.3
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	1.3	1.6	2.4	4.2	11.6	10.6	8.2	19.9	19.5	39.1	2.3	1.9	0.2
1966	0.0	0.0	0.0	0.0	0.8	0.5	0.5	0.9	1.3	3.1	4.0	4.1	7.2	9.4	21.4	35.5	29.6	29.5	3.0	2.5	0.3
1967	0.3	0.0	0.0	0.3	0.0	0.0	0.4	0.9	1.2	1.5	4.5	2.6	5.5	4.6	14.4	21.6	38.5	31.9	2.4	1.8	0.2
1968	0.3	0.0	0.0	0.0	0.0	0.4	0.0	1.7	1.2	1.4	3.3	10.8	4.5	8.1	16.4	14.9	22.1	34.0	2.5	2.1	0.2
1964-68	0.1	0.0	0.1	0.1	0.2	0.2	0.5	0.9	1.6	2.1	3.6	5.2	7.5	8.1	15.7	24.6	26.7	33.3	2.6	2.2	0.2
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	2.1	3.6	2.2	8.6	11.1	10.5	19.4	24.7	25.9	53.0	3.2	2.7	0.3
1970	0.0	0.0	0.0	0.0	0.0	0.8	0.0	2.2	0.9	2.2	2.7	4.7	9.3	12.7	21.1	11.3	13.6	33.0	2.5	2.2	0.3
1971	0.0	0.0	0.0	0.3	0.0	0.0	0.4	1.3	1.3	3.4	3.1	7.5	8.3	7.4	11.3	18.0	17.3	30.8	2.4	2.1	0.2
1972	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	1.3	1.3	4.4	5.7	8.1	8.0	17.2	17.5	18.9	40.8	2.5	2.1	0.2
1973	0.3	0.0	0.0	0.0	0.3	0.6	1.6	1.3	2.6	1.7	5.1	8.0	7.1	10.1	20.6	28.9	28.6	30.6	3.2	2.7	0.3
1969-73	0.1	0.0	0.0	0.1	0.1	0.3	0.7	1.1	1.6	2.4	3.6	6.9	8.7	9.7	17.9	20.2	20.9	37.2	2.8	2.3	0.3
1974	0.0	0.3	0.0	0.0	0.6	1.2	0.4	0.9	2.6	3.0	4.4	5.1	11.3	20.4	17.2	19.3	37.9	60.7	3.7	3.1	0.3
1975	0.0	0.0	0.0	0.0	1.4	0.0	2.1	2.5	2.1	1.7	4.3	6.1	8.4	8.1	9.3	23.7	27.1	17.8	2.7	2.3	0.2
1976	0.0	0.0	0.0	0.5	0.5	0.0	0.3	2.1	0.0	4.3	2.9	4.2	10.1	10.7	12.6	21.8	30.0	19.3	2.7	2.2	0.2
1977	0.7	0.0	0.0	0.0	1.3	0.3	1.3	0.8	1.3	3.4	3.4	3.5	11.7	8.9	18.3	21.3	39.7	39.3	3.2	2.6	0.3
1978	0.0	0.0	0.0	0.0	0.0	0.8	0.9	3.4	1.3	2.1	5.5	3.7	11.0	8.7	14.3	18.3	38.5	37.9	3.0	2.5	0.3
1974-78	0.1	0.1	0.0	0.1	0.7	0.5	1.0	2.0	1.5	2.9	4.1	4.5	10.5	11.2	14.4	20.8	34.8	34.8	3.0	2.6	0.3
1979	0.0	0.0	0.0	0.2	0.0	1.6	1.5	1.1	1.7	3.9	1.3	6.2	7.5	7.1	16.3	15.5	26.8	29.7	2.6	2.2	0.2
1980	0.0	0.0	0.0	0.0	0.5	0.0	1.1	1.4	0.4	0.9	3.4	4.3	6.2	9.9	15.1	16.1	27.5	22.4	2.3	1.9	0.2
1981	0.0	0.0	0.0	0.0	0.3	0.3	1.1	2.8	1.6	3.0	4.7	4.3	5.9	9.1	13.8	18.7	32.9	23.3	2.7	2.2	0.2
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.6	1.9	0.8	2.2	2.6	5.6	7.1	6.6	8.2	16.0	29.1	33.3	2.2	1.8	0.2
1983	0.3	0.0	0.0	0.0	0.2	0.0	0.0	1.5	0.8	2.6	6.0	4.7	5.7	13.2	7.9	24.1	26.4	38.7	2.7	2.2	0.2
1979-83	0.1	0.0	0.0	0.1	0.2	0.4	0.8	1.8	1.1	2.5	3.6	5.0	6.4	9.2	12.1	18.2	28.5	29.8	2.5	2.1	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF PROSTATE (ICD9 185)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	0	0	1	1	0	0	0	3	13	20	52	139	227	294	295	262	146	0	1454
1965	0	0	0	0	0	0	0	0	4	3	23	55	102	179	255	279	233	139	0	1272
1966	1	0	0	0	0	1	0	1	2	5	35	65	103	180	244	258	196	129	0	1220
1967	0	0	0	0	0	1	0	0	0	6	15	55	94	185	238	264	205	154	0	1217
1968	0	0	0	0	0	0	2	0	3	13	23	58	122	158	228	250	196	127	0	1180
1964-68	2	0	0	1	1	2	2	1	12	40	116	285	560	929	1259	1346	1092	695	0	6343
1969	0	0	0	0	1	0	3	0	0	5	14	64	118	211	254	255	207	134	0	1266
1970	0	0	0	0	0	1	0	0	5	9	16	60	117	194	213	231	198	149	0	1193
1971	0	0	1	0	0	1	0	0	2	3	20	72	132	197	235	267	193	136	0	1259
1972	0	0	0	0	1	0	0	1	3	7	27	57	148	218	253	293	216	191	0	1415
1973	1	0	0	0	0	1	0	1	2	5	31	90	157	243	286	322	223	186	11	1559
1969-73	1	0	1	0	2	3	3	2	12	29	108	343	672	1063	1241	1368	1037	796	11	6692
1974	0	0	0	0	0	0	0	2	1	11	31	74	180	256	315	281	235	173	7	1566
1975	0	0	0	0	1	2	0	1	5	11	37	91	171	271	327	321	254	171	7	1670
1976	1	0	0	0	0	1	0	0	1	12	45	73	199	271	352	326	242	182	4	1709
1977	0	0	0	0	0	1	0	1	4	8	45	91	210	295	382	317	253	191	2	1800
1978	1	0	0	0	0	0	0	2	7	11	34	96	254	342	421	396	302	233	7	2106
1974-78	2	0	0	0	1	4	0	6	18	53	192	425	1014	1435	1797	1641	1286	950	27	8851
1979	0	0	0	0	1	0	1	1	1	6	39	120	202	330	404	379	268	232	6	1990
1980	0	0	0	0	0	0	1	2	0	11	39	103	204	340	413	432	281	233	6	2065
1981	0	0	0	0	0	0	0	1	2	5	36	134	248	409	467	430	311	248	7	2298
1982	0	0	0	0	0	0	0	1	3	10	53	120	247	374	501	478	299	214	4	2304
1983	0	0	0	0	0	0	1	0	2	8	45	114	259	390	507	469	326	251	3	2375
1979-83	0	0	0	0	1	0	3	5	8	40	212	591	1160	1843	2292	2188	1485	1178	26	11032

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF PROSTATE (ICD9 185)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.0	0.0	0.4	0.5	0.0	0.0	0.0	1.4	6.9	11.8	37.2	122.0	252.2	430.5	649.8	1031.5	1073.5	44.4	40.8	4.2
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.8	1.6	13.2	38.4	87.4	195.6	367.4	607.8	899.6	1000.0	38.2	34.7	3.5
1966	0.3	0.0	0.0	0.0	0.0	0.5	0.0	0.4	0.9	2.6	19.6	44.2	86.1	193.3	346.6	553.6	753.8	921.4	36.1	33.1	3.4
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	3.0	8.2	36.2	76.5	194.7	335.7	555.8	776.5	1062.1	35.4	32.4	3.2
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	1.2	6.2	12.6	36.9	96.9	162.6	320.7	523.0	731.3	846.7	33.6	30.9	3.1
1964-68	0.1	0.0	0.0	0.1	0.1	0.2	0.2	0.1	1.0	4.1	13.1	38.6	93.5	199.0	359.6	577.2	836.8	978.9	37.4	34.3	3.5
1969	0.0	0.0	0.0	0.0	0.3	0.0	1.3	0.0	0.0	2.3	7.6	39.4	91.5	211.0	353.8	530.1	761.0	853.3	35.4	32.6	3.5
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	2.1	4.0	8.7	35.9	88.4	188.3	292.2	476.3	712.2	908.5	32.6	30.2	3.1
1971	0.0	0.0	0.2	0.0	0.0	0.3	0.0	0.0	0.8	1.3	10.6	42.3	97.0	185.7	315.9	546.0	684.4	786.1	33.6	31.0	3.2
1972	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	1.2	3.0	13.6	33.4	105.0	201.9	329.9	593.1	752.6	1067.0	37.0	34.2	3.4
1973	0.3	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.8	2.1	14.8	53.1	107.9	220.7	359.3	646.6	766.3	1022.0	39.8	36.7	3.7
1969-73	0.1	0.0	0.0	0.0	0.1	0.2	0.2	0.2	1.0	2.5	11.2	40.8	98.3	201.6	330.6	559.1	735.5	931.0	35.7	33.0	3.4
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	4.7	14.2	43.9	119.8	226.7	382.7	555.3	804.8	935.1	39.3	36.3	3.9
1975	0.0	0.0	0.0	0.0	0.3	0.6	0.0	0.4	2.1	4.6	16.6	52.9	110.2	233.6	387.4	612.6	869.9	900.0	40.9	37.6	4.0
1976	0.3	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	5.0	19.8	40.8	126.3	224.9	407.4	599.3	814.8	938.1	40.7	37.7	4.0
1977	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	1.7	3.3	19.8	48.2	133.1	234.3	432.6	561.1	837.7	969.5	41.6	38.9	4.3
1978	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.8	2.9	4.5	14.8	48.2	161.3	262.9	464.2	672.3	980.5	1165.0	47.3	44.3	4.7
1974-78	0.1	0.0	0.0	0.0	0.1	0.2	0.0	0.5	1.5	4.4	17.0	46.8	130.3	237.0	415.9	601.5	862.5	983.4	42.0	39.0	4.2
1979	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.4	0.4	2.5	16.9	57.6	128.9	245.4	431.2	619.3	842.8	1131.7	43.3	40.6	4.3
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.7	0.0	4.6	16.7	48.4	126.6	245.0	425.8	686.8	848.9	1125.6	43.8	40.8	4.2
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	2.1	15.3	62.1	146.9	289.9	463.8	664.6	912.0	1169.8	47.3	44.3	4.8
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	1.2	4.2	22.4	55.0	138.9	262.1	479.0	718.8	837.5	990.7	46.3	43.0	4.7
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.8	3.4	19.0	51.6	138.0	274.1	464.7	684.7	862.4	1125.6	46.4	43.1	4.6
1979-83	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.3	0.6	3.4	18.1	54.9	136.1	263.5	453.8	675.7	860.9	1108.2	45.5	42.4	4.5

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF TESTIS (ICD9 186)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	0	0	5	8	9	15	18	10	5	6	2	1	0	0	1	1	0	0	82
1965	0	1	0	8	9	15	18	19	16	12	6	1	0	1	3	0	1	0	0	110
1966	0	0	0	4	8	8	12	13	14	7	6	3	2	3	2	1	2	1	0	86
1967	0	1	0	4	18	14	12	10	9	2	3	2	1	1	2	1	2	0	0	82
1968	0	0	0	3	10	10	21	12	12	8	3	2	2	0	3	0	2	0	0	88
1964-68	1	2	0	24	53	56	78	72	61	34	24	10	6	5	10	3	8	1	0	448
1969	0	1	0	7	7	14	18	24	17	9	5	3	3	1	0	1	0	3	0	113
1970	0	1	0	2	11	20	16	8	8	9	7	3	3	3	1	2	0	1	0	95
1971	1	3	1	2	15	15	17	16	11	4	7	4	4	0	3	1	1	0	0	105
1972	1	0	0	9	11	18	20	17	16	14	4	1	6	0	1	1	0	3	0	122
1973	1	0	0	6	16	25	15	15	13	10	6	4	4	2	1	3	0	0	0	121
1969-73	3	5	1	26	60	92	86	80	65	46	29	15	20	6	6	8	1	7	0	556
1974	2	0	0	9	22	27	28	16	14	13	1	5	5	4	0	0	0	1	0	147
1975	3	0	1	3	13	25	21	14	15	11	10	2	2	1	0	2	0	1	1	125
1976	2	0	1	13	20	28	19	14	8	12	7	9	2	4	2	2	1	1	0	145
1977	4	0	0	4	21	30	26	19	9	7	6	5	2	4	1	2	2	1	1	144
1978	2	0	0	10	27	34	18	17	12	8	8	3	5	1	3	1	1	2	0	152
1974-78	13	0	2	39	103	144	112	80	58	51	32	24	16	14	6	7	4	6	2	713
1979	3	0	0	13	13	32	20	17	11	11	6	7	2	1	1	0	0	0	0	137
1980	3	0	1	5	21	39	33	14	14	5	10	10	4	5	2	0	0	1	0	167
1981	2	0	0	10	28	31	34	21	16	15	3	7	4	2	4	3	1	1	1	183
1982	3	0	0	4	22	34	38	16	9	13	5	8	5	3	3	1	0	0	0	164
1983	1	0	1	5	31	37	25	24	14	13	11	2	4	2	4	4	3	1	0	182
1979-83	12	0	2	37	115	173	150	92	64	57	35	34	19	13	14	8	4	3	1	833

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF TESTIS (ICD9 186)

	AGE SPECIFIC RATES																		SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.0	0.0	1.8	3.8	4.4	6.7	7.6	4.6	2.7	3.5	1.4	0.9	0.0	0.0	2.2	3.9	0.0	2.4	2.5	0.2
1965	0.0	0.3	0.0	2.8	4.0	7.2	8.0	8.0	7.1	6.3	3.4	0.7	0.0	1.1	4.3	0.0	3.9	0.0	3.2	3.3	0.3
1966	0.0	0.0	0.0	1.3	3.3	3.7	5.3	5.4	6.0	3.6	3.4	2.0	1.7	3.2	2.8	2.1	7.7	7.1	2.4	2.5	0.2
1967	0.0	0.3	0.0	1.3	6.9	6.2	5.3	4.2	3.8	1.0	1.6	1.3	0.8	1.1	2.8	2.1	7.6	0.0	2.3	2.3	0.2
1968	0.0	0.0	0.0	0.9	3.6	4.2	9.2	5.0	5.0	3.8	1.6	1.3	1.6	0.0	4.2	0.0	7.5	0.0	2.4	2.4	0.2
1964-68	0.1	0.1	0.0	1.6	4.3	5.1	6.9	6.0	5.3	3.5	2.7	1.4	1.0	1.1	2.9	1.3	6.1	1.4	2.5	2.6	0.2
1969	0.0	0.2	0.0	2.1	2.4	5.6	7.8	10.1	7.0	4.1	2.7	1.8	2.3	1.0	0.0	2.1	0.0	19.1	3.0	3.0	0.2
1970	0.0	0.2	0.0	0.6	3.5	7.4	6.8	3.4	3.3	4.0	3.8	1.8	2.3	2.9	1.4	4.1	0.0	6.1	2.5	2.5	0.2
1971	0.3	0.7	0.2	0.6	4.5	5.2	7.0	6.7	4.5	1.7	3.7	2.3	2.9	0.0	4.0	2.0	3.5	0.0	2.7	2.7	0.2
1972	0.3	0.0	0.0	2.4	3.3	5.9	8.0	7.2	6.6	6.0	2.0	0.6	4.3	0.0	1.3	2.0	0.0	16.8	3.1	3.1	0.2
1973	0.3	0.0	0.0	1.6	4.7	7.8	5.8	6.3	5.3	4.3	2.9	2.4	2.7	1.8	1.3	6.0	0.0	0.0	3.0	3.0	0.2
1969-73	0.2	0.3	0.0	1.4	3.7	6.4	7.1	6.7	5.4	4.0	3.0	1.8	2.9	1.1	1.6	3.3	0.7	8.2	2.8	2.9	0.2
1974	0.6	0.0	0.0	2.3	6.3	8.1	10.2	6.7	5.7	5.5	0.5	3.0	3.3	3.5	0.0	0.0	0.0	5.4	3.5	3.5	0.3
1975	0.9	0.0	0.2	0.7	3.6	7.2	7.3	5.7	6.2	4.6	4.5	1.2	1.3	0.9	0.0	3.8	0.0	5.3	2.9	2.9	0.2
1976	0.6	0.0	0.2	3.1	5.4	7.9	6.4	5.7	3.3	5.0	3.1	5.0	1.3	3.3	2.3	3.7	3.4	5.2	3.4	3.3	0.3
1977	1.3	0.0	0.0	1.0	5.6	8.5	8.2	7.5	3.8	2.9	2.6	2.6	1.3	3.2	1.1	3.5	6.6	5.1	3.3	3.2	0.2
1978	0.6	0.0	0.0	2.4	7.1	9.6	5.5	6.4	5.0	3.3	3.5	1.5	3.2	0.8	3.3	1.7	3.2	10.0	3.5	3.4	0.3
1974-78	0.8	0.0	0.1	1.9	5.6	8.3	7.5	6.4	4.8	4.3	2.8	2.6	2.1	2.3	1.4	2.6	2.7	6.2	3.3	3.3	0.3
1979	1.0	0.0	0.0	3.1	3.4	9.0	5.9	6.2	4.6	4.6	2.6	3.4	1.3	0.7	1.1	0.0	0.0	0.0	3.1	3.0	0.2
1980	1.0	0.0	0.3	1.2	5.4	10.9	9.4	5.0	5.8	2.1	4.3	4.7	2.5	3.6	2.1	0.0	0.0	4.8	3.6	3.6	0.3
1981	0.7	0.0	0.0	2.4	7.1	8.6	9.5	7.2	6.5	6.3	1.3	3.2	2.4	1.4	4.0	4.6	2.9	4.7	4.0	3.8	0.3
1982	1.0	0.0	0.0	1.0	5.5	9.3	10.8	5.1	3.5	5.5	2.1	3.7	2.8	2.1	2.9	1.5	0.0	0.0	3.5	3.4	0.3
1983	0.3	0.0	0.3	1.3	7.5	9.9	7.1	7.3	5.3	5.5	4.6	0.9	2.1	1.4	3.7	5.8	7.9	4.5	3.9	3.7	0.3
1979-83	0.8	0.0	0.1	1.8	5.8	9.6	8.6	6.2	5.1	4.8	3.0	3.2	2.2	1.9	2.8	2.5	2.3	2.8	3.6	3.5	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF PENIS AND OTHER MALE GENITAL ORGANS (ICD9 187)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	1	0	3	3	1	3	5	4	5	3	6	5	3	0	42
1965	0	0	0	0	0	0	0	2	0	0	4	5	3	4	8	6	6	5	0	43
1966	0	0	0	0	0	0	0	2	1	3	2	6	7	3	4	4	2	3	0	37
1967	0	0	0	0	0	0	0	1	3	4	2	1	5	3	8	0	1	3	0	31
1968	0	0	0	1	0	0	1	0	2	3	1	4	4	2	4	4	1	2	0	29
1964-68	0	0	0	1	0	1	1	8	9	11	12	21	23	17	27	20	15	16	0	182
1969	0	0	0	1	0	0	0	0	2	2	2	4	3	4	10	2	5	3	0	38
1970	0	0	0	0	0	0	1	1	1	1	3	5	6	2	5	6	5	4	0	40
1971	0	0	0	0	0	0	0	2	0	2	5	5	4	4	6	4	3	1	0	36
1972	0	1	0	0	0	0	0	1	0	3	3	5	2	3	6	1	2	*	0	29
1973	0	0	0	0	0	0	0	0	1	1	1	0	3	8	4	4	2	0	0	24
1969-73	0	1	0	1	0	0	1	4	4	9	14	19	18	21	31	17	17	10	0	167
1974	0	0	0	0	0	1	1	1	1	3	5	5	4	5	2	9	3	3	3	46
1975	0	0	0	0	0	0	0	1	0	3	2	0	8	3	3	4	4	5	0	33
1976	0	0	0	0	0	0	1	0	1	3	5	6	7	7	5	3	5	2	0	45
1977	0	0	0	1	0	0	0	0	2	9	2	7	8	6	5	5	4	5	0	54
1978	0	0	0	0	0	1	1	0	0	1	1	2	6	4	3	5	5	2	0	31
1974-78	0	0	0	1	0	2	3	2	4	19	15	20	33	25	18	26	21	17	3	209
1979	0	0	0	0	1	1	1	1	0	1	3	4	5	7	7	4	1	1	0	37
1980	0	0	0	0	0	1	0	1	1	2	0	1	5	11	8	5	2	4	0	41
1981	0	0	0	0	0	0	2	0	2	1	7	6	6	10	6	3	6	2	0	51
1982	1	0	0	1	1	0	0	2	1	4	6	8	5	7	5	7	1	1	0	50
1983	0	0	1	1	0	2	1	0	1	1	3	6	5	7	7	4	2	5	0	46
1979-83	1	0	1	2	2	4	4	4	5	9	19	25	26	42	33	23	12	13	0	225

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF PENIS AND OTHER MALE GENITAL ORGANS (ICD9 187)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.5	0.0	1.3	1.4	0.5	1.8	3.6	3.5	5.6	4.4	13.2	19.7	22.1	1.3	1.2	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	2.3	3.5	2.6	4.4	11.5	13.1	23.2	36.0	1.3	1.2	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	1.5	1.1	4.1	5.9	3.2	5.7	8.5	7.7	21.4	1.1	1.1	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	2.0	1.1	0.7	4.1	3.2	11.3	0.0	3.8	20.7	0.9	0.9	0.1
1968	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	0.8	1.4	0.5	2.5	3.2	2.1	5.6	8.4	3.7	13.3	0.8	0.8	0.1
1964-68	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.7	0.8	1.1	1.4	2.8	3.8	3.6	7.7	8.6	11.5	22.5	1.1	1.0	0.1
1969	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.8	0.9	1.1	2.5	2.3	4.0	13.9	4.2	18.4	19.1	1.0	1.0	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.4	1.6	3.0	4.5	1.9	6.9	12.4	18.0	24.4	1.1	1.0	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.9	2.6	2.9	2.9	3.8	8.1	8.2	10.6	5.8	0.9	0.9	0.1
1972	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.3	1.5	2.9	1.4	2.8	7.8	2.0	7.0	11.2	0.7	0.7	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.0	2.1	7.3	5.0	8.0	6.9	0.0	0.6	0.6	0.1
1969-73	0.0	0.1	0.0	0.1	0.0	0.0	0.1	0.3	0.3	0.8	1.5	2.3	2.6	4.0	8.3	6.9	12.1	11.7	0.9	0.8	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.4	0.4	1.3	2.3	3.0	2.7	4.4	2.4	17.8	10.3	16.2	1.1	1.0	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.3	0.9	0.0	5.2	2.6	3.6	7.6	13.7	26.3	0.8	0.8	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	1.3	2.2	3.4	4.4	5.8	5.8	5.5	16.8	10.3	1.1	1.0	0.1
1977	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.8	3.7	0.9	3.7	5.1	4.8	5.7	8.8	13.2	25.4	1.2	1.2	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.4	0.4	1.0	3.8	3.1	3.3	8.5	16.2	10.0	0.7	0.7	0.1
1974-78	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.2	0.3	1.6	1.3	2.2	4.2	4.1	4.2	9.5	14.1	17.6	1.0	0.9	0.1
1979	0.0	0.0	0.0	0.0	0.3	0.3	0.3	0.4	0.0	0.4	1.3	1.9	3.2	5.2	7.5	6.5	3.1	4.9	0.8	0.8	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.4	0.8	0.0	0.5	3.1	7.9	8.2	7.9	6.0	19.3	0.9	0.9	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.8	0.4	3.0	2.8	3.6	7.1	6.0	4.6	17.6	9.4	1.1	1.0	0.1
1982	0.3	0.0	0.0	0.2	0.2	0.0	0.0	0.6	0.4	1.7	2.5	3.7	2.8	4.9	4.8	10.5	2.8	4.6	1.0	1.0	0.1
1983	0.0	0.0	0.3	0.3	0.0	0.5	0.3	0.0	0.4	0.4	1.3	2.7	2.7	4.9	6.4	5.8	5.3	22.4	0.9	0.9	0.1
1979-83	0.1	0.0	0.1	0.1	0.1	0.2	0.2	0.3	0.4	0.8	1.6	2.3	3.1	6.0	6.5	7.1	7.0	12.2	0.9	0.9	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF BLADDER (ICD9 188)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	1	1	2	0	1	5	6	15	17	46	78	117	109	123	87	65	37	0	710
1965	0	0	0	0	0	2	2	8	15	29	31	61	69	97	115	79	52	34	1	595
1966	0	0	0	1	1	2	4	9	16	34	37	57	81	97	85	70	49	26	0	569
1967	0	0	0	0	1	5	5	12	12	27	39	49	76	93	79	81	56	39	0	574
1968	0	0	0	0	2	2	5	3	15	24	44	68	82	108	91	88	64	44	0	640
1964-68	0	1	1	3	4	12	21	38	73	131	197	313	425	504	493	405	286	180	1	3088
1969	0	0	0	1	6	0	8	10	13	32	45	60	82	93	109	103	60	33	0	655
1970	0	0	0	1	1	2	7	10	13	26	34	80	88	120	98	82	44	59	0	665
1971	0	0	0	0	1	0	2	7	18	35	32	55	91	115	109	75	70	42	0	652
1972	0	1	0	1	0	3	5	6	14	23	53	87	93	98	97	93	80	42	0	696
1973	0	0	0	0	1	5	6	10	20	36	47	77	121	121	125	135	63	44	4	815
1969-73	0	1	0	3	9	10	28	43	78	152	211	359	475	547	538	488	317	220	4	3483
1974	0	0	0	1	5	10	7	16	24	36	46	77	114	144	155	102	68	63	11	879
1975	0	1	0	2	1	2	7	12	18	40	63	94	92	124	132	109	64	51	7	819
1976	0	0	0	0	2	5	7	7	17	38	68	74	113	142	149	112	68	54	3	859
1977	0	0	1	0	1	8	8	15	18	49	73	85	104	161	140	135	80	54	4	936
1978	0	0	0	2	2	7	6	11	27	41	70	96	135	193	150	138	97	81	0	1056
1974-78	0	1	1	5	11	32	35	61	104	204	320	426	558	764	726	596	377	303	25	4549
1979	0	0	0	4	2	9	6	13	24	42	92	130	146	162	167	126	87	68	2	1080
1980	0	0	1	0	2	6	10	14	21	40	65	93	130	181	212	144	95	64	4	1082
1981	1	0	0	0	3	5	11	21	19	41	79	118	126	187	204	148	115	83	1	1162
1982	0	0	0	0	2	8	12	13	26	40	80	111	134	178	198	153	89	68	2	1114
1983	0	0	0	2	3	6	7	18	23	45	60	128	150	192	211	145	97	76	4	1167
1979-83	1	0	1	6	12	34	46	79	113	208	376	580	686	900	992	716	483	359	13	5605

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF BLADDER (ICD9 188)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.3	0.3	0.7	0.0	0.5	2.2	2.5	6.9	9.0	27.1	55.8	102.7	121.1	180.1	191.6	255.9	272.1	21.5	20.9	2.5
1965	0.0	0.0	0.0	0.0	0.0	1.0	0.9	3.4	6.6	15.2	17.7	42.6	59.1	106.0	165.7	172.1	200.8	244.6	17.6	17.0	2.1
1966	0.0	0.0	0.0	0.3	0.4	0.9	1.8	3.8	6.9	17.4	20.7	38.7	67.7	104.2	120.7	150.2	188.5	185.7	16.6	16.1	1.9
1967	0.0	0.0	0.0	0.0	0.4	2.2	2.2	5.0	5.0	13.4	21.4	32.2	61.8	97.9	111.4	170.5	212.1	269.0	16.4	15.9	1.7
1968	0.0	0.0	0.0	0.0	0.7	0.8	2.2	1.3	6.2	11.5	24.1	43.3	65.1	111.1	128.0	184.1	238.8	293.3	18.0	17.3	2.0
1964-68	0.0	0.1	0.1	0.2	0.3	1.1	1.8	3.2	6.3	13.3	22.2	42.3	71.0	108.0	140.8	173.7	219.2	253.5	18.0	17.4	2.0
1969	0.0	0.0	0.0	0.3	2.0	0.0	3.5	4.2	5.4	14.7	24.6	36.9	63.6	93.0	151.8	214.1	220.6	210.2	18.0	17.2	2.0
1970	0.0	0.0	0.0	0.3	0.3	0.7	3.0	4.2	5.4	11.5	18.4	47.9	66.5	116.5	134.4	169.1	158.3	359.8	17.9	17.5	2.0
1971	0.0	0.0	0.0	0.0	0.3	0.0	0.8	2.9	7.4	15.1	16.9	32.3	66.9	108.4	146.5	153.4	248.2	242.8	17.1	16.6	2.0
1972	0.0	0.3	0.0	0.3	0.0	1.0	2.0	2.5	5.8	9.9	26.7	50.9	66.0	90.7	126.5	188.3	278.7	234.6	17.9	17.0	1.9
1973	0.0	0.0	0.0	0.0	0.3	1.6	2.3	4.2	8.2	15.3	22.5	45.4	83.2	109.9	157.0	271.1	216.5	241.8	20.5	19.7	2.2
1969-73	0.0	0.1	0.0	0.2	0.6	0.7	2.3	3.6	6.4	13.3	21.9	42.7	69.5	103.8	143.3	199.4	224.8	257.3	18.3	17.6	2.0
1974	0.0	0.0	0.0	0.3	1.4	3.0	2.5	6.7	9.6	15.2	21.0	45.7	75.8	127.5	188.3	201.6	232.9	340.5	21.5	20.8	2.5
1975	0.0	0.3	0.0	0.5	0.3	0.6	2.4	4.9	7.4	16.8	28.2	54.7	59.3	106.9	156.4	208.0	219.2	268.4	19.7	18.9	2.2
1976	0.0	0.0	0.0	0.0	0.5	1.4	2.4	2.8	7.1	15.9	29.9	41.3	71.7	117.8	172.5	205.9	229.0	278.4	20.1	19.4	2.3
1977	0.0	0.0	0.3	0.0	0.3	2.3	2.5	5.9	7.5	20.4	32.1	45.0	65.9	127.9	158.6	238.9	264.9	274.1	21.4	20.5	2.3
1978	0.0	0.0	0.0	0.5	0.5	2.0	1.8	4.1	11.3	17.0	30.6	48.2	85.7	148.3	165.4	234.3	314.9	405.0	23.6	22.9	2.5
1974-78	0.0	0.1	0.0	0.2	0.6	1.8	2.3	4.9	8.6	17.1	28.4	46.9	71.7	126.2	168.0	218.5	252.9	313.7	21.3	20.5	2.4
1979	0.0	0.0	0.0	1.0	0.5	2.5	1.8	4.7	10.0	17.4	40.0	62.4	93.2	120.4	178.2	205.9	273.6	331.7	23.5	22.8	2.6
1980	0.0	0.0	0.3	0.0	0.5	1.7	2.9	5.0	8.6	16.8	27.9	43.7	80.6	130.4	218.6	228.9	287.0	309.2	22.8	22.1	2.6
1981	0.3	0.0	0.0	0.0	0.8	1.4	3.1	7.2	7.7	17.3	33.5	54.7	74.6	132.5	202.6	228.7	337.2	391.5	24.1	23.1	2.6
1982	0.0	0.0	0.0	0.0	0.5	2.2	3.4	4.2	10.2	17.0	33.8	50.9	75.4	124.7	189.3	230.1	249.3	314.8	22.5	21.7	2.5
1983	0.0	0.0	0.0	0.5	0.7	1.6	2.0	5.5	8.7	19.1	25.3	57.9	79.9	134.9	193.4	211.7	256.6	340.8	22.9	22.1	2.6
1979-83	0.1	0.0	0.1	0.3	0.6	1.9	2.6	5.3	9.0	17.5	32.1	53.9	80.5	128.7	196.4	221.1	280.0	337.7	23.2	22.4	2.6

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF BLADDER (ICD9 188)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	1	0	1	4	7	9	9	16	22	37	47	28	43	20	0	244
1965	0	0	0	0	0	0	3	4	3	7	8	17	20	33	37	28	33	24	0	217
1966	0	0	0	0	0	2	2	2	4	9	11	10	21	24	35	39	30	20	0	209
1967	0	0	0	0	2	0	1	0	8	10	17	22	17	39	42	42	28	24	0	252
1968	0	0	0	1	1	1	0	1	3	10	13	28	28	27	28	30	25	30	0	226
1964-68	0	0	0	1	4	3	7	11	25	45	58	93	108	160	189	167	159	118	0	1148
1969	1	0	0	1	0	2	5	3	7	9	10	14	29	37	51	49	36	24	0	278
1970	0	0	0	2	0	1	2	2	4	16	13	22	25	22	30	40	31	18	0	228
1971	0	0	0	0	2	3	1	5	3	9	7	25	24	27	45	41	33	22	0	247
1972	0	0	0	1	0	1	1	3	4	13	16	23	21	31	35	36	33	29	0	247
1973	0	0	0	1	0	1	4	2	3	22	19	24	34	37	47	42	39	25	2	302
1969-73	1	0	0	5	2	8	13	15	21	69	65	108	133	154	208	208	172	118	2	1302
1974	0	0	0	0	0	1	0	5	8	10	23	16	42	22	55	45	38	40	3	308
1975	0	0	0	1	1	4	1	9	8	13	22	20	48	38	44	44	42	40	1	336
1976	3	0	0	0	3	3	3	3	2	6	24	25	24	56	51	37	42	41	1	321
1977	0	0	0	0	0	1	2	2	11	13	17	23	30	49	58	61	41	29	2	339
1978	1	0	0	1	1	2	6	5	11	14	28	19	42	59	50	54	44	40	0	377
1974-78	1	0	0	2	5	11	12	24	40	56	114	103	186	224	258	241	207	190	7	1681
1979	0	0	0	2	2	3	2	7	10	6	20	34	50	43	50	53	41	47	0	370
1980	0	0	0	0	3	3	3	8	5	9	20	43	54	45	64	57	54	41	0	409
1981	0	0	0	0	1	3	2	7	8	14	28	46	45	61	52	43	43	47	1	401
1982	0	0	0	0	1	3	3	4	4	9	25	34	35	52	62	52	46	46	0	376
1983	0	0	0	0	3	0	0	7	7	13	41	36	41	49	71	43	46	57	0	414
1979-83	0	0	0	2	10	12	10	33	34	51	134	193	225	250	299	248	230	238	1	1970

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF BLADDER (ICD9 188)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.5	0.0	0.5	1.7	3.2	4.9	5.5	11.6	18.6	36.4	56.6	47.8	124.3	90.9	7.4	5.9	0.7
1965	0.0	0.0	0.0	0.0	0.0	0.0	1.4	1.7	1.3	3.7	4.7	12.0	16.5	31.9	43.2	46.4	91.9	104.3	6.4	5.1	0.6
1966	0.0	0.0	0.0	0.0	0.0	0.9	0.9	0.9	1.7	4.6	6.3	6.8	16.9	22.6	39.5	63.0	80.6	84.4	6.0	4.7	0.5
1967	0.0	0.0	0.0	0.0	0.8	0.0	0.4	0.0	3.3	4.9	9.5	14.5	13.2	35.9	46.6	64.9	71.8	95.6	7.0	5.7	0.6
1968	0.0	0.0	0.0	0.3	0.4	0.4	0.0	0.4	1.2	4.7	7.2	17.8	21.2	24.3	30.7	44.8	61.4	113.2	6.1	5.1	0.5
1964-68	0.0	0.0	0.0	0.1	0.3	0.3	0.6	0.9	2.1	4.6	6.7	12.7	17.3	30.1	43.1	53.4	84.8	98.1	6.6	5.3	0.6
1969	0.3	0.0	0.0	0.3	0.0	0.8	2.2	1.3	3.0	4.1	5.5	8.6	21.4	32.3	55.1	71.2	84.9	84.8	7.3	5.9	0.7
1970	0.0	0.0	0.0	0.6	0.0	0.4	0.9	0.9	1.7	6.9	7.0	13.0	17.9	18.6	31.7	56.7	70.1	59.4	5.8	4.7	0.5
1971	0.0	0.0	0.0	0.0	0.6	1.1	0.4	2.2	1.3	3.8	3.6	14.4	16.7	22.2	46.3	56.9	71.4	67.7	6.1	4.9	0.6
1972	0.0	0.0	0.0	0.3	0.0	0.3	0.4	1.3	1.7	5.5	7.8	13.2	14.1	24.8	35.3	48.5	69.5	84.5	5.9	4.8	0.5
1973	0.0	0.0	0.0	0.3	0.0	0.3	1.6	0.9	1.3	9.3	8.8	13.7	22.1	28.8	46.2	55.3	79.6	69.6	7.0	5.8	0.7
1969-73	0.1	0.0	0.0	0.3	0.1	0.6	1.1	1.3	1.8	5.9	6.6	12.6	18.4	25.3	42.9	57.5	75.0	73.2	6.4	5.2	0.6
1974	0.0	0.0	0.0	0.0	0.0	0.3	0.0	2.1	3.4	4.2	10.1	9.1	26.3	16.6	52.4	57.8	75.8	105.5	6.9	5.6	0.6
1975	0.0	0.0	0.0	0.3	0.3	1.1	0.4	3.8	3.4	5.5	9.4	11.0	28.9	28.1	40.7	55.0	81.4	101.5	7.4	6.1	0.7
1976	0.0	0.0	0.0	0.0	0.8	0.8	1.0	1.2	0.9	2.6	10.1	13.1	14.2	39.9	46.0	44.7	78.8	99.0	6.8	5.5	0.7
1977	0.0	0.0	0.0	0.0	0.0	0.3	0.6	0.8	4.7	5.6	7.2	11.4	17.5	33.7	50.5	72.0	74.0	67.0	7.0	5.7	0.7
1978	0.3	0.0	0.0	0.2	0.3	0.6	1.8	1.9	4.7	6.0	11.9	8.9	24.4	39.3	42.1	61.8	76.9	89.1	7.6	6.4	0.7
1974-78	0.1	0.0	0.0	0.1	0.3	0.6	0.8	2.0	3.4	4.8	9.7	10.7	22.2	31.8	46.3	58.4	77.4	91.8	7.1	5.9	0.7
1979	0.0	0.0	0.0	0.5	0.5	0.8	0.6	2.6	4.2	2.6	8.6	15.1	28.9	27.8	40.7	58.8	68.8	99.8	7.2	6.0	0.7
1980	0.0	0.0	0.0	0.0	0.8	0.8	0.8	2.9	2.1	3.9	8.6	18.7	30.2	27.9	50.7	61.1	87.4	83.5	7.8	6.4	0.7
1981	0.0	0.0	0.0	0.0	0.3	0.8	0.5	2.4	3.3	6.0	12.0	19.7	24.0	37.1	39.9	44.7	67.3	91.3	7.4	6.3	0.7
1982	0.0	0.0	0.0	0.0	0.2	0.8	0.8	1.3	1.6	3.9	10.7	14.6	17.7	31.1	46.1	51.9	70.3	85.2	6.7	5.5	0.6
1983	0.0	0.0	0.0	0.0	0.7	0.0	0.0	2.1	2.7	5.6	17.6	15.3	19.5	29.3	50.9	41.5	67.5	100.2	7.3	6.1	0.7
1979-83	0.0	0.0	0.0	0.1	0.5	0.6	0.6	2.2	2.8	4.4	11.5	16.7	23.7	30.7	45.8	51.3	72.1	92.0	7.3	6.1	0.7

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF KIDNEY AND OTHER AND UNSPECIFIED URINARY ORGANS (ICD9 189)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	5	3	0	0	3	0	3	4	10	12	24	37	31	31	27	15	12	3	0	220
1965	11	3	1	1	0	2	1	7	6	15	30	30	27	29	22	19	14	5	0	223
1966	5	0	0	0	0	0	2	3	7	16	17	29	25	24	17	18	11	3	0	177
1967	9	3	0	0	1	3	1	3	7	26	24	28	30	31	26	26	12	5	0	235
1968	8	5	0	0	0	2	2	5	13	14	24	23	37	18	17	14	14	6	0	202
1964-68	38	14	1	1	4	7	9	22	43	83	119	147	150	133	109	92	63	22	0	1057
1969	7	2	0	1	0	1	0	8	15	17	27	29	32	41	24	24	10	4	0	242
1970	12	3	0	1	1	1	3	10	10	18	25	32	31	24	37	26	6	7	0	247
1971	7	2	0	0	0	0	5	6	12	15	19	30	34	33	35	21	10	7	0	236
1972	10	1	0	0	2	2	3	4	11	18	11	31	38	38	31	20	15	7	0	242
1973	6	2	3	1	1	3	7	8	13	23	32	27	42	36	31	21	10	12	0	278
1969-73	42	10	3	3	4	7	18	36	61	91	114	149	177	172	158	112	51	37	0	1245
1974	5	2	1	0	2	1	1	6	16	32	27	23	36	36	35	29	12	7	2	273
1975	5	0	2	2	1	1	3	2	12	30	32	35	45	47	31	29	12	8	0	297
1976	7	0	0	1	0	5	6	16	15	27	30	41	44	42	24	28	11	3	0	300
1977	8	5	0	1	1	1	3	3	11	22	25	44	42	43	27	22	13	6	0	277
1978	2	1	0	1	0	2	0	3	17	18	29	56	40	43	41	34	12	9	1	309
1974-78	27	8	3	5	4	10	13	30	71	129	143	199	207	211	158	142	60	33	3	1456
1979	3	1	1	0	2	1	2	6	17	32	34	52	37	54	40	21	22	8	0	333
1980	5	1	2	1	2	2	5	9	10	16	41	56	47	57	44	35	20	7	0	360
1981	1	2	0	1	1	1	1	6	8	20	40	59	61	59	53	33	16	13	0	375
1982	7	3	1	0	3	1	3	7	13	27	44	61	72	53	42	37	15	9	3	401
1983	7	2	0	0	0	0	3	4	17	29	34	62	61	51	60	34	27	14	2	407
1979-83	23	9	4	2	8	5	14	32	65	124	193	290	278	274	239	160	100	51	5	1876

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASMS OF KIDNEY AND OTHER AND UNSPECIFIED URINARY ORGANS (ICD9 189)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	1.3	0.8	0.0	0.0	1.4	0.0	1.3	1.7	4.6	6.4	14.1	26.5	27.2	34.4	39.5	33.0	47.2	22.1	6.6	6.5	0.8
1965	2.8	0.8	0.3	0.3	0.0	1.0	0.4	2.9	2.7	7.9	17.2	20.9	23.1	31.7	31.7	41.4	54.1	36.0	6.6	6.5	0.7
1966	1.3	0.0	0.0	0.0	0.0	0.0	0.9	1.3	3.0	8.2	9.5	19.7	20.9	25.8	24.1	38.6	42.3	21.4	5.1	5.0	0.6
1967	2.4	0.8	0.0	0.0	0.4	1.3	0.4	1.2	2.9	12.9	13.2	18.4	24.4	32.6	36.7	54.7	45.5	34.5	6.6	6.6	0.7
1968	2.2	1.2	0.0	0.0	0.0	0.8	0.9	2.1	5.4	6.7	13.1	14.6	29.4	18.5	23.9	29.3	52.2	40.0	5.6	5.6	0.6
1964-68	2.0	0.7	0.1	0.1	0.3	0.6	0.8	1.8	3.7	8.4	13.4	19.9	25.0	28.5	31.1	39.5	48.3	31.0	6.1	6.0	0.7
1969	2.0	0.5	0.0	0.3	0.0	0.4	0.0	3.4	6.2	7.8	14.7	17.9	24.8	41.0	33.4	49.9	36.8	25.5	6.5	6.5	0.8
1970	3.6	0.7	0.0	0.3	0.3	0.4	1.3	4.2	4.1	8.0	13.5	19.2	23.4	23.3	50.8	53.6	21.6	42.7	6.5	6.6	0.8
1971	2.1	0.5	0.0	0.0	0.0	0.0	2.1	2.5	5.0	6.5	10.1	17.6	25.0	31.1	47.0	42.9	35.5	40.5	6.1	6.2	0.7
1972	3.1	0.3	0.0	0.0	0.6	0.7	1.2	1.7	4.5	7.7	5.5	18.1	27.0	35.2	40.4	40.5	52.3	39.1	6.2	6.2	0.7
1973	1.9	0.5	0.7	0.3	0.3	0.9	2.7	3.4	5.3	9.8	15.3	15.9	28.9	32.7	38.9	42.2	34.4	65.9	6.9	7.0	0.8
1969-73	2.5	0.5	0.1	0.2	0.2	0.5	1.5	3.0	5.0	7.9	11.8	17.7	25.9	32.6	42.1	45.8	36.2	43.3	6.5	6.5	0.8
1974	1.5	0.6	0.2	0.0	0.6	0.3	0.4	2.5	6.5	13.5	12.3	13.6	24.0	31.9	42.5	57.3	41.1	37.8	6.7	6.6	0.7
1975	1.6	0.0	0.5	0.5	0.3	0.3	1.0	0.8	4.9	12.6	14.3	20.3	29.0	40.5	36.7	55.3	41.1	42.1	7.1	7.1	0.8
1976	2.2	0.0	0.0	0.2	0.0	1.4	2.0	6.5	6.2	11.3	13.2	22.9	27.9	34.9	27.8	51.5	37.0	15.5	7.0	7.0	0.8
1977	2.6	1.5	0.0	0.2	0.3	0.3	0.9	1.2	4.6	9.2	11.0	23.3	26.6	34.2	30.6	38.9	43.0	30.5	6.4	6.4	0.7
1978	0.6	0.3	0.0	0.2	0.0	0.6	0.0	1.1	7.1	7.4	12.7	28.1	25.4	33.1	45.2	57.7	39.0	45.0	6.9	6.8	0.8
1974-78	1.7	0.5	0.1	0.2	0.2	0.6	0.9	2.4	5.9	10.8	12.7	21.9	26.6	34.9	36.6	52.1	40.2	34.2	6.8	6.8	0.8
1979	1.0	0.3	0.3	0.0	0.5	0.3	0.6	2.2	7.1	13.3	14.8	24.9	23.6	40.1	42.7	34.3	69.2	39.0	7.3	7.2	0.9
1980	1.6	0.3	0.6	0.2	0.5	0.6	1.4	3.2	4.1	6.7	17.6	26.3	29.2	41.1	45.4	55.6	60.4	33.8	7.7	7.6	0.9
1981	0.3	0.6	0.0	0.2	0.3	0.3	0.3	2.1	3.2	8.5	17.0	27.3	36.1	41.8	52.6	51.0	46.9	61.3	7.8	7.8	0.9
1982	2.3	1.0	0.3	0.0	0.7	0.3	0.9	2.2	5.1	11.5	18.6	28.0	40.5	37.1	40.2	55.6	42.0	41.7	8.2	8.2	0.9
1983	2.3	0.6	0.0	0.0	0.0	0.0	0.9	1.2	6.4	12.3	14.3	28.1	32.5	35.8	55.0	49.6	71.4	62.8	8.1	8.1	0.9
1979-83	1.5	0.6	0.2	0.1	0.4	0.3	0.8	2.1	5.2	10.4	16.5	26.9	32.6	39.2	47.3	49.4	58.0	48.0	7.8	7.8	0.9

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF KIDNEY AND OTHER AND UNSPECIFIED URINARY ORGANS (ICD9 189)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	10	0	0	1	1	3	2	2	8	7	10	16	14	17	21	19	9	6	0	146
1965	10	4	0	0	0	0	2	1	3	11	11	10	13	15	14	17	8	10	0	129
1966	7	4	1	1	0	1	3	1	9	3	12	15	9	15	19	11	7	2	0	120
1967	7	6	1	1	1	3	1	1	3	6	10	17	19	20	23	8	7	3	0	137
1968	7	4	2	0	0	0	1	6	5	5	9	15	11	16	19	18	5	7	0	130
1964-68	41	18	4	3	2	7	9	11	28	32	52	73	66	83	96	73	36	28	0	662
1969	5	2	0	0	0	2	1	2	1	8	8	19	19	19	20	17	12	7	0	142
1970	7	1	0	0	1	1	1	5	5	10	11	15	14	11	15	11	12	6	1	127
1971	6	0	0	1	0	0	3	2	5	7	17	12	29	14	17	11	17	8	1	150
1972	3	4	0	1	1	1	2	4	4	8	13	15	26	21	15	14	11	3	0	146
1973	7	4	0	0	3	0	0	4	8	13	14	22	24	16	18	16	5	9	0	163
1969-73	28	11	0	2	5	4	7	17	23	46	63	83	112	81	85	69	57	33	2	728
1974	9	3	1	0	0	2	3	0	6	10	19	16	21	29	26	14	15	5	0	179
1975	11	1	0	2	1	0	1	4	5	10	16	13	20	19	18	17	9	10	0	157
1976	7	3	0	0	1	0	0	6	2	9	16	19	24	19	15	20	7	10	0	158
1977	6	0	0	1	0	1	0	5	5	14	12	23	20	32	18	15	14	6	0	174
1978	6	0	0	0	2	1	1	3	6	11	13	21	26	17	24	21	24	6	0	182
1974-78	39	7	1	3	4	4	5	18	24	54	76	92	111	116	101	87	71	37	0	850
1979	4	4	0	1	0	2	2	8	7	8	23	21	22	25	41	17	11	15	0	211
1980	3	6	1	2	0	3	1	11	4	12	14	34	28	27	23	23	12	3	0	207
1981	2	3	1	1	0	0	3	6	5	10	19	25	17	31	39	22	12	8	0	204
1982	6	3	1	0	3	0	2	6	8	9	22	24	34	33	32	27	11	8	0	229
1983	8	3	0	0	2	0	3	3	7	12	18	24	21	37	39	33	18	16	0	244
1979-83	23	19	3	4	5	5	11	34	31	51	96	128	122	153	174	122	64	50	0	1095

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF KIDNEY AND OTHER AND UNSPECIFIED URINARY ORGANS (ICD9 189)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	2.7	0.0	0.0	0.4	0.5	1.4	0.9	0.9	3.6	3.8	6.1	11.6	11.9	16.7	25.3	32.4	26.0	27.3	4.4	3.9	0.4
1965	2.7	1.1	0.0	0.0	0.0	0.0	0.9	0.4	1.3	5.8	6.5	7.0	10.7	14.5	16.4	28.1	22.3	43.5	3.8	3.3	0.3
1966	1.9	1.1	0.3	0.3	0.0	0.5	1.4	0.4	3.8	1.5	6.9	10.3	7.2	14.1	21.4	17.8	18.8	8.4	3.4	3.1	0.4
1967	2.0	1.6	0.3	0.3	0.4	1.3	0.4	0.4	1.2	3.0	5.6	11.2	14.8	18.4	25.5	12.4	17.9	12.0	3.8	3.5	0.4
1968	2.1	1.0	0.6	0.0	0.0	0.0	0.4	2.6	2.1	2.4	5.0	9.5	8.3	14.4	20.8	26.9	12.3	26.4	3.5	3.1	0.3
1964-68	2.3	1.0	0.2	0.2	0.2	0.6	0.8	0.9	2.4	3.2	6.0	9.9	10.6	15.6	21.9	23.4	19.2	23.3	3.8	3.4	0.4
1969	1.5	0.5	0.0	0.0	0.0	0.8	0.4	0.9	0.4	3.6	4.4	11.7	14.0	16.6	21.6	24.7	28.3	24.7	3.7	3.3	0.4
1970	2.2	0.3	0.0	0.0	0.3	0.4	0.4	2.2	2.1	4.3	5.9	8.9	10.0	9.3	15.8	15.6	27.1	19.8	3.2	2.9	0.3
1971	1.9	0.0	0.0	0.3	0.0	0.0	1.3	0.9	2.2	2.9	8.8	6.9	20.2	11.5	17.5	15.3	36.8	24.6	3.7	3.4	0.4
1972	1.0	1.1	0.0	0.3	0.3	0.3	0.8	1.7	1.7	3.4	6.4	8.6	17.4	16.8	15.1	18.8	23.2	8.7	3.6	3.3	0.4
1973	2.3	1.1	0.0	0.0	0.9	0.0	0.0	1.7	3.4	5.5	6.5	12.6	15.6	12.4	17.7	21.1	10.2	25.1	3.9	3.7	0.4
1969-73	1.8	0.6	0.0	0.1	0.3	0.3	0.6	1.5	2.0	4.0	6.4	9.7	15.5	13.3	17.5	19.1	24.9	20.5	3.6	3.3	0.4
1974	2.9	0.9	0.3	0.0	0.0	0.6	1.1	0.0	2.6	4.2	8.3	9.1	13.2	21.9	24.8	18.0	29.9	13.2	4.2	3.8	0.4
1975	3.6	0.3	0.0	0.5	0.3	0.0	0.4	1.7	2.1	4.3	6.8	7.2	12.0	14.1	16.7	21.2	17.4	25.4	3.6	3.3	0.3
1976	2.4	0.9	0.0	0.0	0.3	0.0	0.0	2.5	0.9	3.9	6.7	10.0	14.2	13.5	13.5	24.2	13.1	24.2	3.5	3.2	0.3
1977	2.0	0.0	0.0	0.3	0.0	0.3	0.0	2.0	2.1	6.0	5.1	11.4	11.7	22.0	15.7	17.7	28.9	13.9	3.7	3.4	0.4
1978	2.0	0.0	0.0	0.0	0.5	0.3	0.3	1.1	2.6	4.7	5.5	9.8	15.1	11.3	20.2	24.0	42.0	13.4	3.8	3.4	0.4
1974-78	2.6	0.4	0.1	0.2	0.2	0.2	0.3	1.5	2.1	4.6	6.5	9.5	13.2	16.5	18.1	21.1	26.5	17.9	3.8	3.4	0.4
1979	1.4	1.3	0.0	0.2	0.0	0.5	0.6	2.9	3.0	3.4	9.8	9.3	12.7	16.1	33.4	18.8	18.5	31.8	4.3	3.9	0.5
1980	1.0	2.0	0.3	0.5	0.0	0.8	0.3	3.9	1.7	5.2	6.0	14.8	15.7	16.8	18.2	24.7	19.4	6.1	4.2	3.9	0.4
1981	0.7	1.0	0.3	0.3	0.0	0.0	0.8	2.1	2.0	4.3	8.1	10.7	9.1	18.9	29.9	22.8	18.8	15.5	4.0	3.6	0.4
1982	2.1	1.0	0.3	0.0	0.7	0.0	0.6	1.9	3.2	3.9	9.4	10.3	17.1	19.7	23.8	27.0	16.8	14.8	4.4	4.1	0.5
1983	2.7	1.0	0.0	0.0	0.5	0.0	0.8	0.9	2.7	5.2	7.7	10.2	10.0	22.2	27.9	31.8	26.4	28.1	4.5	4.0	0.5
1979-83	1.6	1.3	0.2	0.2	0.3	0.3	0.6	2.3	2.5	4.4	8.2	11.1	12.9	18.8	26.6	25.2	20.1	19.3	4.3	3.9	0.5

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF EYE (ICD9 190)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	6	0	0	0	0	0	0	4	1	2	2	5	2	4	2	4	0	0	0	32
1965	4	1	0	0	0	0	3	1	2	2	5	4	6	3	2	1	0	1	0	35
1966	9	1	0	0	0	0	0	1	2	1	1	0	8	1	5	1	0	1	0	31
1967	3	0	2	0	0	1	0	0	3	2	1	3	1	1	3	3	0	1	0	24
1968	1	1	0	0	1	2	2	1	0	4	3	4	6	3	2	1	1	0	0	32
1964-68	23	3	2	0	1	3	5	7	8	11	12	16	23	12	14	10	1	3	0	154
1969	7	2	0	0	1	2	0	0	3	1	1	4	5	5	2	2	3	1	0	39
1970	1	0	0	0	0	0	1	2	0	3	6	5	5	5	5	1	2	0	0	36
1971	3	1	0	0	2	1	0	0	1	2	4	2	7	2	2	3	1	0	0	31
1972	3	0	0	0	0	2	1	2	2	5	3	4	2	9	6	2	0	1	0	42
1973	5	1	0	0	0	0	2	1	3	1	5	3	2	0	4	5	2	2	1	37
1969-73	19	4	0	0	3	5	4	5	9	12	19	18	21	21	19	13	8	4	1	185
1974	4	1	0	0	1	1	2	1	3	4	4	0	3	4	0	4	2	1	0	35
1975	1	0	0	3	0	1	2	0	0	5	4	4	3	4	4	4	2	2	0	39
1976	5	1	0	1	1	3	1	0	1	4	1	2	4	2	3	4	1	3	0	37
1977	4	0	0	0	1	1	0	0	2	3	3	7	5	4	5	3	2	0	0	40
1978	3	1	0	2	1	1	0	0	5	6	2	4	6	4	5	5	1	0	2	48
1974-78	17	3	0	6	4	7	5	1	11	22	14	17	21	18	17	20	8	6	2	199
1979	6	0	1	0	0	0	0	3	2	0	2	3	7	5	5	4	2	1	0	41
1980	4	0	0	1	0	2	1	2	1	2	4	7	3	1	2	6	1	1	0	38
1981	7	0	0	3	0	0	1	1	0	3	4	0	8	9	0	1	4	0	1	42
1982	6	1	0	1	0	0	0	1	1	2	3	5	0	7	6	3	0	3	1	40
1983	4	0	0	0	1	0	1	2	3	2	7	5	1	3	4	5	2	0	1	41
1979-83	27	1	1	5	1	2	3	9	7	9	20	20	19	25	17	19	9	5	3	202

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF EYE (ICD9 190)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	1.5	0.0	0.0	0.0	0.0	0.0	0.0	1.7	0.5	1.1	1.2	3.6	1.8	4.4	2.9	8.8	0.0	0.0	1.0	0.9	0.1
1965	1.0	0.3	0.0	0.0	0.0	0.0	1.3	0.4	0.9	1.0	2.9	2.8	5.1	3.3	2.9	2.2	0.0	7.2	1.0	1.0	0.1
1966	2.4	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.9	0.5	0.6	0.0	6.7	1.1	7.1	2.1	0.0	7.1	0.9	0.9	0.1
1967	0.8	0.0	0.6	0.0	0.0	0.4	0.0	0.0	1.3	1.0	0.5	2.0	0.8	1.1	4.2	6.3	0.0	6.9	0.7	0.7	0.1
1968	0.3	0.2	0.0	0.0	0.4	0.8	0.9	0.4	0.0	1.9	1.6	2.5	4.8	3.1	2.8	2.1	3.7	0.0	0.9	0.9	0.1
1964-68	1.2	0.2	0.1	0.0	0.1	0.3	0.4	0.6	0.7	1.1	1.4	2.2	3.8	2.6	4.0	4.3	0.8	4.2	0.9	0.9	0.1
1969	2.0	0.5	0.0	0.0	0.3	0.8	0.0	0.0	1.2	0.5	0.5	2.5	3.9	5.0	2.8	4.2	11.0	6.4	1.1	1.1	0.1
1970	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.8	0.0	1.3	3.2	3.0	3.8	4.9	6.9	2.1	7.2	0.0	0.9	1.0	0.1
1971	0.9	0.2	0.0	0.0	0.6	0.3	0.0	0.0	0.4	0.9	2.1	1.2	5.1	1.9	2.7	6.1	3.5	0.0	0.8	0.8	0.1
1972	0.9	0.0	0.0	0.0	0.0	0.7	0.4	0.8	0.8	2.1	1.5	2.3	1.4	8.3	7.8	4.0	0.0	5.6	1.1	1.1	0.1
1973	1.5	0.3	0.0	0.0	0.0	0.0	0.8	0.4	1.2	0.4	2.4	1.8	1.4	0.0	5.0	10.0	6.9	11.0	0.9	0.9	0.1
1969-73	1.1	0.2	0.0	0.0	0.2	0.3	0.3	0.4	0.7	1.0	2.0	2.1	3.1	4.0	5.1	5.3	5.7	4.7	1.0	1.0	0.1
1974	1.2	0.3	0.0	0.0	0.3	0.3	0.7	0.4	1.2	1.7	1.8	0.0	2.0	3.5	0.0	7.9	6.8	5.4	0.9	0.9	0.1
1975	0.3	0.0	0.0	0.7	0.0	0.3	0.7	0.0	0.0	2.1	1.8	2.3	1.9	3.4	4.7	7.6	6.8	10.5	0.9	0.9	0.1
1976	1.6	0.3	0.0	0.2	0.3	0.8	0.3	0.0	0.4	1.7	0.4	1.1	2.5	1.7	3.5	7.4	3.4	15.5	0.9	0.9	0.1
1977	1.3	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.8	1.2	1.3	3.7	3.2	3.2	5.7	5.3	6.6	0.0	0.9	1.0	0.1
1978	1.0	0.3	0.0	0.5	0.3	0.3	0.0	0.0	2.1	2.5	0.9	2.0	3.8	3.1	5.5	8.5	3.2	0.0	1.1	1.1	0.1
1974-78	1.1	0.2	0.0	0.3	0.2	0.4	0.3	0.1	0.9	1.8	1.2	1.9	2.7	3.0	3.9	7.3	5.4	6.2	0.9	1.0	0.1
1979	1.9	0.0	0.3	0.0	0.0	0.0	0.0	1.1	0.8	0.0	0.9	1.4	4.5	3.7	5.3	6.5	6.3	4.9	0.9	1.0	0.1
1980	1.3	0.0	0.0	0.2	0.0	0.6	0.3	0.7	0.4	0.8	1.7	3.3	1.9	0.7	2.1	9.5	3.0	4.8	0.8	0.8	0.1
1981	2.3	0.0	0.0	0.7	0.0	0.0	0.3	0.3	0.0	1.3	1.7	0.0	4.7	6.4	0.0	1.5	11.7	0.0	0.9	1.0	0.1
1982	2.0	0.3	0.0	0.2	0.0	0.0	0.0	0.3	0.4	0.8	1.3	2.3	0.0	4.9	5.7	4.5	0.0	13.9	0.8	0.9	0.1
1983	1.3	0.0	0.0	0.0	0.2	0.0	0.3	0.6	1.1	0.8	3.0	2.3	0.5	2.1	3.7	7.3	5.3	0.0	0.8	0.8	0.1
1979-83	1.8	0.1	0.1	0.2	0.1	0.1	0.2	0.6	0.6	0.8	1.7	1.9	2.2	3.6	3.4	5.9	5.2	4.7	0.9	0.9	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF EYE (ICD9 190)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	3	1	0	0	0	0	0	1	4	2	5	3	2	0	1	1	0	0	0	23
1965	5	1	0	0	0	1	0	1	4	3	2	0	1	2	5	2	1	1	0	29
1966	5	0	0	0	0	0	2	2	2	1	0	2	4	3	2	1	0	0	0	24
1967	4	0	1	0	0	0	0	1	1	1	3	0	2	3	1	2	1	1	0	21
1968	3	0	1	0	0	1	0	2	3	1	2	3	4	4	3	2	0	1	0	30
1964-68	20	2	2	0	0	2	2	7	14	8	12	8	13	12	12	8	2	3	0	127
1969	3	0	1	1	1	1	1	1	2	1	2	2	4	5	2	3	1	0	0	31
1970	1	0	0	1	0	1	0	0	2	1	2	2	2	1	2	1	0	1	0	17
1971	2	1	0	2	1	2	0	2	0	1	2	1	5	3	1	1	2	0	0	26
1972	3	0	0	0	0	1	0	2	0	3	5	7	2	2	1	1	0	3	0	30
1973	5	0	2	2	1	2	1	1	3	1	0	6	4	1	3	3	3	0	0	38
1969-73	14	1	3	6	3	7	2	6	7	7	11	18	17	12	9	9	6	4	0	142
1974	5	1	0	2	2	2	2	2	4	1	2	2	1	3	3	2	0	2	0	36
1975	0	0	0	1	2	1	2	0	1	8	1	3	3	3	1	2	1	2	0	31
1976	4	0	0	0	2	4	2	3	2	4	3	4	4	2	2	0	1	1	0	36
1977	4	0	1	0	0	0	2	0	3	7	5	1	1	0	5	1	3	0	0	33
1978	2	0	0	0	1	0	3	1	2	1	2	4	3	2	5	2	2	3	0	33
1974-78	15	1	1	3	7	7	11	6	12	21	13	14	12	10	16	7	7	8	0	171
1979	6	0	0	0	0	2	1	0	0	2	2	3	6	4	9	3	3	2	0	43
1980	5	1	0	0	0	2	1	4	0	1	5	4	4	5	4	1	0	2	0	39
1981	2	0	0	0	2	0	0	1	0	1	5	6	2	4	3	3	5	2	0	36
1982	1	0	0	0	1	1	0	1	2	4	5	4	7	7	8	3	1	2	0	47
1983	3	0	0	0	0	0	0	2	1	1	6	8	6	2	4	4	4	4	0	45
1979-83	17	1	0	0	3	5	2	8	3	9	23	25	25	22	28	14	13	12	0	210

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF EYE (ICD9 190)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.8	0.3	0.0	0.0	0.0	0.0	0.0	0.4	1.8	1.1	3.0	2.2	1.7	0.0	1.2	1.7	0.0	0.0	0.7	0.7	0.1
1965	1.3	0.3	0.0	0.0	0.0	0.5	0.0	0.4	1.7	1.6	1.2	0.0	0.8	1.9	5.8	3.3	2.8	4.3	0.8	0.8	0.1
1966	1.4	0.0	0.0	0.0	0.0	0.0	0.9	0.9	0.8	0.5	0.0	1.4	3.2	2.8	2.3	1.6	0.0	0.0	0.7	0.7	0.1
1967	1.1	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.4	0.5	1.7	0.0	1.6	2.8	1.1	3.1	2.6	4.0	0.6	0.6	0.0
1968	0.9	0.0	0.3	0.0	0.0	0.4	0.0	0.9	1.2	0.5	1.1	1.9	3.0	3.6	3.3	3.0	0.0	3.8	0.8	0.8	0.1
1964-68	1.1	0.1	0.1	0.0	0.0	0.2	0.2	0.6	1.2	0.8	1.4	1.1	2.1	2.3	2.7	2.6	1.1	2.5	0.7	0.7	0.1
1969	0.9	0.0	0.3	0.3	0.3	0.4	0.4	0.4	0.8	0.5	1.1	1.2	2.9	4.4	2.2	4.4	2.4	0.0	0.8	0.8	0.1
1970	0.3	0.0	0.0	0.3	0.0	0.4	0.0	0.0	0.9	0.4	1.1	1.2	1.4	0.8	2.1	1.4	0.0	3.3	0.4	0.4	0.0
1971	0.6	0.3	0.0	0.6	0.3	0.7	0.0	0.9	0.0	0.4	1.0	0.6	3.5	2.5	1.0	1.4	4.3	0.0	0.7	0.7	0.1
1972	1.0	0.0	0.0	0.0	0.0	0.3	0.0	0.9	0.0	1.3	2.5	4.0	1.3	1.6	1.0	1.3	0.0	8.7	0.7	0.7	0.1
1973	1.6	0.0	0.5	0.5	0.3	0.6	0.4	0.4	1.3	0.4	0.0	3.4	2.6	0.8	2.9	3.9	6.1	0.0	0.9	0.9	0.1
1969-73	0.9	0.1	0.2	0.3	0.2	0.5	0.2	0.5	0.6	0.6	1.1	2.1	2.4	2.0	1.9	2.5	2.6	2.5	0.7	0.7	0.1
1974	1.6	0.3	0.0	0.5	0.6	0.6	0.7	0.9	1.7	0.4	0.9	1.1	0.6	2.3	2.9	2.6	0.0	5.3	0.9	0.9	0.1
1975	0.0	0.0	0.0	0.3	0.5	0.3	0.7	0.0	0.4	3.4	0.4	1.7	1.8	2.2	0.9	2.5	1.9	5.1	0.7	0.7	0.1
1976	1.4	0.0	0.0	0.0	0.5	1.1	0.7	1.2	0.9	1.7	1.3	2.1	2.4	1.4	1.8	0.0	1.9	2.4	0.9	0.9	0.1
1977	1.4	0.0	0.3	0.0	0.0	0.0	0.6	0.0	1.3	3.0	2.1	0.5	0.6	0.0	4.4	1.2	5.4	0.0	0.7	0.8	0.1
1978	0.7	0.0	0.0	0.0	0.3	0.0	0.9	0.4	0.9	0.4	0.8	1.9	1.7	1.3	4.2	2.3	3.5	6.7	0.7	0.6	0.1
1974-78	1.0	0.1	0.1	0.2	0.4	0.4	0.7	0.5	1.0	1.8	1.1	1.5	1.4	1.4	2.9	1.7	2.6	3.9	0.8	0.8	0.1
1979	2.0	0.0	0.0	0.0	0.0	0.5	0.3	0.0	0.0	0.9	0.9	1.3	3.5	2.6	7.3	3.3	5.0	4.2	0.9	0.9	0.1
1980	1.7	0.3	0.0	0.0	0.0	0.5	0.3	1.4	0.0	0.4	2.1	1.7	2.2	3.1	3.2	1.1	0.0	4.1	0.8	0.9	0.1
1981	0.7	0.0	0.0	0.0	0.5	0.0	0.0	0.3	0.0	0.4	2.1	2.6	1.1	2.4	2.3	3.1	7.8	3.9	0.7	0.6	0.1
1982	0.3	0.0	0.0	0.0	0.2	0.3	0.0	0.3	0.8	1.7	2.1	1.7	3.5	4.2	6.0	3.0	1.5	3.7	0.9	0.9	0.1
1983	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.4	0.4	2.6	3.4	2.9	1.2	2.9	3.9	5.9	7.0	0.9	0.8	0.1
1979-83	1.2	0.1	0.0	0.0	0.2	0.3	0.1	0.5	0.2	0.8	2.0	2.2	2.6	2.7	4.3	2.9	4.1	4.6	0.8	0.8	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF BRAIN (ICD9 191)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	14	8	5	4	10	7	5	8	11	17	28	27	26	16	6	7	1	1	0	201
1965	9	12	9	2	4	6	15	11	13	19	12	16	20	10	7	1	2	0	0	168
1966	8	11	9	5	5	6	6	12	12	17	20	21	22	14	8	6	1	0	0	183
1967	11	8	8	6	6	2	6	14	13	18	16	26	18	15	5	3	3	1	0	179
1968	12	12	6	8	2	11	10	13	16	16	23	23	19	14	8	4	2	1	0	200
1964-68	54	51	37	25	27	32	42	58	65	87	99	113	105	69	34	21	9	3	0	931
1969	14	10	6	5	3	4	14	11	17	22	30	26	24	16	4	2	0	0	0	208
1970	11	9	6	9	4	6	6	12	11	24	22	14	24	13	5	3	1	0	0	180
1971	8	13	8	12	6	5	8	13	20	29	28	25	27	25	14	3	0	1	0	245
1972	9	6	12	10	6	9	5	10	18	16	27	30	35	19	24	7	1	1	0	245
1973	8	7	9	6	5	4	9	6	20	26	19	27	21	19	17	7	0	1	0	211
1969-73	50	45	41	42	24	28	42	52	86	117	126	122	131	92	64	22	2	3	0	1089
1974	5	4	10	5	6	7	11	9	14	20	30	29	31	28	17	12	4	0	1	243
1975	7	13	14	5	12	5	6	14	17	20	30	26	26	20	18	6	5	1	0	245
1976	6	9	14	7	10	12	6	2	16	24	32	28	37	28	13	11	5	1	0	261
1977	14	12	9	4	8	15	11	12	7	25	21	30	25	30	17	12	8	1	0	261
1978	6	5	9	11	11	10	11	7	22	28	32	22	42	34	23	10	5	0	0	288
1974-78	38	43	56	32	47	49	45	44	76	117	145	135	161	140	88	51	27	3	1	1298
1979	8	14	6	13	11	13	27	13	12	22	38	33	28	35	15	10	2	3	0	303
1980	8	5	4	7	11	10	14	17	13	19	40	38	25	44	24	12	3	1	2	297
1981	11	6	10	7	9	12	18	9	9	23	32	45	36	43	34	12	7	1	0	324
1982	11	2	7	10	10	9	17	15	13	21	24	50	48	40	26	20	5	3	0	331
1983	5	22	7	15	9	13	13	15	21	24	26	38	48	33	33	28	13	2	0	365
1979-83	43	49	34	52	50	57	89	69	68	109	160	204	185	195	132	82	30	10	2	1620

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF BRAIN (ICD9 191)

YEAR	AGE SPECIFIC RATES																	SUMMARY RATES		
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	3.6	2.2	1.5	1.5	4.7	3.4	2.2	3.4	5.0	9.0	16.5	19.3	22.8	17.8	8.8	15.4	3.9	7.4	6.0	6.2	0.6
1965	2.3	3.1	2.6	0.7	1.8	2.9	6.7	4.6	5.8	10.0	6.9	11.2	17.1	10.9	10.1	2.2	7.7	0.0	4.9	5.0	0.5
1966	2.1	2.8	2.6	1.6	2.1	2.8	2.6	5.0	5.2	8.7	11.2	14.3	18.4	15.0	11.4	12.9	3.8	0.0	5.2	5.3	0.5
1967	2.9	2.0	2.2	1.9	2.3	0.9	2.6	5.8	5.5	8.9	8.8	17.1	14.6	15.8	7.1	6.3	11.4	6.9	5.0	5.0	0.5
1968	3.3	3.0	1.6	2.4	0.7	4.6	4.4	5.4	6.6	7.7	12.6	14.6	15.1	14.4	11.3	8.4	7.5	6.7	5.5	5.6	0.5
1964-68	2.8	2.6	2.1	1.6	2.2	2.9	3.7	4.9	5.6	8.8	11.1	15.3	17.5	14.8	9.7	9.0	6.9	4.2	5.3	5.4	0.5
1969	4.1	2.5	1.6	1.5	1.0	1.6	6.1	4.6	7.0	10.1	16.4	16.0	18.6	16.0	5.6	4.2	0.0	0.0	5.6	5.7	0.6
1970	3.3	2.2	1.5	2.6	1.3	2.2	2.5	5.0	4.6	10.6	11.9	8.4	18.1	12.6	6.9	6.2	3.6	0.0	4.7	4.9	0.5
1971	2.5	3.2	2.0	3.3	1.8	1.7	3.3	5.4	8.3	12.5	14.8	14.7	19.8	23.6	18.8	6.1	0.0	5.8	6.3	6.4	0.7
1972	2.8	1.5	2.9	2.7	1.8	3.0	2.0	4.2	7.4	6.9	13.6	17.6	24.8	17.6	31.3	14.2	3.5	5.6	6.2	6.3	0.7
1973	2.5	1.9	2.2	1.6	1.5	1.3	3.5	2.5	8.2	11.1	9.1	15.9	14.4	17.3	21.4	14.1	0.0	5.5	5.3	5.3	0.6
1969-73	3.0	2.3	2.0	2.3	1.5	2.0	3.4	4.4	7.1	10.2	13.1	14.5	19.2	17.5	17.0	9.0	1.4	3.5	5.6	5.7	0.6
1974	1.5	1.1	2.4	1.3	1.7	2.1	4.0	3.7	5.7	8.5	13.7	17.2	20.6	24.8	20.7	23.7	13.7	0.0	5.9	5.9	0.6
1975	2.2	3.8	3.3	1.2	3.3	1.4	2.1	5.7	7.0	8.4	13.4	15.1	16.8	17.2	21.3	11.5	17.1	5.3	5.9	5.9	0.6
1976	1.9	2.6	3.4	1.7	2.7	3.4	2.0	0.8	6.6	10.0	14.1	15.6	23.5	23.2	15.0	20.2	16.8	5.2	6.2	6.2	0.6
1977	4.5	3.6	2.3	1.0	2.1	4.3	3.5	4.7	2.9	10.4	9.2	15.9	15.8	23.8	19.3	21.2	26.5	5.1	6.1	6.2	0.6
1978	1.9	1.5	2.4	2.6	2.9	2.8	3.3	2.6	9.2	11.6	14.0	11.1	26.7	26.1	25.4	17.0	16.2	0.0	6.6	6.6	0.7
1974-78	2.4	2.5	2.8	1.6	2.6	2.8	3.0	3.5	6.3	9.8	12.9	14.9	20.7	23.1	20.4	18.7	18.1	3.1	6.1	6.2	0.6
1979	2.6	4.2	1.6	3.1	2.8	3.7	7.9	4.7	5.0	9.1	16.5	15.8	17.9	26.0	16.0	16.3	6.3	14.6	6.8	6.8	0.7
1980	2.6	1.5	1.1	1.7	2.8	2.8	4.0	6.0	5.3	8.0	17.2	17.9	15.5	31.7	24.7	19.1	9.1	4.8	6.4	6.5	0.7
1981	3.6	1.9	2.9	1.7	2.3	3.3	5.0	3.1	3.6	9.7	13.6	20.9	21.3	30.5	33.8	18.5	20.5	4.7	7.0	7.0	0.8
1982	3.6	0.6	2.0	2.5	2.5	2.5	4.8	4.8	5.1	8.9	10.1	22.9	27.0	28.0	24.9	30.1	14.0	13.9	7.0	7.0	0.7
1983	1.6	7.1	2.1	3.9	2.2	3.5	3.7	4.6	8.0	10.2	11.0	17.2	25.6	23.2	30.2	40.9	34.4	9.0	7.7	7.7	0.8
1979-83	2.8	3.1	1.9	2.5	2.5	3.1	5.1	4.6	5.4	9.2	13.6	19.0	21.7	27.9	26.1	25.3	17.4	9.4	7.0	7.0	0.7

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF BRAIN (ICD9 191)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	8	9	4	5	5	5	6	6	6	8	12	13	14	9	8	3	1	0	0	122
1965	5	12	6	3	3	2	5	2	6	14	12	15	12	9	4	2	4	0	0	116
1966	9	10	6	4	2	3	1	6	9	9	15	8	14	10	7	3	2	0	0	118
1967	9	7	2	5	1	1	7	7	18	10	12	11	12	10	5	3	0	0	0	120
1968	8	7	7	1	1	6	5	8	8	13	23	5	16	14	7	4	1	0	0	134
1964-68	39	45	25	18	12	17	24	29	47	54	74	52	68	52	31	15	8	0	0	610
1969	7	5	7	4	7	4	4	3	8	13	12	11	23	18	5	0	2	0	0	133
1970	9	12	6	0	2	3	2	11	10	11	10	20	11	9	12	6	2	0	0	136
1971	6	7	5	4	7	3	5	8	12	26	10	22	11	13	16	3	1	0	0	159
1972	4	7	7	4	3	2	6	6	8	13	15	23	14	13	9	5	1	1	0	141
1973	8	12	9	4	5	7	7	8	5	8	16	20	29	18	19	8	4	1	0	188
1969-73	34	43	34	16	24	19	24	36	43	71	63	96	88	71	61	22	10	2	0	757
1974	8	11	9	2	3	5	8	6	9	11	20	14	30	22	11	9	3	1	0	182
1975	8	8	7	6	5	8	5	3	9	19	13	23	26	18	8	4	4	0	0	174
1976	8	8	6	4	5	10	10	8	8	21	30	22	20	27	16	10	4	1	0	218
1977	5	12	9	4	4	12	10	11	7	16	18	23	22	22	18	8	3	4	0	208
1978	4	8	8	6	2	10	10	9	6	14	11	24	36	28	19	19	8	5	1	228
1974-78	33	47	39	22	19	45	43	37	39	81	92	106	134	117	72	50	22	11	1	1010
1979	4	7	3	5	10	13	4	13	13	14	21	23	26	31	23	15	7	5	0	237
1980	9	6	7	7	6	14	12	9	5	17	24	33	30	28	21	15	8	2	0	253
1981	10	4	3	1	6	8	12	10	14	21	20	35	29	28	32	16	4	6	0	258
1982	11	7	4	4	3	11	14	11	14	7	21	36	30	14	28	13	13	7	0	248
1983	8	10	15	5	6	8	9	10	16	9	17	35	37	24	23	13	14	2	0	261
1979-83	42	34	32	22	31	54	51	53	62	68	103	162	151	125	127	72	46	22	0	1257

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF BRAIN (ICD9 191)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	2.1	2.5	1.3	1.9	2.3	2.4	2.7	2.6	2.7	4.3	7.3	9.4	11.9	8.9	9.6	5.1	2.9	0.0	3.7	3.7	0.4
1965	1.3	3.3	1.8	1.1	1.3	0.9	2.3	0.9	2.6	7.4	7.1	10.6	9.9	8.7	4.7	3.3	11.1	0.0	3.5	3.3	0.3
1966	2.5	2.7	1.8	1.4	0.8	1.4	0.5	2.6	3.8	4.6	8.6	5.5	11.2	9.4	7.9	4.8	5.4	0.0	3.4	3.3	0.3
1967	2.5	1.8	0.6	1.6	0.4	0.4	3.1	3.0	7.5	4.9	6.7	7.3	9.3	9.2	5.5	4.6	0.0	0.0	3.3	3.3	0.3
1968	2.3	1.8	2.0	0.3	0.4	2.5	2.2	3.5	3.3	6.1	12.7	3.2	12.1	12.6	7.7	6.0	2.5	0.0	3.6	3.7	0.4
1964-68	2.2	2.4	1.5	1.2	1.0	1.5	2.2	2.5	4.0	5.5	8.5	7.1	10.9	9.8	7.1	4.8	4.3	0.0	3.5	3.4	0.3
1969	2.1	1.3	1.9	1.2	2.4	1.6	1.8	1.3	3.4	5.9	6.5	6.8	16.9	15.7	5.4	0.0	4.7	0.0	3.6	3.6	0.4
1970	2.8	3.1	1.6	0.0	0.6	1.1	0.9	4.8	4.3	4.8	5.4	11.9	7.9	7.6	12.7	8.5	4.5	0.0	3.5	3.5	0.3
1971	1.9	1.8	1.3	1.1	2.1	1.1	2.1	3.5	5.2	11.0	5.2	12.7	7.6	10.7	16.5	4.2	2.2	0.0	4.0	4.0	0.4
1972	1.3	1.9	1.8	1.1	0.9	0.7	2.5	2.6	3.4	5.5	7.4	13.2	9.4	10.4	9.1	6.7	2.1	2.9	3.5	3.4	0.4
1973	2.6	3.4	2.3	1.1	1.5	2.2	2.7	3.5	2.2	3.4	7.4	11.4	18.8	14.0	18.7	10.5	8.2	2.8	4.6	4.5	0.5
1969-73	2.2	2.3	1.8	0.9	1.5	1.3	2.0	3.1	3.7	6.1	6.4	11.2	12.2	11.7	12.6	6.1	4.4	1.2	3.9	3.8	0.4
1974	2.6	3.3	2.3	0.5	0.8	1.5	3.0	2.6	3.9	4.7	8.8	7.9	18.8	16.6	10.5	11.6	6.0	2.6	4.4	4.3	0.4
1975	2.6	2.4	1.8	1.6	1.4	2.3	1.8	1.3	3.9	8.1	5.6	12.7	15.7	13.3	7.4	5.0	7.8	0.0	4.1	4.1	0.4
1976	2.7	2.5	1.5	1.0	1.3	2.8	3.4	3.3	3.4	9.0	12.6	11.6	11.8	19.2	14.4	12.1	7.5	2.4	5.0	4.9	0.5
1977	1.7	3.7	2.4	1.0	1.0	3.3	3.2	4.4	3.0	6.9	7.6	11.4	12.9	15.1	15.7	9.4	5.4	9.2	4.6	4.6	0.5
1978	1.4	2.5	2.2	1.5	0.5	2.8	3.0	3.4	2.6	6.0	4.7	11.2	20.9	18.6	16.0	21.7	14.0	11.1	4.9	4.7	0.5
1974-78	2.2	2.9	2.0	1.1	1.0	2.5	2.9	3.0	3.3	6.9	7.9	11.0	16.0	16.6	12.9	12.1	8.2	5.3	4.6	4.5	0.5
1979	1.4	2.2	0.9	1.2	2.6	3.5	1.2	4.8	5.5	6.0	9.0	10.2	15.0	20.0	18.7	16.6	11.7	10.6	5.0	4.8	0.5
1980	3.1	2.0	2.1	1.7	1.5	3.7	3.4	3.2	2.1	7.3	10.3	14.3	16.8	17.4	16.6	16.1	12.9	4.1	5.3	5.2	0.5
1981	3.5	1.3	0.9	0.3	1.5	2.1	3.3	3.5	5.7	9.1	8.6	15.0	14.9	17.0	24.6	16.6	6.3	11.7	5.2	5.1	0.6
1982	3.8	2.4	1.2	1.0	0.7	2.9	3.9	3.5	5.6	3.0	9.0	15.4	15.1	8.4	20.8	13.0	19.9	13.0	4.9	4.8	0.5
1983	2.7	3.4	4.7	1.4	1.5	2.1	2.5	3.0	6.1	3.9	7.3	14.9	17.6	14.4	16.5	12.5	20.6	3.5	5.3	5.1	0.5
1979-83	2.9	2.3	1.9	1.1	1.6	2.9	2.9	3.6	5.0	5.9	8.8	14.0	15.9	15.3	19.4	14.9	14.4	8.5	5.2	5.0	0.5

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF NERVOUS SYSTEM (ICD9 192)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	2	1	2	1	0	1	1	4	2	0	4	3	1	4	1	0	0	0	0	27
1965	3	0	2	1	0	0	0	1	1	1	4	0	2	0	1	0	0	0	0	16
1966	2	0	1	1	3	1	2	1	2	1	1	1	1	2	0	0	0	0	0	19
1967	1	0	1	3	1	1	0	1	1	2	2	1	1	0	0	1	0	0	0	16
1968	1	2	1	1	3	1	0	3	1	2	1	0	1	2	0	1	0	0	0	20
1964-68	9	3	7	7	7	4	3	10	7	6	12	5	6	8	2	2	0	0	0	98
1969	3	0	3	1	0	0	2	0	1	4	2	0	1	0	0	2	0	0	0	19
1970	2	3	0	0	1	1	3	0	2	0	2	2	3	4	1	0	0	0	0	24
1971	1	1	2	1	0	0	1	3	0	2	0	1	0	1	1	1	0	0	0	15
1972	1	0	3	0	1	0	0	0	2	0	2	0	2	0	0	1	1	1	0	14
1973	2	2	1	1	2	2	1	0	2	1	1	0	2	3	3	1	0	0	0	24
1969-73	9	6	9	3	4	3	7	3	7	7	7	3	8	8	5	5	1	1	0	96
1974	2	1	1	0	2	1	3	1	0	4	6	3	3	3	0	5	1	0	1	37
1975	4	1	0	2	2	0	0	4	1	0	5	2	2	1	6	1	1	1	0	33
1976	4	1	1	2	5	1	0	4	2	6	3	4	1	3	1	1	0	0	0	39
1977	1	3	0	5	3	0	0	2	3	6	1	4	2	3	2	1	0	0	0	36
1978	3	0	2	2	3	0	4	4	1	4	3	4	8	7	3	0	0	0	0	48
1974-78	14	6	4	11	15	2	7	15	7	20	18	17	16	17	12	8	2	1	1	193
1979	3	0	1	0	3	3	5	1	0	5	0	2	3	4	3	2	1	1	0	37
1980	0	2	1	2	4	3	3	0	2	1	0	1	4	4	4	2	1	0	0	34
1981	1	0	1	2	2	4	0	4	0	3	2	1	4	3	2	0	1	0	0	30
1982	2	0	1	0	2	4	2	1	0	5	3	1	2	3	0	3	0	0	0	29
1983	3	3	0	1	0	1	0	0	1	1	1	2	4	1	6	0	1	0	0	25
1979-83	9	5	4	5	11	15	10	6	3	15	6	7	17	15	15	7	4	1	0	155

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF NERVOUS SYSTEM (ICD9 192)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.5	0.3	0.6	0.4	0.0	0.5	0.4	1.7	0.9	0.0	2.4	2.1	0.9	4.4	1.5	0.0	0.0	0.0	0.8	0.8	0.1
1965	0.8	0.0	0.6	0.3	0.0	0.0	0.0	0.4	0.4	0.5	2.3	0.0	1.7	0.0	1.4	0.0	0.0	0.0	0.5	0.5	0.0
1966	0.5	0.0	0.3	0.3	1.2	0.5	0.9	0.4	0.9	0.5	0.6	0.7	0.8	2.1	0.0	0.0	0.0	0.0	0.5	0.6	0.0
1967	0.3	0.0	0.3	0.9	0.4	0.4	0.0	0.4	0.4	1.0	1.1	0.7	0.8	0.0	0.0	2.1	0.0	0.0	0.4	0.5	0.0
1968	0.3	0.5	0.3	0.3	1.1	0.4	0.0	1.3	0.4	1.0	0.5	0.0	0.8	2.1	0.0	2.1	0.0	0.0	0.5	0.6	0.0
1964-68	0.5	0.2	0.4	0.5	0.6	0.4	0.3	0.8	0.6	0.6	1.4	0.7	1.0	1.7	0.6	0.9	0.0	0.0	0.6	0.6	0.1
1969	0.9	0.0	0.8	0.3	0.0	0.0	0.9	0.0	0.4	1.8	1.1	0.0	0.8	0.0	0.0	4.2	0.0	0.0	0.5	0.5	0.0
1970	0.6	0.7	0.0	0.0	0.3	0.4	1.3	0.0	0.8	0.0	1.1	1.2	2.3	3.9	1.4	0.0	0.0	0.0	0.6	0.7	0.1
1971	0.3	0.2	0.5	0.3	0.0	0.0	0.4	1.3	0.0	0.9	0.0	0.6	0.0	0.9	1.3	2.0	0.0	0.0	0.4	0.4	0.0
1972	0.3	0.0	0.7	0.0	0.3	0.0	0.0	0.0	0.8	0.0	1.0	0.0	1.4	0.0	0.0	2.0	3.5	5.6	0.4	0.3	0.0
1973	0.6	0.5	0.2	0.3	0.6	0.6	0.4	0.0	0.8	0.4	0.5	0.0	1.4	2.7	3.8	2.0	0.0	0.0	0.6	0.6	0.1
1969-73	0.5	0.3	0.4	0.2	0.2	0.2	0.6	0.3	0.6	0.6	0.7	0.4	1.2	1.5	1.3	2.0	0.7	1.2	0.5	0.5	0.0
1974	0.6	0.3	0.2	0.0	0.6	0.3	1.1	0.4	0.0	1.7	2.7	1.8	2.0	2.7	0.0	9.9	3.4	0.0	0.9	0.9	0.1
1975	1.2	0.3	0.0	0.5	0.6	0.0	0.0	1.6	0.4	0.0	2.2	1.2	1.3	0.9	7.1	1.9	3.4	5.3	0.8	0.8	0.1
1976	1.3	0.3	0.2	0.5	1.4	0.3	0.0	1.6	0.8	2.5	1.3	2.2	0.6	2.5	1.2	1.8	0.0	0.0	0.9	1.0	0.1
1977	0.3	0.9	0.0	1.2	0.8	0.0	0.0	0.8	1.3	2.5	0.4	2.1	1.3	2.4	2.3	1.8	0.0	0.0	0.9	0.9	0.1
1978	1.0	0.0	0.5	0.5	0.8	0.0	1.2	1.5	0.4	1.7	1.3	2.0	5.1	5.4	3.3	0.0	0.0	0.0	1.1	1.1	0.1
1974-78	0.9	0.3	0.2	0.5	0.8	0.1	0.5	1.2	0.6	1.7	1.6	1.9	2.1	2.8	2.8	2.9	1.3	1.0	0.9	0.9	0.1
1979	1.0	0.0	0.3	0.0	0.8	0.8	1.5	0.4	0.0	2.1	0.0	1.0	1.9	3.0	3.2	3.3	3.1	4.9	0.8	0.8	0.1
1980	0.0	0.6	0.3	0.5	1.0	0.8	0.9	0.0	0.8	0.4	0.0	0.5	2.5	2.9	4.1	3.2	3.0	0.0	0.8	0.7	0.1
1981	0.3	0.0	0.3	0.5	0.5	1.1	0.0	1.4	0.0	1.3	0.8	0.5	2.4	2.1	2.0	0.0	2.9	0.0	0.7	0.7	0.1
1982	0.7	0.0	0.3	0.0	0.5	1.1	0.6	0.3	0.0	2.1	1.3	0.5	1.1	2.1	0.0	4.5	0.0	0.0	0.6	0.6	0.1
1983	1.0	1.0	0.0	0.3	0.0	0.3	0.0	0.0	0.4	0.4	0.4	0.9	2.1	0.7	5.5	0.0	2.6	0.0	0.5	0.6	0.1
1979-83	0.6	0.3	0.2	0.2	0.6	0.8	0.6	0.4	0.2	1.3	0.5	0.7	2.0	2.1	3.0	2.2	2.3	0.9	0.7	0.7	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF NERVOUS SYSTEM (ICD9 192)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	0	0	2	0	0	3	2	1	2	1	2	1	0	2	0	0	0	0	17
1965	2	4	1	2	1	0	3	2	0	0	1	2	3	1	1	0	1	0	0	24
1966	1	1	0	2	0	1	0	0	4	1	2	1	0	0	1	1	0	0	0	15
1967	1	2	1	1	1	0	2	3	4	2	0	0	0	1	0	0	1	0	0	19
1968	1	1	1	0	0	0	0	0	0	3	2	2	0	3	0	0	0	0	0	13
1964-68	6	8	3	7	2	1	8	7	9	8	6	7	4	5	4	1	2	0	0	88
1969	0	3	2	2	1	0	0	0	1	2	2	0	3	3	2	0	0	0	0	21
1970	3	0	6	1	0	0	2	0	0	0	3	3	2	1	1	0	0	1	0	23
1971	3	1	0	0	0	0	0	1	0	1	1	1	3	0	0	1	2	0	0	14
1972	2	2	1	0	1	1	0	0	2	2	0	4	2	4	3	1	0	0	0	25
1973	6	1	1	0	1	0	0	1	1	1	5	1	4	2	6	0	0	0	0	30
1969-73	14	7	10	3	3	1	2	2	4	6	11	9	14	10	12	2	2	1	0	113
1974	2	1	2	1	1	3	1	2	1	1	0	1	4	2	4	1	0	0	0	27
1975	0	2	0	1	1	3	0	0	1	1	1	0	4	4	4	1	1	0	1	25
1976	1	1	2	0	1	2	1	1	1	3	2	3	5	2	1	1	4	1	0	32
1977	3	1	1	1	2	3	3	3	0	4	5	4	2	3	0	2	3	1	0	41
1978	1	0	0	2	2	1	2	1	0	2	3	3	6	1	4	3	1	1	0	33
1974-78	7	5	5	5	7	12	7	7	3	11	11	11	21	12	13	8	9	3	1	158
1979	1	0	0	0	1	3	2	1	1	3	3	4	5	2	4	6	1	1	0	38
1980	1	0	0	0	2	2	3	2	0	2	3	2	7	4	3	7	1	1	0	40
1981	2	0	1	0	1	2	1	2	1	0	4	3	4	0	5	3	2	0	0	31
1982	1	0	0	2	1	3	0	2	2	3	1	4	1	5	3	2	2	0	0	32
1983	2	1	2	2	0	0	1	2	2	2	1	4	3	0	0	2	1	1	0	26
1979-83	7	1	3	4	5	10	7	9	6	10	12	17	20	11	15	20	7	3	0	167

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED PARTS OF NERVOUS SYSTEM (ICD9 192)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.0	0.0	0.8	0.0	0.0	1.4	0.9	0.5	1.1	0.6	1.4	0.8	0.0	2.4	0.0	0.0	0.0	0.5	0.5	0.1
1965	0.5	1.1	0.3	0.7	0.4	0.0	1.4	0.9	0.0	0.0	0.6	1.4	2.5	1.0	1.2	0.0	2.8	0.0	0.7	0.7	0.1
1966	0.3	0.3	0.0	0.7	0.0	0.5	0.0	0.0	1.7	0.5	1.1	0.7	0.0	0.0	1.1	6	0.0	0.0	0.4	0.4	0.0
1967	0.3	0.5	0.3	0.3	0.4	0.0	0.9	1.3	1.7	1.0	0.0	0.0	0.0	0.9	0.0	9	2.6	0.0	0.5	0.5	0.0
1968	0.3	0.3	0.3	0.0	0.0	0.0	0.0	0.0	0.0	1.4	1.1	1.3	0.0	2.7	0.0	.0	0.0	0.0	0.4	0.4	0.0
1964-68	0.3	0.4	0.2	0.5	0.2	0.1	0.7	0.6	0.8	0.8	0.7	1.0	0.6	0.9	0.9	0.3	1.1	0.0	0.5	0.5	0.0
1969	0.0	0.8	0.5	0.6	0.3	0.0	0.0	0.0	0.4	0.9	1.1	0.0	2.2	2.6	2.2	0.0	0.0	0.0	0.6	0.6	0.1
1970	0.9	0.0	1.6	0.3	0.0	0.0	0.9	0.0	0.0	0.0	1.6	1.8	1.4	0.8	1.1	0.0	0.0	3.3	0.6	0.6	0.1
1971	1.0	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.5	0.6	2.1	0.0	0.0	1.4	4.3	0.0	0.4	0.4	0.0
1972	0.6	0.5	0.3	0.0	0.3	0.3	0.0	0.0	0.9	0.8	0.0	2.3	1.3	3.2	3.0	1.3	0.0	0.0	0.6	0.6	0.1
1973	2.0	0.3	0.3	0.0	0.3	0.0	0.0	0.4	0.4	0.4	2.3	0.6	2.6	1.6	5.9	0.0	0.0	0.0	0.7	0.8	0.1
1969-73	0.9	0.4	0.5	0.2	0.2	0.1	0.2	0.2	0.3	0.5	1.1	1.1	1.9	1.6	2.5	0.6	0.9	0.6	0.6	0.6	0.1
1974	0.7	0.3	0.5	0.3	0.3	0.9	0.4	0.9	0.4	0.4	0.0	0.6	2.5	1.5	3.8	1.3	0.0	0.0	0.6	0.7	0.1
1975	0.0	0.6	0.0	0.3	0.3	0.9	0.0	0.0	0.4	0.4	0.4	0.0	2.4	3.0	3.7	1.2	1.9	0.0	0.6	0.5	0.1
1976	0.3	0.3	0.5	0.0	0.3	0.6	0.3	0.4	0.4	1.3	0.8	1.6	3.0	1.4	0.9	1.2	7.5	2.4	0.7	0.7	0.1
1977	1.0	0.3	0.3	0.3	0.5	0.8	1.0	1.2	0.0	1.7	2.1	2.0	1.2	2.1	0.0	2.4	5.4	2.3	0.9	0.9	0.1
1978	0.3	0.0	0.0	0.5	0.5	0.3	0.6	0.4	0.0	0.9	1.3	1.4	3.5	0.7	3.4	3.4	1.7	2.2	0.7	0.7	0.1
1974-78	0.5	0.3	0.3	0.3	0.4	0.7	0.5	0.6	0.3	0.9	0.9	1.1	2.5	1.7	2.3	1.9	3.4	1.4	0.7	0.7	0.1
1979	0.3	0.0	0.0	0.0	0.3	0.8	0.6	0.4	0.4	1.3	1.3	1.8	2.9	1.3	3.3	6.7	1.7	2.1	0.8	0.7	0.1
1980	0.3	0.0	0.0	0.0	0.5	0.5	0.8	0.7	0.0	0.9	1.3	0.9	3.9	2.5	2.4	7.5	1.6	2.0	0.8	0.7	0.1
1981	0.7	0.0	0.3	0.0	0.3	0.5	0.3	0.7	0.4	0.0	1.7	1.3	2.1	0.0	3.8	3.1	3.1	0.0	0.6	0.6	0.1
1982	0.3	0.0	0.0	0.5	0.2	0.8	0.0	0.6	0.8	1.3	0.4	1.7	0.5	3.0	2.2	2.0	3.1	0.0	0.6	0.6	0.1
1983	0.7	0.3	0.6	0.5	0.0	0.0	0.3	0.6	0.8	0.9	0.4	1.7	1.4	0.0	0.0	1.9	1.5	1.8	0.6	0.6	0.0
1979-83	0.5	0.1	0.2	0.2	0.3	0.5	0.4	0.6	0.5	0.9	1.0	1.5	2.1	1.4	2.3	4.1	2.2	1.2	0.7	0.6	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF THYROID GLAND (ICD9 193)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	1	1	1	4	0	4	5	1	1	1	6	3	3	1	3	1	0	36
1965	0	1	1	2	1	2	2	3	3	1	2	7	7	6	0	2	1	0	0	41
1966	0	0	0	1	2	3	2	2	5	0	3	3	4	9	3	3	1	0	0	41
1967	0	0	1	0	2	2	1	2	2	4	4	5	0	3	2	1	4	2	0	35
1968	0	1	3	5	8	11	5	13	17	13	13	19	18	26	15	10	11	4	0	188
1964-68																				
1969	0	1	0	3	0	3	2	1	7	0	0	4	4	2	6	2	2	0	0	37
1970	0	0	1	0	4	6	0	7	2	2	2	3	3	6	2	3	3	0	0	44
1971	0	0	0	0	3	3	0	5	3	2	3	6	2	6	0	1	0	1	1	36
1972	0	1	0	1	0	4	1	1	3	5	4	4	1	0	5	1	2	0	1	34
1973	0	0	0	1	2	3	3	3	2	4	2	2	2	4	7	2	0	0	1	38
1969-73																				
1974	0	2	1	5	9	19	6	17	17	13	11	19	12	18	20	9	7	1	3	189
1975	0	0	0	1	6	2	4	3	6	5	6	6	3	5	1	2	0	1	0	51
1976	0	1	0	3	3	4	4	2	2	5	8	4	3	6	5	2	0	3	1	56
1977	0	0	1	2	3	4	5	3	3	2	4	4	4	7	5	4	1	1	0	53
1978	0	1	0	4	4	4	4	4	8	5	6	6	4	8	3	2	0	3	0	66
1974-78																				
1979	0	2	2	13	18	21	20	18	26	24	32	25	18	29	18	12	1	8	1	288
1980	0	1	1	2	3	6	9	6	2	5	3	8	5	6	5	5	1	0	1	69
1981	0	0	1	5	0	3	9	9	8	7	3	9	7	8	5	5	1	0	0	80
1982	0	0	1	4	2	6	8	5	6	9	3	5	7	9	5	5	0	1	0	76
1983	0	0	0	3	6	8	6	8	3	6	4	6	3	3	4	3	1	1	0	65
1979-83																				
	0	1	3	15	12	25	40	34	23	33	20	35	29	37	26	20	3	4	1	361

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF THYROID GLAND (ICD9 193)

YEAR	AGE SPECIFIC RATES																	SUMMARY RATES		
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.3	0.4	0.5	2.0	0.0	1.7	2.3	0.5	0.6	0.7	5.3	3.3	4.4	2.2	11.8	7.4	1.1	1.1	0.1
1965	0.0	0.3	0.3	0.7	0.4	1.0	0.9	1.3	1.3	0.5	1.1	4.9	6.0	6.6	0.0	4.4	3.9	0.0	1.2	1.2	0.1
1966	0.0	0.0	0.0	0.3	0.8	1.4	0.9	0.8	2.1	0.0	1.7	2.0	3.3	9.7	4.3	6.4	3.8	0.0	1.2	1.2	0.1
1967	0.0	0.0	0.0	0.3	0.8	0.0	0.0	0.8	0.8	3.5	1.6	2.0	0.8	5.3	4.2	6.3	7.6	6.9	1.0	1.0	0.1
1968	0.0	0.0	0.3	0.0	0.7	0.8	0.4	0.8	0.8	1.9	2.2	3.2	0.0	3.1	2.8	2.1	14.9	13.3	1.0	0.9	0.1
1964-68	0.0	0.1	0.2	0.3	0.7	1.0	0.4	1.1	1.5	1.3	1.5	2.6	3.0	5.6	3.1	4.3	8.4	5.6	1.1	1.1	0.1
1969	0.0	0.2	0.3	0.9	0.0	1.2	0.9	0.4	2.9	0.0	0.0	2.5	3.1	2.0	8.4	4.2	7.4	0.0	1.0	1.0	0.1
1970	0.0	0.0	0.3	0.0	1.3	2.2	0.0	2.9	0.8	0.9	1.1	1.8	2.3	5.8	2.7	6.2	10.8	0.0	1.2	1.1	0.1
1971	0.0	0.0	0.0	0.0	0.9	1.0	0.0	2.1	1.2	0.9	1.6	3.5	1.5	5.7	0.0	2.0	0.0	5.8	0.9	0.9	0.1
1972	0.0	0.3	0.0	0.3	0.0	1.3	0.4	0.4	1.2	2.1	2.0	2.3	0.7	0.0	6.5	2.0	7.0	0.0	0.8	0.8	0.1
1973	0.0	0.0	0.0	0.3	0.6	0.9	1.2	1.3	0.8	1.7	1.0	1.2	1.4	3.6	8.8	4.0	0.0	0.0	0.9	0.9	0.1
1969-73	0.0	0.1	0.0	0.3	0.6	1.3	0.5	1.4	1.4	1.1	1.1	2.3	1.8	3.4	5.3	3.7	5.0	1.2	1.0	1.0	0.1
1974	0.0	0.0	0.0	0.3	1.7	0.6	1.5	1.2	2.5	2.1	2.7	3.6	2.0	4.4	1.2	4.0	0.0	5.4	1.2	1.2	0.1
1975	0.0	0.3	0.0	0.7	0.8	1.2	1.4	0.8	0.8	2.1	3.6	2.3	1.9	5.2	5.9	3.8	0.0	15.8	1.3	1.3	0.1
1976	0.0	0.0	0.2	0.5	0.8	1.1	1.7	1.2	1.2	0.8	1.8	2.2	2.5	5.8	5.8	7.4	3.4	5.2	1.2	1.2	0.1
1977	0.0	0.3	0.0	1.0	1.1	1.1	1.3	1.6	3.3	2.1	2.6	3.2	2.5	6.4	3.4	3.5	0.0	15.2	1.5	1.5	0.1
1978	0.0	0.0	0.3	0.7	0.5	2.0	0.9	2.3	2.9	2.9	3.5	2.5	2.5	2.3	4.4	3.4	0.0	0.0	1.4	1.4	0.1
1974-78	0.0	0.1	0.1	0.6	1.0	1.2	1.3	1.4	2.2	2.0	2.8	2.8	2.3	4.8	4.2	4.4	0.7	8.3	1.3	1.3	0.1
1979	0.0	0.3	0.3	0.5	0.8	1.7	2.6	2.2	0.8	2.1	1.3	3.8	3.2	4.5	5.3	8.2	3.1	0.0	1.5	1.4	0.1
1980	0.0	0.0	0.3	1.2	0.0	0.8	2.6	3.2	3.3	2.9	1.3	4.2	4.3	5.8	5.2	7.9	3.0	0.0	1.7	1.7	0.2
1981	0.0	0.0	0.3	1.0	0.5	1.7	2.2	1.7	2.4	3.8	1.3	2.3	4.1	6.4	5.0	7.7	0.0	4.7	1.6	1.6	0.2
1982	0.0	0.0	0.0	0.7	1.5	2.2	1.7	2.6	1.2	2.5	1.7	2.7	1.7	2.1	3.8	4.5	2.8	4.6	1.4	1.3	0.1
1983	0.0	0.0	0.0	0.3	0.2	0.5	2.3	1.8	1.5	2.5	3.0	3.2	3.7	7.7	6.4	2.9	0.0	9.0	1.4	1.4	0.2
1979-83	0.0	0.1	0.2	0.7	0.6	1.4	2.3	2.3	1.8	2.8	1.7	3.3	3.4	5.3	5.1	6.2	1.7	3.8	1.5	1.5	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF THYROID GLAND (ICD9 193)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	6	5	6	6	6	12	7	8	7	8	4	12	4	9	2	0	102
1965	0	0	1	3	8	11	7	14	9	6	9	6	11	7	6	9	4	2	0	113
1966	0	0	0	3	10	6	8	6	8	9	7	8	8	10	5	4	4	7	0	103
1967	0	0	0	4	3	5	11	8	7	11	4	9	11	8	9	5	4	2	0	101
1968	0	0	1	5	5	7	8	6	8	4	4	5	5	6	4	8	2	2	0	80
1964-68	0	0	2	21	31	35	40	40	44	37	32	35	43	35	36	30	23	15	0	499
1969	0	0	0	2	9	8	16	10	16	6	11	12	7	13	8	5	8	2	0	133
1970	0	0	0	7	9	10	11	10	10	10	5	7	11	8	9	0	6	3	0	116
1971	0	0	2	5	5	7	11	8	4	18	9	8	5	8	2	6	7	4	0	109
1972	0	0	1	6	7	9	7	10	4	9	10	5	8	9	5	3	3	3	0	99
1973	0	0	1	2	20	23	7	14	17	6	16	12	10	9	8	8	3	6	1	163
1969-73	0	0	4	22	50	57	52	52	51	49	51	44	41	47	32	22	27	18	1	620
1974	0	0	2	6	14	17	10	12	10	8	17	18	10	6	8	8	3	3	0	152
1975	2	0	3	2	12	5	23	12	10	20	16	14	4	7	9	5	3	5	1	153
1976	0	0	3	7	13	17	19	10	16	8	11	11	10	11	6	5	6	4	0	157
1977	0	2	0	10	8	11	17	16	14	15	7	10	11	15	10	8	6	4	0	164
1978	0	0	3	6	16	20	23	18	15	14	11	10	6	6	9	10	5	7	0	179
1974-78	2	2	11	31	63	70	92	68	65	65	62	63	41	45	42	36	23	23	1	805
1979	0	1	3	8	11	26	33	16	16	12	16	15	14	16	13	9	3	4	2	218
1980	0	0	2	12	14	21	25	22	13	16	15	22	10	9	13	4	9	6	0	213
1981	0	1	0	9	22	21	23	26	14	16	12	20	12	15	8	12	10	2	1	224
1982	0	0	1	6	12	21	13	16	10	13	16	10	15	12	14	16	8	4	1	188
1983	0	0	4	4	20	17	29	23	17	18	12	21	13	10	16	8	9	3	2	226
1979-83	0	2	10	39	79	106	123	103	70	75	71	88	64	62	64	49	39	19	6	1069

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF THYROID GLAND (ICD9 193)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	2.3	2.3	2.9	2.7	2.6	5.4	3.8	4.9	5.1	6.8	3.9	14.5	6.8	26.0	9.1	3.0	2.9	0.3
1965	0.0	0.0	0.3	1.1	3.5	5.2	3.2	6.0	3.9	3.2	5.3	4.2	9.1	6.8	7.0	14.9	11.1	8.7	3.3	3.2	0.3
1966	0.0	0.0	0.0	1.0	4.1	2.7	3.6	2.6	3.4	4.6	4.0	5.5	6.4	9.4	5.6	6.5	10.8	29.5	3.0	2.8	0.3
1967	0.0	0.0	0.0	1.3	1.1	2.2	4.9	3.5	2.9	5.4	2.2	5.9	8.6	7.4	10.0	7.7	10.3	8.0	2.8	2.7	0.3
1968	0.0	0.0	0.3	1.6	1.8	2.9	3.6	2.6	3.3	1.9	2.2	3.2	3.8	5.4	4.4	11.9	4.9	7.5	2.2	2.0	0.2
1964-68	0.0	0.0	0.1	1.4	2.5	3.2	3.6	3.4	3.8	3.8	3.7	4.8	6.9	6.6	8.2	9.6	12.3	12.5	2.8	2.7	0.3
1969	0.0	0.0	0.0	0.6	3.0	3.2	7.1	4.4	6.7	2.7	6.0	7.4	5.2	11.3	8.6	7.3	18.9	7.1	3.5	3.3	0.3
1970	0.0	0.0	0.0	2.0	2.8	3.8	4.8	4.4	4.3	4.3	2.7	4.2	7.9	6.8	9.5	0.0	13.6	9.9	3.0	2.9	0.3
1971	0.0	0.0	0.5	1.4	1.5	2.5	4.7	3.5	1.7	7.6	4.7	4.6	3.5	6.6	2.1	8.3	15.2	12.3	2.7	2.6	0.2
1972	0.0	0.0	0.3	1.7	2.1	3.0	2.9	4.4	1.7	3.8	4.9	2.9	5.4	7.2	5.0	4.0	6.3	8.7	2.4	2.3	0.2
1973	0.0	0.0	0.3	0.5	5.9	7.2	2.7	6.1	7.3	2.5	7.4	6.9	6.5	7.3	7.9	10.5	6.1	16.7	3.9	3.7	0.3
1969-73	0.0	0.0	0.2	1.3	3.1	4.0	4.4	4.5	4.4	4.2	5.2	5.2	5.7	7.7	6.6	6.1	11.8	11.2	3.1	3.0	0.3
1974	0.0	0.0	0.5	1.6	4.0	5.1	3.7	5.1	4.3	3.4	7.5	10.2	6.3	4.5	7.6	10.3	6.0	7.9	3.6	3.4	0.3
1975	0.7	0.0	0.8	0.5	3.3	1.4	8.2	5.0	4.3	8.5	6.8	7.7	2.4	5.2	8.3	6.3	5.8	12.7	3.5	3.4	0.3
1976	0.0	0.0	0.8	1.8	3.5	4.7	6.5	4.1	6.8	3.4	4.6	5.8	5.9	7.8	5.4	6.0	11.3	9.7	3.5	3.3	0.3
1977	0.0	0.6	0.0	2.5	2.1	3.1	5.4	6.4	6.0	6.4	3.0	4.9	6.4	10.3	8.7	9.4	10.8	9.2	3.6	3.4	0.3
1978	0.0	0.0	0.8	1.5	4.1	5.5	7.0	6.9	6.4	6.0	4.7	4.7	3.5	4.0	7.6	11.4	8.7	15.6	3.9	3.6	0.3
1974-78	0.1	0.1	0.6	1.6	3.4	4.0	6.2	5.5	5.6	5.6	5.3	6.5	4.9	6.4	7.5	8.7	8.6	11.1	3.6	3.4	0.3
1979	0.0	0.3	0.9	2.0	2.8	7.1	9.7	5.9	6.8	5.1	6.8	6.7	8.1	10.3	10.6	10.0	5.0	8.5	4.5	4.3	0.4
1980	0.0	0.0	0.6	3.0	3.6	5.6	7.1	7.9	5.4	6.9	6.4	9.6	5.6	5.6	10.3	4.3	14.6	12.2	4.5	4.2	0.4
1981	0.0	0.3	0.0	2.3	5.5	5.6	6.3	9.0	5.7	6.9	5.1	8.6	6.4	9.1	6.1	12.5	15.6	3.9	4.6	4.3	0.4
1982	0.0	0.0	0.3	1.6	3.0	5.5	3.6	5.1	4.0	5.6	6.8	4.3	7.6	7.2	10.4	16.0	12.2	7.4	3.7	3.4	0.3
1983	0.0	0.0	1.2	1.1	4.9	4.4	7.9	7.0	6.5	7.7	5.1	8.9	6.2	6.0	11.5	7.7	13.2	5.3	4.5	4.1	0.4
1979-83	0.0	0.1	0.6	2.0	4.0	5.6	6.9	6.9	5.7	6.5	6.1	7.6	6.7	7.6	9.8	10.1	12.2	7.3	4.4	4.1	0.4

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OTHER ENDOCRINE GLANDS AND RELATED STRUCTURES (ICD9 194)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	3	0	1	0	0	1	0	0	1	2	1	1	4	1	1	1	0	0	0	17
1965	6	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	8
1966	2	0	0	0	0	1	0	0	1	0	1	3	1	1	0	0	0	1	0	11
1967	3	0	0	0	0	1	0	1	0	3	0	0	4	0	0	1	1	0	0	14
1968	4	0	0	1	1	0	0	2	0	1	0	1	0	1	1	0	0	0	0	12
1964-68	18	0	1	1	1	3	0	3	2	6	2	7	9	3	2	2	1	1	0	62
1969	3	1	0	0	0	0	0	1	0	0	3	0	2	2	1	0	1	0	0	14
1970	4	1	0	0	1	0	1	0	2	0	0	0	0	1	0	0	0	0	0	10
1971	2	1	1	0	0	0	1	1	0	0	1	3	2	2	0	0	0	0	0	14
1972	2	0	1	1	2	0	0	0	0	0	1	2	0	2	1	0	1	1	0	14
1973	1	0	0	0	0	1	0	0	1	0	3	3	3	1	1	1	0	0	0	15
1969-73	12	3	2	1	3	1	2	2	3	0	8	8	7	8	3	1	2	1	0	67
1974	2	0	0	2	0	0	0	1	1	2	0	1	2	1	2	0	0	0	1	15
1975	2	0	1	0	0	2	0	0	2	0	1	4	3	1	0	0	0	0	1	17
1976	0	2	0	0	1	1	1	0	1	3	4	1	5	2	1	1	0	0	0	23
1977	0	2	1	1	1	0	1	0	1	1	2	1	3	2	0	0	0	0	0	16
1978	2	2	3	3	6	0	1	1	1	4	2	1	2	5	1	3	1	0	0	38
1974-78	6	6	5	6	8	3	3	2	6	10	9	8	15	11	4	4	1	0	2	109
1979	12	1	3	1	0	0	0	1	1	1	1	0	5	1	0	0	0	0	0	27
1980	9	1	0	2	0	0	1	2	3	2	2	1	5	3	3	0	1	0	0	35
1981	7	3	0	2	2	1	2	2	1	4	3	3	1	3	2	3	0	0	0	39
1982	4	2	0	2	2	2	0	1	2	3	1	1	2	1	0	1	1	0	0	25
1983	6	0	2	0	1	0	0	2	1	3	1	3	5	1	1	1	0	0	0	27
1979-83	38	7	5	7	5	3	3	8	8	13	8	8	18	9	6	5	2	0	0	153

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OTHER ENDOCRINE GLANDS AND RELATED STRUCTURES (ICD9 194)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.8	0.0	0.3	0.0	0.0	0.5	0.0	0.0	0.5	1.1	0.6	0.7	3.5	1.1	1.5	2.2	0.0	0.0	0.5	0.5	0.1
1965	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1966	0.5	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.4	0.0	0.6	2.0	0.8	1.1	0.0	0.0	0.0	7.1	0.3	0.3	0.0
1967	0.8	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.0	1.5	0.0	0.0	3.3	0.0	0.0	2.1	3.8	0.0	0.4	0.4	0.0
1968	1.1	0.0	0.0	0.3	0.4	0.0	0.0	0.8	0.0	0.5	0.0	0.6	0.0	1.0	1.4	0.0	0.0	0.0	0.3	0.4	0.0
1964-68	0.9	0.0	0.1	0.1	0.1	0.3	0.0	0.3	0.2	0.6	0.2	0.9	1.5	0.6	0.6	0.9	0.8	1.4	0.4	0.4	0.0
1969	0.9	0.2	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.6	0.0	1.6	2.0	1.4	0.0	3.7	0.0	0.4	0.4	0.0
1970	1.2	0.2	0.0	0.0	0.3	0.0	0.4	0.0	0.8	0.0	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1971	0.6	0.2	0.2	0.0	0.0	0.0	0.4	0.4	0.0	0.0	0.5	1.8	1.5	1.9	0.0	0.0	0.0	0.0	0.4	0.4	0.0
1972	0.6	0.0	0.2	0.3	0.6	0.0	0.0	0.0	0.0	0.0	0.5	1.2	0.0	1.9	1.3	0.0	3.5	5.6	0.4	0.4	0.0
1973	0.3	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	1.4	1.8	2.1	0.9	1.3	2.0	0.0	0.0	0.4	0.4	0.0
1969-73	0.7	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2	0.0	0.8	1.0	1.0	1.5	0.8	0.4	1.4	1.2	0.4	0.4	0.0
1974	0.6	0.0	0.0	0.5	0.0	0.0	0.0	0.4	0.4	0.8	0.0	0.6	1.3	0.9	2.4	0.0	0.0	0.0	0.3	0.4	0.0
1975	0.6	0.0	0.2	0.0	0.0	0.6	0.0	0.0	0.8	0.0	0.4	2.3	1.9	0.9	0.0	0.0	0.0	0.0	0.4	0.4	0.0
1976	0.0	0.6	0.0	0.0	0.3	0.3	0.3	0.0	0.4	1.3	1.8	0.6	3.2	1.7	1.2	1.8	0.0	0.0	0.5	0.6	0.1
1977	0.0	0.6	0.3	0.2	0.3	0.0	0.3	0.0	0.4	0.4	0.9	0.5	1.9	1.6	0.0	0.0	0.0	0.0	0.4	0.4	0.0
1978	0.6	0.6	0.8	0.7	1.6	0.0	0.3	0.4	0.4	1.7	0.9	0.5	1.3	3.8	1.1	5.1	3.2	0.0	0.9	0.9	0.1
1974-78	0.4	0.3	0.2	0.3	0.4	0.2	0.2	0.2	0.5	0.8	0.8	0.9	1.9	1.8	0.9	1.5	0.7	0.0	0.5	0.5	0.0
1979	3.9	0.3	0.8	0.2	0.0	0.0	0.0	0.4	0.4	0.4	0.4	0.0	3.2	0.7	0.0	0.0	0.0	0.0	0.7	0.8	0.1
1980	2.9	0.3	0.0	0.5	0.0	0.0	0.3	0.7	1.2	0.8	0.9	0.5	3.1	2.2	3.1	0.0	3.0	0.0	0.8	0.9	0.1
1981	2.3	0.9	0.0	0.5	0.5	0.3	0.6	0.7	0.4	1.7	1.3	1.4	0.6	2.1	2.0	4.6	0.0	0.0	0.9	1.0	0.1
1982	1.3	0.6	0.0	0.5	0.5	0.5	0.0	0.3	0.8	1.3	0.4	0.5	1.1	0.7	0.0	1.5	2.8	0.0	0.6	0.6	0.0
1983	1.9	0.0	0.6	0.0	0.2	0.0	0.0	0.6	0.4	1.3	0.4	1.4	2.7	0.7	0.9	1.5	0.0	0.0	0.6	0.7	0.1
1979-83	2.5	0.4	0.3	0.3	0.3	0.2	0.2	0.5	0.6	1.1	0.7	0.7	2.1	1.3	1.2	1.5	1.2	0.0	0.7	0.8	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OTHER ENDOCRINE GLANDS AND RELATED STRUCTURES (ICD9 194)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	3	0	1	0	0	0	0	0	0	0	0	1	1	0	1	0	1	1	0	9
1965	5	0	0	0	0	0	2	1	1	1	0	0	1	1	0	0	0	0	0	12
1966	2	1	0	0	1	0	0	0	1	0	0	1	0	0	0	1	0	0	0	7
1967	2	0	0	1	0	0	0	0	1	0	3	0	1	1	0	1	0	0	0	10
1968	1	0	0	0	0	0	0	1	0	2	0	0	0	1	1	1	0	0	0	7
1964-68	13	1	1	1	1	0	2	2	3	3	3	2	3	3	2	3	1	1	0	45
1969	0	0	0	0	0	0	0	1	1	1	0	2	0	1	0	1	0	0	0	7
1970	0	0	0	0	0	0	0	0	0	1	2	0	0	0	0	0	1	0	0	4
1971	1	0	0	0	0	1	0	1	0	2	0	0	2	1	0	0	0	0	0	8
1972	1	0	1	0	1	0	0	0	0	2	0	2	0	0	1	0	0	0	0	8
1973	3	0	0	0	0	0	0	2	1	0	1	1	1	1	0	1	0	0	0	11
1969-73	5	0	1	0	1	1	0	4	2	6	3	5	3	3	1	2	1	0	0	38
1974	1	0	0	2	1	1	1	1	1	4	1	2	1	0	1	0	0	0	0	17
1975	1	0	0	0	1	2	0	0	2	3	1	0	1	2	0	0	0	0	0	13
1976	1	0	0	0	2	6	0	0	1	0	2	1	1	1	0	1	1	1	0	18
1977	0	1	0	0	1	0	0	1	1	5	1	0	1	1	0	0	0	1	0	13
1978	0	0	2	1	1	1	2	1	1	1	3	4	1	0	2	1	1	0	0	22
1974-78	3	1	2	3	6	10	3	3	6	13	8	7	5	4	3	2	2	2	0	83
1979	8	0	0	0	2	2	3	1	3	1	3	3	4	3	2	3	1	1	0	40
1980	6	2	0	0	1	2	4	0	3	0	2	3	6	2	2	2	0	0	0	35
1981	5	0	1	1	0	1	1	0	2	3	3	4	2	5	3	1	0	0	0	32
1982	3	2	0	0	1	1	1	0	4	1	1	2	2	3	3	0	0	0	0	23
1983	4	0	0	0	1	1	1	0	2	1	2	3	3	1	4	2	1	2	1	29
1979-83	26	4	1	1	5	7	8	7	10	7	13	15	16	17	9	7	3	2	1	159

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OTHER ENDOCRINE GLANDS AND RELATED STRUCTURES (ICD9 194)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.8	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.8	0.0	1.2	0.0	2.9	4.5	0.3	0.2	0.0
1965	1.3	0.0	0.0	0.0	0.0	0.0	0.9	0.4	0.4	0.5	0.0	0.0	0.8	1.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0
1966	0.6	0.3	0.0	0.0	0.4	0.0	0.0	0.0	0.4	0.0	0.0	0.7	0.0	0.0	0.0	1.6	0.0	0.0	0.2	0.2	0.0
1967	0.6	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.0	1.7	0.0	0.8	0.9	0.0	1.5	0.0	0.0	0.3	0.3	0.0
1968	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	0.0	0.0	0.0	0.9	1.1	1.5	0.0	0.0	0.2	0.2	0.0
1964-68	0.7	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.3	0.3	0.3	0.3	0.5	0.6	0.5	1.0	0.5	0.8	0.3	0.3	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.0	1.2	0.0	0.9	0.0	1.5	0.0	0.0	0.2	0.2	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.1	0.0	0.0	0.0	0.0	0.0	2.3	0.0	0.1	0.1	0.0
1971	0.3	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.0	0.8	0.0	0.0	1.4	0.8	0.0	0.0	0.0	0.0	0.2	0.2	0.0
1972	0.3	0.0	0.3	0.0	0.3	0.0	0.0	0.0	0.0	0.8	0.0	1.1	0.0	0.0	1.0	0.0	0.0	0.0	0.2	0.2	0.0
1973	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.4	0.0	0.5	0.6	0.6	0.8	0.0	1.3	0.0	0.0	0.3	0.3	0.0
1969-73	0.3	0.0	0.1	0.0	0.1	0.1	0.0	0.3	0.2	0.5	0.3	0.6	0.4	0.5	0.2	0.6	0.4	0.0	0.2	0.2	0.0
1974	0.3	0.0	0.0	0.5	0.3	0.3	0.4	0.4	0.4	1.7	0.4	1.1	0.6	0.0	1.0	0.0	0.0	0.0	0.4	0.4	0.0
1975	0.3	0.0	0.0	0.0	0.3	0.6	0.0	0.0	0.9	1.3	0.4	0.0	0.6	1.5	0.0	0.0	0.0	0.0	0.3	0.3	0.0
1976	0.3	0.0	0.0	0.0	0.5	1.7	0.0	0.0	0.4	0.0	0.8	0.5	0.6	0.7	0.0	1.2	1.9	2.4	0.4	0.4	0.0
1977	0.0	0.3	0.0	0.0	0.3	0.0	0.0	0.4	0.4	2.1	0.4	0.0	0.6	0.7	0.0	0.0	0.0	2.3	0.3	0.3	0.0
1978	0.0	0.0	0.6	0.2	0.3	0.3	0.6	0.4	0.4	0.4	1.3	1.9	0.6	0.0	1.7	1.1	1.7	0.0	0.5	0.4	0.0
1974-78	0.2	0.1	0.1	0.2	0.3	0.6	0.2	0.2	0.5	1.1	0.7	0.7	0.6	0.6	0.5	0.5	0.7	1.0	0.4	0.4	0.0
1979	2.7	0.0	0.0	0.0	0.5	0.5	0.9	0.4	1.3	0.4	1.3	1.3	2.3	1.9	1.6	3.3	1.7	2.1	0.9	0.9	0.1
1980	2.1	0.7	0.0	0.0	0.3	0.5	1.1	0.0	1.3	0.0	0.9	1.3	3.4	1.2	1.6	2.1	0.0	0.0	0.8	0.8	0.1
1981	1.7	0.0	0.3	0.3	0.0	0.3	0.3	0.0	0.8	1.3	1.3	1.7	1.1	3.0	2.3	1.0	0.0	0.0	0.7	0.7	0.1
1982	1.0	0.7	0.0	0.0	0.2	0.3	0.0	1.3	0.4	0.4	0.9	0.9	1.5	1.8	0.0	0.0	0.0	0.0	0.5	0.5	0.0
1983	1.4	0.0	0.0	0.0	0.2	0.3	0.0	0.6	0.4	0.9	1.3	1.3	0.5	2.4	1.4	1.0	2.9	1.8	0.6	0.6	0.1
1979-83	1.8	0.3	0.1	0.1	0.3	0.4	0.4	0.5	0.8	0.6	1.1	1.3	1.7	2.1	1.4	1.4	0.9	0.8	0.7	0.7	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

LYMPHOSARCOMA AND RETICULOSARCOMA (ICD9 200)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	6	2	2	3	2	2	5	12	15	14	24	24	32	21	21	15	8	6	0	214
1965	0	3	4	4	5	4	3	5	9	17	20	20	24	28	19	13	9	5	0	192
1966	0	4	1	5	3	4	1	7	10	10	17	24	26	22	18	17	10	5	0	184
1967	2	4	7	1	4	4	4	5	17	9	20	26	26	22	15	12	9	5	0	192
1968	2	2	6	4	5	4	12	5	13	13	15	27	27	14	14	16	6	6	0	191
1964-68	10	15	20	17	19	18	25	34	64	63	96	121	135	107	87	73	42	27	0	973
1969	5	6	4	4	3	10	6	7	12	9	9	28	33	30	17	10	16	0	0	209
1970	1	7	7	1	4	7	7	10	14	11	17	25	21	20	18	13	6	9	0	198
1971	4	3	4	1	5	6	7	7	13	27	15	29	31	28	26	19	7	2	0	234
1972	0	0	4	4	3	3	5	14	16	21	30	24	30	28	10	28	9	1	0	230
1973	3	1	5	4	7	5	10	6	18	19	19	28	34	45	27	22	18	4	0	275
1969-73	13	17	24	14	22	31	35	44	73	87	90	134	149	151	98	92	56	16	0	1146
1974	9	4	4	4	10	2	9	8	9	19	25	40	30	29	30	20	15	6	0	273
1975	4	4	4	4	5	8	8	13	21	12	25	26	38	33	34	18	14	9	0	280
1976	4	8	3	8	3	6	7	10	16	18	24	37	34	26	34	20	14	7	0	279
1977	3	5	2	0	6	6	13	8	17	20	26	31	37	38	30	31	11	7	1	292
1978	1	8	6	11	6	5	6	9	16	16	29	37	41	42	32	23	18	9	2	317
1974-78	21	29	19	27	30	27	43	48	79	85	129	171	180	168	160	112	72	38	3	1441
1979	3	2	7	10	7	2	6	11	12	27	21	37	33	42	37	36	13	3	0	309
1980	0	3	3	8	2	9	15	7	8	11	17	41	35	41	29	26	14	7	0	276
1981	0	1	3	9	7	4	7	10	9	17	27	25	39	37	33	31	19	8	0	286
1982	0	2	0	4	7	2	1	11	12	16	23	30	42	47	42	30	24	11	3	307
1983	2	1	0	6	4	3	5	9	8	18	27	37	45	37	39	34	23	6	1	305
1979-83	5	9	13	37	27	20	34	48	49	89	115	170	194	204	180	157	93	35	4	1483

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

LYMPHOSARCOMA AND RETICULOSARCOMA (ICD9 200)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	1.5	0.5	0.6	1.1	0.9	1.0	2.2	5.1	6.9	7.4	14.1	17.2	28.1	23.3	30.7	33.0	31.5	44.1	6.4	6.4	0.7
1965	0.0	0.8	1.2	1.4	2.2	1.9	1.3	2.1	4.0	8.9	11.4	14.0	20.6	30.6	27.4	28.3	34.7	36.0	5.7	5.7	0.6
1966	0.0	1.0	0.3	1.6	1.2	1.9	0.4	2.9	4.3	5.1	9.5	16.3	21.7	23.6	25.6	36.5	38.5	35.7	5.3	5.2	0.6
1967	0.5	1.0	1.9	0.3	1.5	1.8	1.8	2.1	7.1	4.5	11.0	17.1	21.2	23.2	21.2	25.3	34.1	34.5	5.4	5.3	0.6
1968	0.6	0.5	1.6	1.2	1.8	1.7	5.2	2.1	5.4	6.2	8.2	17.2	21.4	14.4	19.7	33.5	22.4	40.0	5.3	5.2	0.5
1964-68	0.5	0.8	1.1	1.1	1.6	1.6	2.2	2.8	5.5	6.4	10.8	16.4	22.5	22.9	24.9	31.3	32.2	38.0	5.6	5.6	0.6
1969	1.4	1.5	1.0	1.2	1.0	4.0	2.6	2.9	5.0	4.1	4.9	17.2	25.6	30.0	23.7	20.8	58.8	0.0	5.7	5.6	0.6
1970	0.3	1.7	1.8	0.3	1.3	2.6	3.0	4.2	5.8	4.9	9.2	15.0	15.9	19.4	24.7	26.8	21.6	54.9	5.2	5.2	0.5
1971	1.2	0.7	1.0	0.3	1.5	2.1	2.9	2.9	5.4	11.6	7.9	17.0	22.8	26.4	34.9	38.9	24.8	11.6	6.1	6.0	0.7
1972	3.0	0.0	1.0	1.1	0.9	1.0	2.0	5.9	6.6	9.0	15.1	14.1	21.3	25.9	13.0	56.7	31.4	5.6	5.8	5.7	0.6
1973	0.9	0.3	1.2	1.0	2.1	1.6	3.8	2.5	7.4	8.1	9.1	16.5	23.4	40.9	33.9	44.2	61.9	22.0	6.9	6.8	0.8
1969-73	0.8	0.9	1.2	0.8	1.4	2.2	2.9	3.7	6.0	7.6	9.3	15.9	21.8	28.6	26.1	37.6	39.7	18.7	6.0	5.9	0.6
1974	2.8	1.1	1.0	1.0	2.9	0.6	3.3	3.3	3.7	8.0	11.4	23.7	20.0	25.7	36.5	39.5	51.4	32.4	6.7	6.6	0.7
1975	1.2	1.2	1.0	1.0	1.4	2.3	2.8	5.3	8.7	5.0	11.2	15.1	24.5	28.4	40.3	34.4	47.9	47.4	6.7	6.7	0.7
1976	1.3	2.3	0.7	1.9	0.8	1.7	2.4	4.0	6.6	7.5	10.6	20.7	21.6	21.6	39.4	36.8	47.1	36.1	6.6	6.5	0.7
1977	1.0	1.5	0.5	0.0	1.6	1.7	4.1	3.1	7.1	8.3	11.4	16.4	23.4	30.2	34.0	54.9	36.4	35.5	6.7	6.6	0.7
1978	0.3	2.4	1.6	2.6	1.6	1.4	1.8	3.4	6.7	6.6	12.7	18.6	26.0	32.3	35.3	39.0	58.4	45.0	7.2	7.0	0.8
1974-78	1.3	1.7	0.9	1.3	1.6	1.5	2.9	3.8	6.6	7.1	11.5	18.8	23.1	27.8	37.0	41.1	48.3	39.3	6.8	6.7	0.7
1979	1.0	0.6	1.9	2.4	1.8	0.6	1.8	4.0	5.0	11.2	9.1	17.7	21.1	31.2	39.5	58.8	40.9	14.6	6.9	6.7	0.7
1980	0.0	0.9	0.8	1.9	0.5	2.5	4.3	2.5	3.3	4.6	7.3	19.3	21.7	29.5	29.9	41.3	42.3	33.8	5.9	5.7	0.6
1981	0.0	0.3	0.9	2.2	1.8	1.1	2.0	3.4	3.6	7.2	11.5	11.6	23.1	26.2	32.8	47.9	55.7	37.7	6.1	5.9	0.6
1982	0.0	0.6	0.0	1.0	1.7	0.5	0.3	3.5	4.7	6.8	9.7	13.7	23.6	32.9	40.2	45.1	67.2	50.9	6.3	6.1	0.7
1983	0.6	0.3	0.0	1.6	1.0	0.8	1.4	2.7	3.0	7.6	11.4	16.7	24.0	26.0	35.7	49.6	60.8	26.9	6.1	5.9	0.7
1979-83	0.3	0.6	0.7	1.8	1.4	1.1	1.9	3.2	3.9	7.5	9.8	15.8	22.8	29.2	35.6	48.5	53.9	32.9	6.3	6.1	0.7

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

LYMPHOSARCOMA AND RETICULOSARCOMA (ICD9 200)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	2	1	0	3	1	2	5	1	9	10	10	19	14	21	17	13	9	2	0	139
1965	0	0	1	2	2	3	2	5	8	12	12	22	18	17	20	13	7	4	0	148
1966	0	1	0	1	2	4	3	3	12	13	15	16	13	23	16	13	8	7	0	150
1967	1	1	1	0	1	2	2	7	12	10	11	17	21	28	18	17	8	0	0	157
1968	0	0	0	6	4	1	6	6	11	10	12	12	23	22	16	20	5	2	0	156
1964-68	3	3	2	12	10	12	18	22	52	55	60	86	89	111	87	76	37	15	0	750
1969	1	1	1	3	3	1	1	2	6	10	13	20	17	16	22	15	20	7	0	159
1970	0	2	2	1	2	2	2	6	11	19	18	25	22	30	21	11	11	3	0	188
1971	0	3	0	1	4	5	4	5	10	21	19	5	25	23	21	19	11	3	0	179
1972	3	5	2	2	3	3	7	5	6	16	19	22	18	22	20	23	10	8	1	195
1973	2	3	2	1	5	6	3	6	15	8	14	27	38	31	29	28	20	7	1	246
1969-73	6	14	7	8	17	17	17	24	48	74	83	99	120	122	113	96	72	28	2	967
1974	2	1	1	3	2	8	6	8	10	25	20	21	32	30	33	34	13	10	0	259
1975	1	0	1	2	4	10	3	7	10	15	24	31	22	38	30	24	20	11	0	253
1976	0	0	3	2	3	7	3	7	5	12	15	19	28	45	33	28	14	14	0	238
1977	0	1	3	4	0	8	7	10	8	19	29	28	28	28	34	30	25	10	1	273
1978	1	3	1	4	4	1	6	10	7	22	30	31	41	41	35	27	21	6	1	292
1974-78	4	5	9	15	13	34	25	42	40	93	118	130	151	182	165	143	93	51	2	1315
1979	2	2	3	1	2	6	3	6	10	8	6	23	24	38	28	23	28	15	0	228
1980	0	1	1	2	2	4	3	7	8	19	27	35	25	31	31	25	22	19	2	264
1981	0	1	2	2	1	3	6	8	8	13	23	27	35	26	34	27	27	17	0	260
1982	1	0	0	1	2	3	6	10	4	20	18	25	37	36	26	25	22	14	2	252
1983	0	1	1	2	2	3	10	6	11	11	15	30	24	31	35	32	23	18	0	255
1979-83	3	5	7	8	9	19	28	37	41	71	89	140	145	162	154	132	122	83	4	1259

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

LYMPHOSARCOMA AND RETICULOSARCOMA (ICD9 200)

YEAR	AGE SPECIFIC RATES																	SUMMARY RATES		
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.5	0.3	0.0	1.1	0.5	1.0	2.3	0.4	4.1	5.4	6.1	13.8	11.9	20.7	20.5	22.2	26.0	9.1	4.2	3.8	0.4
1965	0.0	0.0	0.3	0.7	0.9	1.4	0.9	2.1	3.5	6.3	7.1	15.5	14.9	16.4	23.4	21.5	19.5	17.4	4.4	4.0	0.5
1966	0.0	0.3	0.0	0.3	0.8	1.8	1.4	1.3	5.1	6.6	8.6	10.9	10.4	21.7	18.1	21.0	21.5	29.5	4.3	3.9	0.4
1967	0.3	0.3	0.3	0.0	0.4	0.9	0.9	3.0	5.0	4.9	6.2	11.2	16.4	25.8	20.0	26.3	20.5	0.0	4.3	4.0	0.5
1968	0.0	0.0	0.0	1.9	1.4	0.4	2.7	2.6	4.6	4.7	6.6	7.6	17.4	19.8	17.5	29.9	12.3	7.5	4.2	3.9	0.4
1964-68	0.2	0.2	0.1	0.8	0.8	1.1	1.6	1.9	4.5	5.6	6.9	11.7	14.3	20.9	19.8	24.3	19.7	12.5	4.3	3.9	0.5
1969	0.3	0.3	0.3	0.9	1.0	0.4	0.4	0.9	2.5	4.5	7.1	12.3	12.5	14.0	23.8	21.8	47.2	24.7	4.2	3.6	0.4
1970	0.0	0.5	0.5	0.3	0.6	0.8	0.9	2.6	4.7	8.2	9.6	14.8	15.8	25.4	22.2	15.6	24.9	9.9	4.8	4.5	0.5
1971	0.0	0.8	0.0	0.3	1.2	1.8	1.7	2.2	4.3	8.8	9.9	2.9	17.4	18.9	21.6	26.4	23.8	9.2	4.5	4.1	0.5
1972	1.0	1.4	0.5	0.6	0.9	1.0	2.9	2.2	2.6	6.8	9.3	12.6	12.1	17.6	20.2	31.0	21.1	23.3	4.7	4.3	0.5
1973	0.7	0.9	0.5	0.3	1.5	1.9	1.2	2.6	6.5	3.4	6.5	15.4	24.7	24.1	28.5	36.8	40.8	19.5	5.8	5.2	0.6
1969-73	0.4	0.7	0.4	0.5	1.0	1.2	1.4	2.1	4.1	6.4	8.4	11.6	16.6	20.1	23.3	26.5	31.4	17.4	4.8	4.3	0.5
1974	0.7	0.3	0.3	0.8	0.6	2.4	2.2	3.4	4.3	10.6	8.8	11.9	20.1	22.7	31.5	43.6	25.9	26.4	6.0	5.4	0.6
1975	0.3	0.0	0.3	0.5	1.1	2.9	1.1	2.9	4.3	6.4	10.3	17.1	13.3	28.1	27.8	30.0	38.8	27.9	5.7	5.1	0.6
1976	0.0	0.0	0.8	0.5	0.8	1.9	1.0	2.9	2.1	5.2	6.3	10.0	16.6	32.1	29.8	33.9	26.3	33.8	5.2	4.6	0.5
1977	0.0	0.3	0.8	1.0	0.0	2.2	2.2	4.0	3.4	8.2	12.3	13.8	16.4	19.2	29.6	35.4	45.1	23.1	5.8	5.1	0.6
1978	0.3	0.9	0.3	1.0	1.0	0.3	1.8	3.8	3.0	9.4	12.7	14.5	23.8	27.3	29.5	30.9	36.7	13.4	6.1	5.6	0.6
1974-78	0.3	0.3	0.5	0.8	0.7	1.9	1.7	3.4	3.4	8.0	10.1	13.5	18.0	25.9	29.6	34.6	34.8	24.6	5.8	5.1	0.6
1979	0.7	0.6	0.9	0.2	0.5	1.6	0.9	2.2	4.2	3.4	2.6	10.2	13.8	24.5	22.8	25.5	47.0	31.8	4.6	4.0	0.4
1980	0.0	0.3	0.3	0.5	0.5	1.1	0.8	2.5	3.3	8.2	11.6	15.2	14.0	19.2	24.6	26.8	35.6	38.7	5.2	4.6	0.5
1981	0.0	0.3	0.6	0.5	0.3	0.8	1.6	2.8	3.3	5.6	9.9	11.6	18.6	15.8	26.1	28.0	42.3	33.0	5.0	4.4	0.5
1982	0.3	0.0	0.0	0.3	0.5	0.8	1.7	3.2	1.6	8.6	7.7	10.7	18.7	21.5	19.3	25.0	33.6	25.9	4.7	4.2	0.5
1983	0.0	0.3	0.3	0.5	0.5	0.8	2.7	1.8	4.2	4.7	6.4	12.8	11.4	18.6	25.1	30.9	33.8	31.6	4.6	4.0	0.4
1979-83	0.2	0.3	0.4	0.4	0.5	1.0	1.6	2.5	3.3	6.1	7.6	12.1	15.3	19.9	23.6	27.3	38.3	32.1	4.8	4.2	0.5

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

HODGKIN'S DISEASE (ICD9 201)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	2	2	4	14	10	6	13	13	3	5	8	9	14	9	6	4	1	1	1	125
1965	0	4	3	6	14	11	9	8	13	8	9	13	7	8	6	5	6	3	0	133
1966	1	3	6	8	7	8	9	10	12	9	15	6	7	6	3	4	0	0	0	114
1967	0	1	6	4	13	9	5	5	17	9	8	13	13	11	8	3	0	1	0	126
1968	1	2	6	11	8	17	12	7	11	5	2	6	6	10	2	7	2	0	0	115
1964-68	4	12	25	43	52	51	48	43	56	36	42	47	47	44	25	23	9	5	1	613
1969	0	2	5	6	18	16	9	5	5	6	6	9	10	6	4	2	2	3	0	114
1970	2	3	5	12	14	20	13	11	12	13	7	7	7	9	5	3	9	1	0	153
1971	0	0	1	10	17	14	9	11	7	9	9	5	6	6	5	4	10	0	0	123
1972	0	6	5	10	14	16	10	5	10	11	7	15	10	5	6	5	5	3	0	143
1973	1	4	7	7	20	14	9	9	12	6	6	9	6	4	8	6	2	2	0	132
1969-73	3	15	23	45	83	80	50	41	46	45	35	45	39	30	28	20	28	9	0	665
1974	1	0	6	12	17	14	16	5	13	13	7	5	9	6	5	6	0	2	1	138
1975	1	2	2	15	21	15	7	8	10	4	13	5	8	8	3	2	1	3	0	128
1976	0	2	7	8	17	13	14	4	5	7	6	7	9	8	6	2	0	0	0	115
1977	1	5	5	8	18	17	8	7	11	8	5	7	6	6	7	4	3	0	0	126
1978	0	1	6	11	15	21	13	8	13	8	12	13	11	5	2	6	3	1	0	149
1974-78	3	10	26	54	88	80	58	32	52	40	43	37	43	33	23	20	7	6	1	656
1979	1	3	11	13	19	12	10	15	10	11	11	9	5	10	10	5	1	1	0	157
1980	2	2	5	18	16	23	16	12	5	6	8	11	11	14	8	4	0	3	0	164
1981	0	1	6	13	15	18	16	13	11	7	5	10	12	7	10	5	1	4	0	154
1982	0	1	6	18	15	19	14	10	15	10	10	8	7	5	11	5	5	2	0	161
1983	0	5	8	13	27	25	10	19	7	6	6	11	12	7	7	2	3	4	0	172
1979-83	3	12	36	75	92	97	66	69	48	40	40	49	47	43	46	21	10	14	0	808

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

HODGKIN'S DISEASE (ICD9 201)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.5	0.5	1.2	5.1	4.7	2.9	5.8	5.5	1.4	2.7	4.7	6.4	12.3	10.0	8.8	8.8	3.9	7.4	3.7	3.8	0.4
1965	0.0	1.0	0.9	2.1	6.2	5.3	4.0	3.4	5.8	4.2	5.1	9.1	6.0	8.7	8.6	10.9	23.2	21.6	3.9	4.0	0.4
1966	0.3	0.8	1.7	2.6	2.9	3.7	4.0	4.2	5.2	4.6	8.4	4.1	5.9	6.4	4.3	8.6	0.0	0.0	3.2	3.3	0.3
1967	0.0	0.3	1.7	1.3	5.0	4.0	2.2	2.1	7.1	4.5	4.4	8.6	10.6	11.6	11.3	6.3	0.0	6.9	3.5	3.6	0.4
1968	0.3	0.5	1.6	3.3	2.9	7.1	5.2	2.9	4.6	2.4	1.1	3.8	4.8	10.3	2.8	14.6	7.5	0.0	3.2	3.2	0.3
1964-68	0.2	0.6	1.4	2.8	4.3	4.7	4.2	3.6	4.8	3.7	4.7	6.4	7.8	9.4	7.1	9.9	6.9	7.0	3.5	3.6	0.3
1969	0.0	0.5	1.3	1.8	6.0	6.4	3.9	2.1	2.1	2.8	3.3	5.5	7.8	6.0	5.6	4.2	7.4	19.1	3.1	3.1	0.3
1970	0.6	0.7	1.3	3.4	4.4	7.4	5.5	4.6	5.0	5.7	3.8	4.2	5.3	8.7	6.9	6.2	32.4	6.1	4.1	4.0	0.3
1971	0.0	0.0	0.2	2.8	5.1	4.9	3.7	4.6	2.9	3.9	4.8	2.9	4.4	5.7	6.7	8.2	35.5	0.0	3.2	3.1	0.3
1972	0.0	1.5	1.2	2.7	4.2	5.2	4.0	2.1	4.1	4.7	3.5	8.8	7.1	4.6	7.8	10.1	17.4	16.8	3.6	3.5	0.3
1973	0.3	1.1	1.7	1.8	5.9	4.4	3.5	3.8	4.9	2.6	2.9	5.3	4.1	3.6	10.1	12.0	6.9	11.0	3.3	3.2	0.3
1969-73	0.2	0.8	1.1	2.5	5.1	5.6	4.1	3.4	3.8	3.9	3.6	5.4	5.7	5.7	7.5	8.2	19.9	10.5	3.5	3.4	0.3
1974	0.3	0.0	1.4	3.1	4.9	4.2	5.8	2.1	5.3	5.5	3.2	3.0	6.0	5.3	6.1	11.9	0.0	10.8	3.3	3.3	0.3
1975	0.3	0.6	0.5	3.7	5.8	4.3	2.4	3.3	4.1	1.7	5.8	2.9	5.2	6.9	3.6	3.8	3.4	15.8	3.1	3.0	0.3
1976	0.0	0.6	1.7	1.9	4.6	3.6	4.7	1.6	2.1	2.9	2.6	3.9	5.7	6.6	6.9	3.7	0.0	0.0	2.7	2.6	0.2
1977	0.3	1.5	1.3	1.9	4.8	4.8	2.5	2.8	4.6	3.3	2.2	3.7	3.8	4.8	7.9	7.1	9.9	0.0	3.0	2.9	0.3
1978	0.0	0.3	1.6	2.6	3.9	6.0	3.9	3.0	5.4	3.3	5.2	6.5	7.0	3.8	2.2	10.2	9.7	5.0	3.4	3.3	0.3
1974-78	0.2	0.6	1.3	2.6	4.8	4.6	3.9	2.6	4.3	3.3	3.8	4.1	5.5	5.5	5.3	7.3	4.7	6.2	3.1	3.0	0.3
1979	0.3	0.9	3.0	3.1	4.9	3.4	2.9	5.5	4.2	4.6	4.8	4.3	3.2	7.4	10.7	8.2	3.1	4.9	3.6	3.5	0.3
1980	0.7	0.6	1.4	4.3	4.1	6.4	4.6	4.2	2.1	2.5	3.4	5.2	6.8	10.1	8.2	6.4	0.0	14.5	3.7	3.6	0.3
1981	0.0	0.3	1.7	3.1	3.8	5.0	4.5	4.5	4.4	3.0	2.1	4.6	7.1	5.0	9.9	7.7	2.9	18.9	3.4	3.3	0.3
1982	0.0	0.3	1.7	4.5	3.7	5.2	4.0	3.2	5.9	4.2	4.2	3.7	3.9	3.5	10.5	7.5	14.0	9.3	3.6	3.4	0.3
1983	0.0	1.6	2.3	3.4	6.6	6.7	2.8	5.8	2.7	2.5	2.5	5.0	6.4	4.9	6.4	2.9	7.9	17.9	3.8	3.6	0.3
1979-83	0.2	0.8	2.1	3.7	4.6	5.4	3.8	4.6	3.8	3.4	3.4	4.6	5.5	6.1	9.1	6.5	5.8	13.2	3.6	3.5	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

HODGKIN'S DISEASE (ICD9 201)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	3	8	11	10	7	9	2	4	6	4	3	6	7	3	2	2	0	87
1965	1	0	5	5	14	2	4	5	11	3	6	6	4	7	4	2	2	2	0	83
1966	0	1	4	12	12	5	8	5	4	3	8	8	7	6	1	7	2	1	0	94
1967	1	1	1	5	12	9	6	2	7	6	3	5	7	5	6	5	1	0	0	82
1968	1	1	1	6	10	8	3	5	11	1	11	7	4	7	2	3	3	1	0	85
1964-68	3	3	14	36	59	34	28	26	35	17	34	30	25	31	20	20	10	6	0	431
1969	1	1	0	14	14	13	4	4	7	5	7	3	6	4	9	3	0	0	0	95
1970	0	2	4	17	12	6	4	10	6	6	10	7	8	2	3	3	2	1	0	103
1971	0	4	4	6	9	5	5	11	9	3	2	6	5	4	11	3	4	1	0	92
1972	1	1	8	8	22	10	9	9	2	2	7	3	6	4	4	7	2	1	0	106
1973	0	0	6	13	14	12	5	5	4	7	1	7	4	3	6	3	3	0	0	93
1969-73	2	8	22	58	71	46	27	39	28	23	27	26	29	17	33	19	11	3	0	489
1974	1	2	6	7	11	14	10	5	7	6	7	6	1	11	6	4	1	1	0	106
1975	0	0	3	15	18	8	11	3	6	5	7	1	4	8	2	6	3	2	1	103
1976	0	1	5	9	15	14	10	3	6	7	2	7	3	5	3	2	0	0	1	93
1977	0	2	5	8	22	14	8	5	5	3	5	6	6	6	2	2	3	2	0	104
1978	0	0	3	12	14	12	4	4	3	4	8	6	3	5	7	3	5	2	0	95
1974-78	1	5	22	51	80	62	43	20	27	25	29	26	17	35	20	17	12	7	2	501
1979	2	2	5	10	9	14	10	8	5	1	3	7	7	3	17	4	1	1	0	109
1980	0	1	3	13	15	17	9	5	6	9	4	3	8	7	3	3	2	4	0	112
1981	1	0	3	8	12	8	10	5	10	4	4	4	3	8	3	2	7	1	0	96
1982	0	1	9	8	19	19	12	11	6	9	3	3	9	9	4	6	5	4	0	137
1983	0	1	2	8	12	16	11	12	2	6	4	7	7	6	8	4	1	2	0	109
1979-83	3	5	22	47	67	74	52	41	29	29	18	24	37	33	35	19	16	12	0	563

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

HODGKIN'S DISEASE (ICD9 201)

YEAR	AGE SPECIFIC RATES																		SUMMARY RATES		
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.9	3.0	5.1	4.8	3.2	3.8	0.9	2.2	3.6	2.9	2.5	5.9	8.4	5.1	5.8	9.1	2.6	2.6	0.2
1965	0.3	0.0	1.5	1.8	6.1	0.9	1.8	2.1	4.8	1.6	3.5	4.2	3.3	6.8	4.7	3.3	5.6	8.7	2.4	2.4	0.2
1966	0.0	0.3	1.2	4.1	4.9	2.3	3.6	2.1	1.7	1.5	4.6	5.5	5.6	5.7	1.1	11.3	5.4	4.2	2.7	2.6	0.2
1967	0.3	0.3	0.3	1.6	4.6	4.0	2.7	0.9	2.9	3.0	1.7	3.3	5.5	4.6	6.7	7.7	2.6	0.0	2.3	2.3	0.2
1968	0.3	0.3	0.3	1.9	3.6	3.4	1.3	2.2	4.6	0.5	6.1	4.5	3.0	6.3	2.2	4.5	7.4	3.8	2.3	2.3	0.2
1964-68	0.2	0.2	0.8	2.4	4.8	3.1	2.5	2.2	3.0	1.7	3.9	4.1	4.0	5.8	4.6	6.4	5.3	5.0	2.5	2.4	0.2
1969	0.3	0.3	0.0	4.2	4.7	5.2	1.8	1.8	3.0	2.3	3.8	1.8	4.4	3.5	9.7	4.4	0.0	0.0	2.6	2.5	0.2
1970	0.0	0.5	1.1	5.0	3.8	2.3	1.7	4.4	2.6	2.6	5.4	4.2	5.7	1.7	3.2	4.3	4.5	3.3	2.7	2.6	0.2
1971	0.0	1.0	1.0	1.7	2.7	1.8	2.1	4.8	3.9	1.3	1.0	3.5	3.5	3.3	11.3	4.2	8.7	3.1	2.3	2.2	0.2
1972	0.3	0.3	2.1	2.2	6.5	3.3	3.7	3.9	0.9	0.8	3.4	1.7	4.0	3.2	4.0	9.4	4.2	2.9	2.7	2.5	0.2
1973	0.0	0.0	1.5	3.5	4.1	3.8	2.0	2.2	1.7	3.0	0.5	4.0	2.6	2.3	5.9	3.9	6.1	0.0	2.3	2.2	0.2
1969-73	0.1	0.4	1.2	3.3	4.3	3.2	2.3	3.4	2.4	2.0	2.7	3.0	4.0	2.8	6.8	5.3	4.8	1.9	2.5	2.4	0.2
1974	0.3	0.6	1.5	1.8	3.1	4.2	3.7	2.1	3.0	2.5	3.1	3.4	0.6	8.3	5.7	5.1	2.0	2.6	2.5	2.4	0.2
1975	0.0	0.0	0.8	3.9	4.9	2.3	3.9	1.3	2.6	2.1	3.0	0.6	2.4	5.9	1.9	7.5	5.8	5.1	2.4	2.2	0.2
1976	0.0	0.3	1.3	2.3	4.0	3.9	3.4	1.2	2.6	3.0	0.8	3.7	1.8	3.6	2.7	2.4	0.0	0.0	2.1	2.0	0.2
1977	0.0	0.6	1.3	2.0	5.7	3.9	2.6	2.0	2.1	1.3	2.1	3.0	3.5	4.1	1.7	2.4	5.4	4.6	2.4	2.2	0.2
1978	0.0	0.0	0.8	3.0	3.6	3.3	1.2	1.5	1.3	1.7	3.4	2.8	1.7	3.3	5.9	3.4	8.7	4.5	2.1	1.9	0.2
1974-78	0.1	0.3	1.1	2.6	4.3	3.5	2.9	1.6	2.3	2.1	2.5	2.7	2.0	5.0	3.6	4.1	4.5	3.4	2.3	2.2	0.2
1979	0.7	0.6	1.4	2.5	2.3	3.8	2.9	2.9	2.1	0.4	1.3	3.1	4.0	1.9	13.8	4.4	1.7	2.1	2.4	2.2	0.2
1980	0.0	0.3	0.9	3.2	3.8	4.5	2.5	1.8	2.5	3.9	1.7	1.3	4.5	4.3	2.4	3.2	3.2	8.1	2.4	2.3	0.2
1981	0.3	0.0	0.9	2.0	3.0	2.1	2.7	1.7	4.1	1.7	1.7	1.7	3.2	4.9	2.3	2.1	11.0	1.9	2.1	1.9	0.2
1982	0.0	0.3	2.8	2.1	4.7	5.0	3.3	3.5	2.4	3.9	1.3	1.3	4.5	5.4	3.0	6.0	7.6	7.4	2.9	2.7	0.2
1983	0.0	0.3	0.6	2.2	2.9	4.1	3.0	3.7	0.8	2.6	1.7	3.0	3.3	3.6	5.7	3.9	1.5	3.5	2.2	2.1	0.2
1979-83	0.2	0.3	1.3	2.4	3.4	3.9	2.9	2.8	2.3	2.5	1.5	2.1	3.9	4.1	5.4	3.9	5.0	4.6	2.4	2.2	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

OTHER MALIGNANT NEOPLASM OF LYMPHOID AND HISTIOCYTIC TISSUE (ICD9 202)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	5	1	1	2	2	2	3	3	3	5	6	8	9	6	4	8	3	3	0	74
1965	2	0	0	2	0	4	0	3	1	2	10	12	3	6	7	3	2	2	0	59
1966	0	0	0	2	1	0	3	2	8	7	1	6	3	8	5	2	2	1	0	51
1967	1	0	0	2	0	4	0	6	4	2	3	5	3	14	12	7	2	3	0	68
1968	5	0	2	3	2	1	4	0	1	5	8	4	6	6	7	6	5	2	2	69
1964-68	13	1	3	11	5	11	10	14	17	21	28	35	24	40	35	26	14	11	2	321
1969	2	0	1	0	3	3	2	3	4	7	2	8	2	7	11	9	3	4	0	71
1970	0	0	1	1	2	2	0	3	3	5	4	7	5	11	11	6	4	1	0	66
1971	1	1	0	3	4	5	3	2	6	6	6	8	5	7	10	5	2	1	0	75
1972	2	1	0	3	4	0	1	5	7	8	7	10	12	10	6	5	3	9	0	84
1973	3	1	2	2	2	3	3	3	7	9	9	15	8	14	9	5	2	2	1	100
1969-73	8	3	4	9	15	13	9	16	27	35	28	48	32	49	47	30	14	8	1	396
1974	0	1	2	1	2	5	4	8	5	12	13	13	11	7	7	8	6	8	5	118
1975	1	0	2	3	5	4	4	4	5	10	10	3	5	10	12	11	5	2	0	96
1976	2	2	1	7	8	8	5	6	7	10	15	13	8	8	14	4	5	8	0	131
1977	2	1	1	0	3	4	6	9	4	13	5	9	10	17	9	9	2	2	1	107
1978	0	0	1	6	0	3	4	8	8	13	7	12	18	23	12	10	11	6	0	142
1974-78	5	4	7	17	18	24	23	35	29	58	50	50	52	65	54	42	29	26	6	594
1979	2	0	1	0	4	6	6	10	9	9	17	19	31	20	17	15	7	6	1	180
1980	3	1	3	1	4	7	2	13	13	19	19	28	26	34	20	15	11	5	1	225
1981	3	2	3	5	4	4	6	10	12	14	14	21	28	23	22	15	9	7	3	205
1982	4	4	1	3	5	0	7	14	17	15	18	35	24	24	18	25	13	3	0	230
1983	3	1	2	2	4	5	10	12	13	15	23	31	27	27	32	21	15	11	2	256
1979-83	15	8	10	11	21	22	31	59	64	72	91	134	136	128	109	91	55	32	7	1096

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

OTHER MALIGNANT NEOPLASM OF LYMPHOID AND HISTIOCYTIC TISSUE (ICD9 202)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	1.3	0.3	0.3	0.7	0.9	1.0	1.3	1.3	1.4	2.7	3.5	5.7	7.9	6.7	5.9	17.6	11.8	22.1	2.2	2.2	0.2
1965	0.5	0.0	0.0	0.7	0.0	1.9	0.0	1.3	0.4	1.0	5.7	8.4	2.6	6.6	10.1	6.5	7.7	14.4	1.7	1.7	0.2
1966	0.0	0.0	0.0	0.7	0.4	0.0	1.3	0.8	3.4	3.6	0.6	4.1	2.5	8.6	7.1	4.3	7.7	7.1	1.4	1.5	0.2
1967	0.3	0.0	0.0	0.6	0.0	1.8	0.0	2.5	1.7	1.0	1.6	3.3	2.4	14.7	16.9	14.7	7.6	20.7	1.9	1.9	0.2
1968	1.4	0.0	0.5	0.9	0.7	0.4	1.7	0.0	0.4	2.4	4.4	2.5	4.8	6.2	9.8	12.6	18.7	13.3	1.9	1.8	0.2
1964-68	0.7	0.1	0.2	0.7	0.4	1.0	0.9	1.2	1.5	2.1	3.2	4.7	4.0	8.6	10.0	11.1	10.7	15.5	1.8	1.8	0.2
1969	0.6	0.0	0.3	0.0	1.0	1.2	0.9	1.3	1.7	3.2	1.1	4.9	1.6	7.0	15.3	18.7	11.0	25.5	1.9	1.9	0.2
1970	0.0	0.0	0.3	0.3	0.6	0.7	0.0	1.3	1.2	2.2	2.2	4.2	3.8	10.7	15.1	12.4	14.4	6.1	1.8	1.7	0.2
1971	0.3	0.2	0.0	0.8	1.2	1.7	1.2	0.8	2.5	2.6	3.2	4.7	3.7	6.6	13.4	10.2	7.1	5.8	1.9	1.9	0.2
1972	0.6	0.3	0.0	0.8	1.2	0.0	0.4	2.1	2.9	3.4	3.5	5.9	8.5	9.3	7.8	10.1	10.5	0.0	2.1	2.1	0.2
1973	0.9	0.3	0.5	0.5	0.6	0.9	1.2	1.3	2.9	3.8	4.3	8.8	5.5	12.7	11.3	10.0	6.9	11.0	2.5	2.5	0.3
1969-73	0.5	0.2	0.2	0.5	0.9	0.9	0.7	1.3	2.2	3.1	2.9	5.7	4.7	9.3	12.5	12.3	9.9	9.4	2.1	2.0	0.2
1974	0.0	0.3	0.5	0.3	0.6	1.5	1.5	3.3	2.0	5.1	5.9	7.7	7.3	6.2	8.5	15.8	20.5	43.2	2.8	2.7	0.3
1975	0.3	0.0	0.5	0.7	1.4	1.2	1.4	1.6	2.1	4.2	4.5	1.7	3.2	8.6	14.2	21.0	17.1	10.5	2.3	2.2	0.2
1976	0.6	0.6	0.2	1.7	2.2	2.2	1.7	2.4	2.9	4.2	6.6	7.3	5.1	6.6	16.2	7.4	16.8	41.2	3.1	3.0	0.3
1977	0.6	0.3	0.3	0.0	0.8	1.1	1.9	3.5	1.7	5.4	2.2	4.8	6.3	13.5	10.2	15.9	6.6	10.2	2.4	2.4	0.3
1978	0.0	0.0	0.3	1.4	0.0	0.9	1.2	3.0	3.4	5.4	3.1	6.0	11.4	17.7	13.2	17.0	35.7	30.0	3.2	3.1	0.3
1974-78	0.3	0.2	0.3	0.8	1.0	1.4	1.5	2.8	2.4	4.9	4.4	5.5	6.7	10.7	12.5	15.4	19.5	26.9	2.8	2.7	0.3
1979	0.6	0.0	0.3	0.0	1.0	1.7	1.8	3.6	3.7	3.7	7.4	9.1	19.8	14.9	18.1	24.5	22.0	29.3	3.9	3.9	0.4
1980	1.0	0.3	0.8	0.2	1.0	2.0	0.6	4.6	5.3	8.0	8.2	13.2	16.1	24.5	20.6	23.8	33.2	24.2	4.8	4.8	0.5
1981	1.0	0.6	0.9	1.2	1.0	1.1	1.7	3.4	4.9	5.9	5.9	9.7	16.6	16.3	21.8	23.2	26.4	33.0	4.3	4.3	0.5
1982	1.3	1.3	0.3	0.7	1.2	0.0	2.0	4.5	6.7	6.4	7.6	16.0	13.5	16.8	17.2	37.6	36.4	13.9	4.8	4.7	0.5
1983	1.0	0.3	0.6	0.5	1.0	1.3	2.8	3.7	4.9	6.4	9.7	14.0	14.4	19.0	29.3	30.7	39.7	49.3	5.1	5.0	0.5
1979-83	1.0	0.5	0.6	0.5	1.1	1.2	1.8	4.0	5.1	6.1	7.8	12.5	15.0	18.3	21.6	28.1	31.9	30.1	4.6	4.6	0.5

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

OTHER MALIGNANT NEOPLASM OF LYMPHOID AND HISTIOCYTIC TISSUE (ICD9 202)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	3	0	0	0	0	1	4	2	6	1	3	1	5	5	5	5	4	1	0	46
1965	1	0	0	0	0	2	1	1	5	3	3	5	6	6	5	2	3	1	0	44
1966	2	0	0	0	0	3	2	0	2	5	6	2	8	10	2	6	3	3	0	54
1967	0	0	1	0	4	1	0	2	6	2	1	7	4	7	6	5	2	4	0	52
1968	2	0	1	1	1	0	3	4	4	2	4	4	7	2	7	3	5	2	0	52
1964-68	8	0	2	1	5	7	10	9	23	13	17	19	30	30	25	21	17	11	0	248
1969	0	0	1	0	0	0	0	1	1	4	4	7	4	8	6	6	4	6	0	52
1970	0	0	0	0	0	1	0	3	2	5	6	0	4	4	5	5	3	5	0	43
1971	1	0	1	1	2	0	1	0	2	2	3	6	10	8	8	6	6	3	0	60
1972	1	0	3	6	2	1	3	4	2	1	4	6	6	10	8	9	7	7	0	80
1973	0	1	1	2	5	3	2	4	4	5	10	6	7	5	7	6	6	2	0	76
1969-73	2	1	6	9	9	5	6	12	11	17	27	25	31	35	34	32	26	23	0	311
1974	0	1	1	3	1	2	1	3	5	8	8	8	14	12	12	10	8	11	0	106
1975	1	1	0	1	1	1	4	4	4	4	9	4	8	16	7	8	6	8	1	88
1976	2	0	0	0	6	7	2	6	8	8	10	16	12	13	9	7	8	9	1	124
1977	0	1	0	1	3	7	3	6	4	11	9	16	12	11	24	13	7	6	0	134
1978	1	0	1	0	2	2	2	2	9	5	6	10	13	16	16	13	13	9	0	120
1974-78	4	3	2	5	13	19	12	21	30	36	42	54	59	68	68	51	42	43	2	574
1979	0	0	0	2	1	3	3	5	8	12	9	16	22	20	15	16	13	10	3	158
1980	3	0	1	1	0	5	5	5	14	12	9	27	18	18	19	15	16	9	2	179
1981	2	0	0	1	2	4	9	8	8	17	17	16	20	24	30	20	13	7	1	199
1982	2	1	0	0	5	9	7	10	13	11	26	19	38	21	28	13	19	13	1	236
1983	1	1	1	1	3	4	8	10	9	9	25	28	27	21	33	18	11	14	1	225
1979-83	8	2	2	5	11	25	32	38	52	61	86	106	125	104	125	82	72	53	8	997

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

OTHER MALIGNANT NEOPLASM OF LYMPHOID AND HISTIOCYTIC TISSUE (ICD9 202)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.8	0.0	0.0	0.0	0.0	0.5	1.8	0.9	2.7	0.5	1.8	0.7	4.2	4.9	6.0	8.5	11.6	4.5	1.4	1.2	0.1
1965	0.3	0.0	0.0	0.0	0.0	0.9	0.5	0.4	2.2	1.6	1.8	3.5	5.0	5.8	5.8	3.3	8.4	4.3	1.3	1.2	0.1
1966	0.6	0.0	0.0	0.0	0.0	1.4	0.9	0.0	0.8	2.6	3.4	1.4	6.4	9.4	2.3	9.7	8.1	12.7	1.6	1.4	0.1
1967	0.0	0.0	0.3	0.0	1.5	0.4	0.0	0.9	2.5	1.0	0.6	4.6	3.1	6.5	6.7	7.7	5.1	15.9	1.4	1.3	0.1
1968	0.6	0.0	0.3	0.3	0.4	0.0	1.3	1.7	1.7	0.9	2.2	2.5	5.3	1.8	7.7	4.5	12.3	7.5	1.4	1.3	0.1
1964-68	0.4	0.0	0.1	0.1	0.4	0.6	0.9	0.8	2.0	1.3	2.0	2.6	4.8	5.6	5.7	6.7	9.1	9.1	1.4	1.3	0.1
1969	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.4	1.8	2.2	4.3	2.9	7.0	6.5	8.7	5.4	21.2	1.4	1.2	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.3	0.9	2.2	3.2	0.0	2.9	3.4	5.3	7.1	6.8	16.5	1.1	1.0	0.1
1971	0.3	0.0	0.3	0.3	0.6	0.0	0.4	0.0	0.9	0.8	1.6	3.5	6.9	6.6	8.2	8.3	13.0	9.2	1.5	1.3	0.2
1972	0.3	0.0	0.8	1.7	0.6	0.3	1.2	1.7	0.9	0.4	2.0	3.4	4.0	8.0	8.1	12.1	14.7	20.4	1.9	1.7	0.2
1973	0.0	0.3	0.3	0.5	1.5	0.9	0.8	1.7	1.7	2.1	4.6	3.4	4.5	3.9	6.9	7.9	12.2	5.6	1.8	1.6	0.2
1969-73	0.1	0.1	0.3	0.5	0.6	0.4	0.5	1.0	0.9	1.5	2.7	2.9	4.3	5.8	7.0	8.8	11.3	14.3	1.5	1.4	0.1
1974	0.0	0.3	0.3	0.8	0.3	0.6	0.4	1.3	2.1	3.4	3.5	4.5	8.8	9.1	11.4	12.8	16.0	29.0	2.5	2.2	0.2
1975	0.3	0.3	0.0	0.3	0.3	0.3	1.4	1.7	1.7	1.7	3.8	2.2	4.8	11.8	6.5	10.0	11.6	20.3	1.9	1.7	0.2
1976	0.7	0.0	0.0	0.0	1.6	1.9	0.7	2.5	3.4	3.4	4.2	8.4	7.1	9.3	8.1	8.5	15.0	21.7	2.7	2.5	0.3
1977	0.0	0.3	0.0	0.3	0.8	2.0	1.0	2.4	1.7	4.7	3.8	7.9	7.0	7.6	20.9	15.3	12.6	13.9	2.9	2.6	0.3
1978	0.3	0.0	0.3	0.0	0.5	0.6	0.6	0.8	3.9	2.1	2.5	4.7	7.5	10.7	13.5	14.9	22.7	20.0	2.5	2.2	0.2
1974-78	0.3	0.2	0.1	0.3	0.7	1.1	0.8	1.7	2.6	3.1	3.6	5.6	7.0	9.7	12.2	12.4	15.7	20.8	2.5	2.2	0.2
1979	0.0	0.0	0.0	0.5	0.3	0.8	0.9	1.8	3.4	5.1	3.9	7.1	12.7	12.9	12.2	17.7	21.8	21.2	3.1	2.8	0.3
1980	1.0	0.0	0.3	0.2	0.0	1.3	1.4	1.8	5.8	5.2	3.9	11.7	10.1	11.2	15.1	16.1	25.9	18.3	3.5	3.2	0.3
1981	0.7	0.0	0.0	0.3	0.5	1.1	2.5	2.8	3.3	7.3	7.3	6.9	10.7	14.6	23.0	20.8	20.3	13.6	3.9	3.5	0.4
1982	0.7	0.3	0.0	0.0	1.2	2.4	1.9	3.2	5.2	4.8	11.1	8.1	19.2	12.6	20.8	13.0	29.1	24.1	4.5	4.1	0.5
1983	0.3	0.3	0.3	0.3	0.7	1.0	2.2	3.0	3.4	3.9	10.7	11.9	12.8	12.6	23.6	17.4	16.2	24.6	4.2	3.8	0.4
1979-83	0.5	0.1	0.1	0.3	0.6	1.3	1.8	2.6	4.2	5.3	7.4	9.2	13.2	12.8	19.1	17.0	22.6	20.5	3.8	3.5	0.4

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MULTIPLE MYELOMA AND IMMUNOPROLIFERATIVE NEOPLASMS (ICD9 203)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	1	0	4	6	9	7	18	10	21	16	7	1	0	100
1965	0	0	0	0	0	0	0	2	2	2	7	6	15	25	10	14	6	3	0	92
1966	0	0	0	0	0	0	2	0	3	7	6	7	14	14	12	8	4	3	0	80
1967	0	0	0	0	0	0	0	3	2	6	7	14	17	15	10	18	7	1	0	100
1968	0	0	0	0	0	0	0	1	4	1	6	10	15	14	22	16	4	7	0	100
1964-68	0	0	0	0	0	0	3	6	15	22	35	44	79	78	75	72	28	15	0	472
1969	0	0	0	0	1	0	0	1	4	8	13	10	16	19	28	10	10	7	0	127
1970	0	0	0	0	0	0	1	1	2	3	7	12	12	14	19	7	13	2	0	93
1971	0	0	0	0	0	0	2	2	3	7	5	9	11	15	17	12	10	7	0	100
1972	0	0	0	0	0	0	0	1	4	5	12	20	18	25	22	25	8	6	0	146
1973	1	0	1	1	0	0	1	2	6	4	7	12	24	11	18	25	13	5	0	131
1969-73	1	0	1	1	1	0	4	7	19	27	44	63	81	84	104	79	54	27	0	597
1974	0	1	0	1	1	0	0	0	4	8	7	10	19	19	27	17	8	4	1	127
1975	0	0	0	0	0	1	1	1	2	6	13	8	13	25	24	20	9	6	2	131
1976	0	0	0	2	0	0	0	2	4	6	9	26	19	24	32	22	13	7	2	168
1977	0	0	1	0	0	2	0	1	4	11	6	24	33	27	26	15	11	9	0	170
1978	0	0	0	1	0	1	1	0	2	8	14	21	25	32	31	27	14	4	0	181
1974-78	0	1	1	4	1	4	2	4	16	39	49	89	109	127	140	101	55	30	5	777
1979	0	0	0	0	0	0	0	2	2	8	19	9	20	32	20	31	16	16	1	176
1980	0	0	0	0	0	0	0	3	3	9	15	15	25	24	21	23	23	16	0	177
1981	0	0	0	0	1	1	0	2	3	4	9	23	31	34	37	22	16	17	1	201
1982	0	0	0	0	0	0	1	2	5	4	15	18	29	33	20	30	19	13	1	190
1983	0	0	0	0	0	0	0	2	0	7	17	20	26	39	46	35	27	15	1	235
1979-83	0	0	0	0	1	1	1	11	13	32	75	85	131	162	144	141	101	77	4	979

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MULTIPLE MYELOMA AND IMMUNOPROLIFERATIVE NEOPLASMS (ICD9 203)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.8	3.2	5.3	5.0	15.8	11.1	30.7	35.2	27.6	7.4	3.0	2.9	0.4
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.9	1.0	4.0	4.2	12.9	27.3	14.4	30.5	23.2	21.6	2.7	2.7	0.3
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	1.3	3.6	3.4	4.8	11.7	15.0	17.0	17.2	15.4	21.4	2.3	2.3	0.3
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	0.8	3.0	3.8	9.2	13.8	15.8	14.1	37.9	26.5	6.9	2.8	2.7	0.3
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	0.5	3.3	6.4	11.9	14.4	30.9	33.5	14.9	46.7	2.8	2.7	0.3
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.5	1.3	2.2	3.9	6.0	13.2	16.7	21.4	30.9	21.5	21.1	2.7	2.7	0.3
1969	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	1.7	3.7	7.1	6.2	12.4	19.0	39.0	20.8	36.8	44.6	3.5	3.4	0.4
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.8	1.3	3.8	7.2	9.1	13.6	26.1	14.4	46.8	12.2	2.5	2.4	0.3
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.8	1.2	3.0	2.6	5.3	8.1	14.1	22.8	24.5	35.5	40.5	2.6	2.5	0.3
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.6	2.1	6.0	11.7	12.8	23.1	28.7	50.6	27.9	33.5	3.8	3.6	0.4
1973	0.3	0.0	0.2	0.3	0.0	0.0	0.4	0.8	2.5	1.7	3.4	7.1	16.5	10.0	22.6	50.2	44.7	27.5	3.3	3.1	0.3
1969-73	0.1	0.0	0.0	0.1	0.1	0.0	0.3	0.6	1.6	2.4	4.6	7.5	11.8	15.9	27.7	32.3	38.3	31.6	3.1	3.0	0.4
1974	0.0	0.3	0.0	0.3	0.3	0.0	0.0	0.0	1.6	3.4	3.2	5.9	12.6	16.8	32.8	33.6	27.4	21.6	3.1	3.0	0.4
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.8	2.5	5.8	4.7	8.4	21.6	28.4	38.2	30.8	31.6	3.1	3.0	0.4
1976	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.8	1.7	2.5	4.0	14.5	12.1	19.9	37.0	40.4	43.8	36.1	3.9	3.7	0.5
1977	0.0	0.0	0.3	0.0	0.0	0.6	0.0	0.4	1.7	4.6	2.6	12.7	20.9	21.4	29.4	26.5	36.4	45.7	3.9	3.9	0.5
1978	0.0	0.0	0.0	0.2	0.0	0.3	0.3	0.0	0.8	3.3	6.1	10.6	15.9	24.6	34.2	45.8	45.5	20.0	4.0	3.9	0.5
1974-78	0.0	0.1	0.0	0.2	0.1	0.2	0.1	0.3	1.3	3.3	4.4	9.8	14.0	21.0	32.4	37.0	36.9	31.1	3.6	3.5	0.4
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.8	3.3	8.3	4.3	12.8	23.8	21.3	50.7	50.3	78.0	3.8	3.7	0.4
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	1.2	3.8	6.4	7.0	15.5	17.3	21.6	36.6	69.5	77.3	3.8	3.6	0.4
1981	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.7	1.2	1.7	3.8	10.7	18.4	24.1	36.7	34.0	46.9	80.2	4.1	4.0	0.5
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.6	2.0	1.7	6.3	8.2	16.3	23.1	19.1	45.1	53.2	60.2	3.8	3.7	0.4
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	3.0	7.2	9.0	13.9	27.4	42.2	51.1	71.4	67.3	4.6	4.4	0.5
1979-83	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.7	1.0	2.7	6.4	7.9	15.4	23.2	28.5	43.5	58.6	72.4	4.1	3.9	0.4

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MULTIPLE MYELOMA AND IMMUNOPROLIFERATIVE NEOPLASMS (ICD9 203)

YEAR OF DIAGNOSIS	AGE (YEARS)																			ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	1	0	3	5	8	9	15	12	12	9	3	0	0	77
1965	0	0	0	0	0	0	1	1	3	3	2	9	15	5	10	8	3	2	0	62
1966	0	0	0	0	0	0	0	1	1	6	4	9	5	9	10	11	4	1	0	61
1967	0	0	0	0	0	0	0	0	3	4	4	7	8	13	17	12	3	3	0	74
1968	0	0	0	0	0	0	0	0	1	3	14	10	14	11	16	7	10	7	0	93
1964-68	0	0	0	0	0	0	2	2	11	21	32	44	57	50	65	47	23	13	0	367
1969	0	0	0	0	0	0	0	0	3	4	11	12	14	22	12	9	11	6	0	104
1970	0	0	0	0	0	0	0	0	1	6	3	11	18	19	12	13	8	3	0	94
1971	0	0	0	0	0	0	0	1	5	2	7	10	14	14	22	15	9	10	0	109
1972	0	0	0	0	0	0	1	0	4	3	7	14	12	19	17	25	8	8	0	118
1973	0	0	0	0	0	1	0	1	4	2	5	15	13	25	20	26	11	5	0	128
1969-73	0	0	0	0	0	1	1	2	17	17	33	62	71	99	83	88	47	32	0	553
1974	0	0	0	0	0	0	3	2	2	7	9	12	16	26	20	16	17	12	0	142
1975	0	0	0	0	0	0	1	1	2	9	14	8	18	17	20	17	24	9	0	140
1976	0	0	0	1	0	1	1	2	3	11	11	12	21	13	20	23	15	8	0	142
1977	0	0	0	2	0	0	1	1	2	2	11	12	21	26	30	17	12	8	0	145
1978	0	0	0	0	0	0	0	1	4	9	9	13	26	24	26	32	24	17	0	185
1974-78	0	0	0	3	0	1	6	7	13	38	54	57	102	106	116	105	92	54	0	754
1979	0	0	0	0	0	0	1	0	3	4	9	19	25	22	24	22	27	13	1	170
1980	0	0	0	0	0	0	2	1	4	7	7	18	21	32	31	32	30	16	0	201
1981	0	0	0	0	0	0	1	2	3	2	9	20	17	24	31	21	19	14	1	164
1982	0	0	0	0	0	0	0	0	5	7	7	15	15	18	35	27	23	20	0	172
1983	0	0	0	0	0	0	0	1	4	3	13	11	20	25	44	26	23	22	1	193
1979-83	0	0	0	0	0	0	4	4	19	23	45	83	98	121	165	128	122	85	3	900

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MULTIPLE MYELOMA AND IMMUNOPROLIFERATIVE NEOPLASMS (ICD9 203)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	1.4	2.7	4.9	6.5	12.7	11.8	14.5	15.4	8.7	0.0	2.3	2.1	0.3
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.4	1.3	1.6	1.2	6.3	12.4	4.8	11.7	13.2	8.4	8.7	1.8	1.6	0.2
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	3.1	2.3	6.2	4.0	8.5	11.3	17.8	10.8	4.2	1.7	1.5	0.2
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	2.0	2.2	4.6	6.2	12.0	18.9	18.5	7.7	12.0	2.0	1.8	0.2
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.4	7.7	6.4	10.6	9.9	17.5	10.4	24.6	26.4	2.5	2.2	0.3
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.9	2.1	3.7	6.0	9.1	9.4	14.8	15.0	12.3	10.8	2.1	1.8	0.2
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	1.8	6.0	7.4	10.3	19.2	15.0	13.1	25.9	21.2	2.7	2.4	0.3
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.6	1.6	6.5	12.9	16.1	12.7	18.4	18.1	9.9	2.4	2.1	0.3
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.2	0.8	3.6	5.8	9.7	11.5	22.7	20.8	19.5	30.8	2.7	2.3	0.3
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.7	1.3	3.4	8.0	8.1	15.2	17.2	33.6	16.8	23.3	2.8	2.4	0.3
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	1.7	0.8	2.3	8.6	8.4	19.4	19.6	34.2	22.4	13.9	3.0	2.5	0.3
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.2	1.5	1.5	3.4	7.3	9.8	16.3	17.1	24.3	20.5	19.8	2.7	2.3	0.3
1974	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.9	0.9	3.0	4.0	6.8	10.0	19.6	19.1	20.5	33.9	31.7	3.2	2.7	0.3
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.9	3.8	6.0	4.4	10.8	12.6	18.5	21.2	46.5	22.8	3.1	2.5	0.3
1976	0.0	0.0	0.0	0.3	0.0	0.3	0.3	0.8	1.3	4.7	4.6	6.3	12.4	9.3	18.0	27.8	28.1	19.3	3.1	2.6	0.3
1977	0.0	0.0	0.0	0.5	0.0	0.0	0.3	0.4	0.9	0.9	4.7	5.9	12.3	17.9	26.1	20.1	21.7	18.5	3.0	2.6	0.3
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	3.9	3.8	6.1	15.1	16.0	21.9	36.6	42.0	37.9	3.7	3.1	0.3
1974-78	0.0	0.0	0.0	0.2	0.0	0.1	0.4	0.6	1.1	3.2	4.6	5.9	12.2	15.1	20.8	25.4	34.4	26.1	3.2	2.7	0.3
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	1.3	1.7	3.9	8.4	14.4	14.2	19.5	24.4	45.3	27.6	3.3	2.7	0.3
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.4	1.7	3.0	3.0	7.8	11.7	19.9	24.6	34.3	48.5	32.6	3.8	3.1	0.4
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.7	1.2	0.9	3.9	8.6	9.1	14.6	23.8	21.8	29.7	27.2	3.0	2.5	0.3
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.0	3.0	3.0	6.4	7.6	10.8	26.0	27.0	35.2	37.0	3.1	2.5	0.3
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	1.5	1.3	5.6	4.7	9.5	15.0	31.5	25.1	33.8	38.7	3.3	2.7	0.3
1979-83	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.3	1.5	2.0	3.9	7.2	10.3	14.9	25.3	26.5	38.3	32.9	3.3	2.7	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

LYMPHOID LEUKAEMIA (ICD9 204)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	11	5	7	3	0	1	1	1	4	3	9	11	18	19	15	23	7	7	0	145
1965	11	2	5	4	1	1	0	1	8	5	11	6	14	13	14	15	6	6	0	123
1966	12	7	0	1	0	2	2	1	4	3	2	13	8	12	28	11	11	6	0	123
1967	14	3	1	4	2	1	2	0	3	7	5	4	11	10	18	18	6	2	0	111
1968	17	8	3	0	3	0	0	2	0	6	8	11	16	19	14	14	13	6	0	140
1964-68	65	25	16	12	6	5	5	5	19	24	35	45	67	73	89	81	43	27	0	642
1969	13	12	8	4	3	0	1	1	3	6	5	7	18	18	18	16	10	4	0	147
1970	11	4	3	6	1	2	4	2	4	9	5	13	12	25	18	7	7	10	0	143
1971	17	6	5	5	2	4	1	1	6	8	8	9	12	14	14	11	13	16	0	152
1972	19	12	6	6	1	2	2	1	3	4	7	14	16	20	21	11	14	11	0	170
1973	25	13	6	7	6	2	7	2	5	7	13	8	15	18	12	15	5	9	0	175
1969-73	85	47	28	28	13	10	15	7	21	34	38	51	73	95	83	60	49	50	0	787
1974	25	10	11	8	5	2	2	6	1	6	7	10	16	19	26	12	13	7	0	186
1975	28	6	8	14	2	1	4	0	2	2	16	5	17	22	26	15	10	5	1	184
1976	20	12	5	8	1	2	1	0	5	3	10	16	21	20	19	11	12	5	0	171
1977	23	8	5	6	3	3	1	0	2	3	7	12	16	17	23	18	14	11	0	172
1978	15	13	7	6	5	4	3	5	1	3	15	20	30	35	45	30	20	18	1	276
1974-78	111	49	36	42	16	12	11	11	11	17	55	63	100	113	139	86	69	46	2	989
1979	23	14	14	11	2	2	0	1	7	6	9	22	24	27	38	35	28	16	1	280
1980	25	11	6	10	0	2	6	3	2	5	16	21	21	36	35	28	25	15	0	267
1981	21	14	8	6	1	6	1	3	4	4	13	19	31	29	33	34	15	15	1	258
1982	30	18	8	2	7	4	1	3	3	9	16	14	22	34	28	26	15	17	0	257
1983	21	16	5	8	3	2	3	2	4	9	12	21	32	34	33	31	20	20	0	276
1979-83	120	73	41	37	13	16	11	12	20	33	66	97	130	160	167	154	103	83	2	1338

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

LYMPHOID LEUKAEMIA (ICD9 204)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	2.8	1.3	2.1	1.1	0.0	0.5	0.4	0.4	1.8	1.6	5.3	7.9	15.8	21.1	22.0	50.7	27.6	51.5	4.4	4.2	0.4
1965	2.8	0.5	1.5	1.4	0.4	0.5	0.0	0.4	3.5	2.6	6.3	4.2	12.0	14.2	20.2	32.7	23.2	43.2	3.6	3.6	0.4
1966	3.1	1.8	0.0	0.3	0.0	0.9	0.9	0.4	1.7	1.5	1.1	8.8	6.7	12.9	39.8	23.6	42.3	42.9	3.6	3.5	0.4
1967	3.8	0.8	0.3	1.3	0.8	0.4	0.9	0.0	1.3	3.5	2.7	2.6	9.0	10.5	25.4	37.9	22.7	13.8	3.2	3.1	0.3
1968	4.7	2.0	0.8	0.0	1.1	0.0	0.0	0.8	0.0	2.9	4.4	7.0	12.7	19.5	19.7	29.3	48.5	40.0	3.9	3.9	0.4
1964-68	3.4	1.3	0.9	0.8	0.5	0.5	0.4	0.4	1.6	2.4	3.9	6.1	11.2	15.6	25.4	34.7	33.0	38.0	3.7	3.6	0.4
1969	3.8	3.0	2.1	1.2	1.0	0.0	0.4	0.4	1.2	2.8	2.7	4.3	14.0	18.0	25.1	33.3	36.8	25.5	4.0	4.0	0.4
1970	3.3	1.0	0.8	1.7	0.3	0.7	1.7	0.8	1.7	4.0	2.7	7.8	9.1	24.3	24.7	14.4	25.2	61.0	3.8	3.9	0.4
1971	5.2	1.5	1.2	1.4	0.6	1.4	0.4	0.4	2.5	3.4	4.2	5.3	8.8	13.2	18.8	22.5	46.1	92.5	4.0	4.0	0.3
1972	5.9	3.1	1.5	1.6	0.3	0.7	0.8	0.4	1.2	1.7	3.5	8.2	11.3	18.5	27.4	22.3	48.8	61.5	4.4	4.5	0.4
1973	7.7	3.5	1.4	1.8	1.8	0.6	2.7	0.8	2.1	3.0	6.2	4.7	10.3	16.3	15.1	30.1	17.2	49.5	4.5	4.6	0.4
1969-73	5.1	2.4	1.4	1.5	0.8	0.7	1.2	0.6	1.7	3.0	3.9	6.1	10.7	18.0	22.1	24.5	34.8	58.5	4.2	4.2	0.4
1974	7.7	2.8	2.6	2.0	1.4	0.6	0.7	2.5	0.4	2.5	3.2	5.9	10.6	16.8	31.6	23.7	44.5	37.8	4.7	4.8	0.5
1975	8.7	1.7	1.9	3.5	0.6	0.3	1.4	0.0	0.8	0.8	7.2	2.9	11.0	19.0	30.8	28.6	34.2	26.3	4.5	4.6	0.5
1976	6.4	3.5	1.2	1.9	0.3	0.6	0.3	0.0	2.1	1.3	4.4	8.9	13.3	16.6	22.0	20.2	40.4	25.8	4.2	4.3	0.4
1977	7.4	2.4	1.3	1.4	0.8	0.9	0.3	0.0	0.8	1.2	3.1	6.4	10.1	13.5	26.0	31.9	46.4	55.8	4.1	4.2	0.4
1978	4.9	3.9	1.8	1.4	1.3	1.1	0.9	1.9	0.4	1.2	6.6	10.1	19.0	26.9	49.6	50.9	64.9	90.0	6.3	6.3	0.7
1974-78	7.1	2.9	1.8	2.1	0.9	0.7	0.7	0.9	0.9	1.4	4.9	6.9	12.8	18.7	32.2	31.5	46.3	47.6	4.8	4.9	0.5
1979	7.5	4.2	3.8	2.6	0.5	0.6	0.0	0.4	2.9	2.5	3.9	10.6	15.3	20.1	40.6	57.2	88.1	78.0	6.4	6.4	0.6
1980	8.2	3.4	1.7	2.4	0.0	0.6	1.7	1.1	0.8	2.1	6.9	9.9	13.0	25.9	36.1	44.5	75.5	72.5	6.0	6.0	0.6
1981	6.9	4.4	2.3	1.5	0.3	1.7	0.3	1.0	1.6	1.7	5.5	8.8	18.4	20.6	32.8	52.6	44.0	70.8	5.7	5.8	0.5
1982	9.8	5.8	2.3	0.5	1.7	1.1	0.3	1.0	1.2	3.8	6.8	6.4	12.4	23.8	26.8	39.1	42.0	78.7	5.7	5.9	0.5
1983	6.8	5.2	1.5	2.1	0.7	0.5	0.9	0.6	1.5	3.8	5.1	9.5	17.0	23.9	30.2	45.3	52.9	89.7	5.9	6.0	0.5
1979-83	7.8	4.6	2.3	1.8	0.7	0.9	0.6	0.8	1.6	2.8	5.6	9.0	15.3	22.9	33.1	47.6	59.7	78.1	5.9	6.0	0.5

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

LYMPHOID LEUKAEMIA (ICD9 204)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	7	2	2	1	0	1	1	1	3	1	4	8	11	10	16	10	7	0	0	85
1965	4	5	2	1	1	0	1	1	0	0	2	6	4	8	13	11	4	5	0	68
1966	8	4	1	3	0	0	3	1	1	5	3	4	6	14	9	14	4	5	0	85
1967	6	4	2	4	0	0	0	0	3	2	6	6	4	10	8	7	5	3	0	70
1968	12	6	2	0	1	1	1	0	2	3	3	10	4	8	7	7	4	8	0	79
1964-68	37	21	9	9	2	2	6	3	9	11	18	34	29	50	53	49	24	21	0	387
1969	13	8	2	3	1	2	0	1	2	6	3	6	9	11	11	16	10	8	0	112
1970	15	5	1	3	0	0	2	0	2	5	2	6	5	7	8	12	5	4	0	82
1971	13	11	4	1	1	4	1	2	0	4	5	7	9	11	13	9	8	2	0	105
1972	12	9	3	2	3	2	0	1	2	2	3	6	12	16	12	9	5	7	0	106
1973	19	6	6	2	2	0	0	2	2	3	3	4	9	9	13	7	8	11	0	106
1969-73	72	39	16	11	7	8	3	6	8	20	16	29	44	54	57	53	36	32	0	511
1974	10	8	4	5	0	0	0	0	1	3	7	2	6	6	19	11	9	9	0	100
1975	13	9	2	6	2	1	0	0	1	6	7	9	8	14	15	14	3	11	0	121
1976	15	6	7	5	2	0	1	0	1	2	6	13	9	12	16	14	11	14	0	134
1977	13	10	4	3	4	0	1	3	1	2	8	7	17	14	11	9	16	11	0	134
1978	26	14	3	2	2	0	1	0	0	2	6	11	16	15	17	19	10	21	0	165
1974-78	77	47	20	21	10	1	3	3	4	15	34	42	56	61	78	67	49	66	0	654
1979	20	9	5	3	1	1	0	2	2	4	9	9	11	17	18	16	26	18	0	171
1980	23	17	4	1	1	1	3	2	4	4	4	6	12	30	25	32	22	16	0	207
1981	12	12	4	4	2	1	3	6	2	2	2	10	14	20	22	17	19	14	4	170
1982	14	7	3	3	1	3	4	3	6	4	3	9	11	18	28	26	14	18	0	175
1983	17	10	2	3	1	3	1	1	3	1	4	13	15	21	21	24	20	27	0	187
1979-83	86	55	18	14	6	9	11	14	17	15	22	47	63	106	114	115	101	93	4	910

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

LYMPHOID LEUKAEMIA (ICD9 204)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	1.9	0.6	0.6	0.4	0.0	0.5	0.5	0.4	1.4	0.5	2.4	5.8	9.3	9.8	19.3	17.1	20.2	0.0	2.6	2.3	0.3
1965	1.1	1.4	0.6	0.4	0.4	0.0	0.5	0.4	0.0	0.0	1.2	4.2	3.3	7.7	15.2	18.2	11.1	21.7	2.0	1.7	0.2
1966	2.2	1.1	0.3	1.0	0.0	0.0	1.4	0.4	0.4	2.6	1.7	2.7	4.8	13.2	10.2	22.6	10.8	21.1	2.4	2.1	0.2
1967	1.7	1.0	0.6	1.3	0.0	0.0	0.0	0.0	1.2	1.0	3.4	4.0	3.1	9.2	8.9	10.8	12.8	12.0	2.0	1.7	0.2
1968	3.5	1.6	0.6	0.0	0.4	0.4	0.4	0.0	0.8	1.4	1.7	6.4	3.0	7.2	7.7	10.4	9.8	30.2	2.2	2.0	0.2
1964-68	2.1	1.1	0.5	0.6	0.2	0.2	0.5	0.3	0.8	1.1	2.1	4.6	4.6	9.4	12.1	15.7	12.8	17.5	2.2	2.0	0.2
1969	4.0	2.1	0.5	0.9	0.3	0.8	0.0	0.4	0.8	2.7	1.6	3.7	6.6	9.6	11.9	23.3	23.6	28.3	3.0	2.7	0.2
1970	4.7	1.3	0.3	0.9	0.0	0.0	0.9	0.0	0.9	2.2	1.1	3.6	3.6	5.9	8.4	17.0	11.3	13.2	2.1	2.0	0.2
1971	4.2	2.9	1.0	0.3	0.3	1.4	0.4	0.9	0.0	1.7	2.6	4.0	6.3	9.0	13.4	12.5	17.3	6.2	2.7	2.5	0.2
1972	3.9	2.4	0.8	0.6	0.9	0.7	0.0	0.4	0.9	0.8	1.5	3.4	8.1	12.8	12.1	12.1	10.5	20.4	2.6	2.5	0.2
1973	6.2	1.7	1.5	0.5	0.6	0.0	0.0	0.9	0.9	1.3	1.4	2.3	5.8	7.0	12.8	9.2	16.3	30.6	2.6	2.5	0.2
1969-73	4.6	2.1	0.8	0.6	0.4	0.6	0.3	0.5	0.7	1.7	1.6	3.4	6.1	8.9	11.7	14.7	15.7	19.8	2.6	2.4	0.2
1974	3.3	2.4	1.0	1.3	0.0	0.0	0.0	0.0	0.4	1.3	3.1	1.1	3.8	4.5	18.1	14.1	18.0	23.7	2.4	2.1	0.2
1975	4.3	2.7	0.5	1.6	0.5	0.3	0.0	0.0	0.4	2.6	3.0	5.0	4.8	10.4	13.9	17.5	5.8	27.9	2.8	2.7	0.2
1976	5.1	1.8	1.8	1.3	0.5	0.0	0.3	0.0	0.4	0.9	2.5	6.8	5.3	8.5	14.4	16.9	20.6	33.8	3.0	2.8	0.2
1977	4.4	3.1	1.1	0.8	1.0	0.0	0.3	1.2	0.4	0.9	3.4	3.5	9.9	9.6	9.6	10.6	28.9	25.4	3.0	2.8	0.2
1978	8.8	4.4	0.8	0.5	0.5	0.0	0.3	0.0	0.0	0.9	2.5	5.1	9.3	10.0	14.3	21.7	17.5	46.8	3.6	3.6	0.3
1974-78	5.2	2.9	1.0	1.1	0.5	0.1	0.2	0.2	0.3	1.3	2.9	4.4	6.7	8.7	14.0	16.2	18.3	31.9	3.0	2.9	0.2
1979	6.8	2.9	1.4	0.7	0.3	0.3	0.0	0.7	0.8	1.7	3.9	4.0	6.3	11.0	14.6	17.7	43.6	38.2	3.6	3.4	0.3
1980	7.9	5.5	1.2	0.2	0.3	0.3	0.8	0.7	1.7	1.7	1.7	2.6	6.7	18.6	19.8	34.3	35.6	32.6	4.3	4.1	0.3
1981	4.1	4.0	1.2	1.0	0.5	0.3	0.8	2.1	0.8	0.9	0.9	4.3	7.5	12.2	16.9	17.7	29.7	27.2	3.3	3.1	0.3
1982	4.8	2.4	0.9	0.8	0.2	0.8	1.1	1.0	2.4	1.7	1.3	3.9	5.5	10.8	20.8	26.0	21.4	33.3	3.4	3.1	0.3
1983	5.8	3.4	0.6	0.8	0.2	0.8	0.3	0.3	1.1	0.4	1.7	5.5	7.1	12.6	15.0	23.1	29.4	47.5	3.5	3.3	0.3
1979-83	5.9	3.6	1.1	0.7	0.3	0.5	0.6	0.9	1.4	1.3	1.9	4.1	6.6	13.0	17.4	23.8	31.7	36.0	3.6	3.4	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MYELOID LEUKAEMIA (ICD9 205)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	1	0	0	0	1	3	3	9	8	7	12	10	7	5	9	2	1	0	79
1965	2	0	2	1	0	6	2	5	2	5	4	11	6	8	9	4	6	2	0	75
1966	4	2	5	3	2	1	3	3	1	1	3	6	12	7	7	9	2	0	0	71
1967	2	0	3	2	0	1	4	5	8	3	3	4	8	8	7	12	2	2	0	74
1968	6	2	1	0	4	2	0	0	4	3	5	7	3	3	8	5	6	4	0	63
1964-68	15	5	11	6	6	11	12	16	24	20	22	40	39	33	36	39	18	9	0	362
1969	1	2	4	8	2	7	3	7	7	5	8	5	6	11	8	8	9	1	0	102
1970	1	3	2	8	5	4	3	6	7	14	6	8	7	10	7	8	8	8	0	115
1971	3	2	1	2	4	6	8	6	11	5	2	14	5	10	12	9	3	4	0	107
1972	2	4	4	2	3	6	4	6	6	7	14	11	6	15	20	8	8	5	0	131
1973	1	3	2	5	3	8	3	7	5	9	6	5	13	8	12	7	15	7	0	119
1969-73	8	14	13	25	17	31	21	32	36	40	36	43	37	54	59	40	43	25	0	574
1974	2	1	5	3	9	7	7	9	8	7	11	11	15	18	14	15	10	3	0	155
1975	2	5	2	2	4	9	10	2	6	13	11	14	7	13	16	12	5	8	1	142
1976	3	2	9	5	12	1	2	6	9	8	9	14	16	20	10	9	10	7	0	148
1977	5	2	5	2	5	3	7	6	8	9	14	12	18	17	18	17	8	6	1	163
1978	1	0	4	6	6	7	9	5	11	13	11	16	11	20	18	14	5	12	0	169
1974-78	13	10	25	18	36	27	35	28	42	50	56	67	67	88	76	67	38	32	2	777
1979	3	0	1	3	1	6	2	9	9	8	11	13	19	26	26	30	8	8	0	183
1980	3	0	0	3	6	6	10	4	10	10	15	15	18	26	25	22	19	14	0	206
1981	3	1	0	3	4	6	8	4	5	7	11	20	19	28	27	19	16	11	1	193
1982	4	2	3	2	6	3	10	12	7	11	15	16	22	25	36	19	18	9	2	222
1983	3	3	3	1	4	6	11	11	9	10	10	13	21	21	37	23	17	8	1	212
1979-83	16	6	7	12	21	27	41	40	40	46	62	77	99	126	151	113	78	50	4	1016

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MYELOID LEUKAEMIA (ICD9 205)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.3	0.0	0.0	0.0	0.5	1.3	1.3	4.1	4.2	4.1	8.6	8.8	7.8	7.3	19.8	7.9	7.4	2.3	2.3	0.2
1965	0.5	0.0	0.6	0.3	0.0	2.9	0.9	2.1	0.9	2.6	2.3	7.7	5.1	8.7	13.0	8.7	23.2	14.4	2.2	2.2	0.2
1966	1.0	0.5	1.4	1.0	0.8	0.5	1.3	1.3	0.4	0.5	1.7	4.1	10.0	7.5	9.9	19.3	7.7	0.0	2.1	2.0	0.2
1967	0.5	0.0	0.8	0.6	0.0	0.4	1.8	2.1	3.4	1.5	1.6	2.6	6.5	8.4	9.9	25.3	7.6	13.8	2.1	2.0	0.2
1968	1.7	0.5	0.3	0.0	1.4	0.8	0.0	0.0	1.7	1.4	2.7	4.5	2.4	3.1	11.3	10.5	22.4	26.7	1.8	1.7	0.2
1964-68	0.8	0.3	0.6	0.4	0.5	1.0	1.1	1.3	2.1	2.0	2.5	5.4	6.5	7.1	10.3	16.7	13.8	12.7	2.1	2.0	0.2
1969	0.3	0.5	1.0	2.4	0.7	2.8	1.3	2.9	2.9	2.3	4.4	3.1	4.7	11.0	11.1	16.6	33.1	6.4	2.8	2.7	0.3
1970	0.3	0.7	0.5	2.3	1.6	1.5	1.3	2.5	2.9	6.2	3.2	4.8	5.3	9.7	9.6	16.5	28.8	48.8	3.1	3.0	0.3
1971	0.9	0.5	0.2	0.6	1.2	2.1	3.3	2.5	4.5	2.2	1.1	8.2	3.7	9.4	16.1	18.4	10.6	23.1	2.8	2.7	0.3
1972	0.6	1.0	1.0	0.5	0.9	2.0	1.6	2.5	2.5	3.0	7.0	6.4	4.3	13.9	26.1	16.2	27.9	27.9	3.3	3.3	0.4
1973	0.3	0.8	0.5	1.3	0.9	2.5	1.2	3.0	2.1	3.8	2.9	2.9	8.9	7.3	15.1	14.1	51.5	38.5	3.0	2.9	0.3
1969-73	0.5	0.7	0.6	1.4	1.0	2.2	1.7	2.7	3.0	3.5	3.7	5.1	5.4	10.2	15.7	16.3	30.5	29.2	3.0	2.9	0.3
1974	0.6	0.3	1.2	0.8	2.6	2.1	2.5	3.7	3.3	3.0	5.0	6.5	10.0	15.9	17.0	29.6	34.2	16.2	3.8	3.7	0.4
1975	0.6	1.4	0.5	0.5	1.1	2.6	3.5	0.8	2.5	5.5	4.9	8.1	4.5	11.2	19.0	22.9	17.1	42.1	3.4	3.3	0.3
1976	1.0	0.6	2.2	1.2	3.3	0.3	0.7	2.4	3.7	3.3	4.0	7.8	10.2	16.6	11.6	16.5	33.7	15.5	3.6	3.4	0.3
1977	1.6	0.6	1.3	0.5	1.3	0.9	2.2	2.4	3.3	3.7	6.2	6.4	11.4	13.5	20.4	30.1	26.5	30.5	3.8	3.7	0.4
1978	0.3	0.0	1.0	1.4	1.6	2.0	2.7	1.9	4.6	5.4	4.8	8.0	7.0	15.4	19.8	23.8	16.2	60.0	3.8	3.7	0.4
1974-78	0.8	0.6	1.2	0.9	2.0	1.5	2.3	2.2	3.5	4.2	5.0	7.4	8.6	14.5	17.6	24.6	25.5	33.1	3.7	3.6	0.4
1979	1.0	0.0	0.3	0.7	0.3	1.7	0.6	3.3	3.7	3.3	4.8	6.2	12.1	19.3	27.7	49.0	25.2	39.0	4.0	3.9	0.4
1980	1.0	0.0	0.0	0.7	1.5	1.7	2.9	1.4	4.1	4.2	6.4	7.0	11.2	18.7	25.8	35.0	57.4	67.6	4.4	4.3	0.4
1981	1.0	0.3	0.0	0.7	1.0	1.7	2.2	1.4	2.0	3.0	4.7	9.3	11.3	19.8	26.8	29.4	46.9	51.9	4.0	3.9	0.4
1982	1.3	0.6	0.9	0.5	1.5	0.8	2.8	3.8	2.8	4.7	6.3	7.3	12.4	17.5	34.4	28.6	50.4	41.7	4.6	4.4	0.5
1983	1.0	1.0	0.9	0.3	1.0	1.6	3.1	3.4	3.4	4.2	4.2	5.9	11.2	14.8	33.9	33.6	45.0	35.9	4.3	4.1	0.4
1979-83	1.0	0.4	0.4	0.6	1.1	1.5	2.3	2.7	3.2	3.9	5.3	7.2	11.6	18.0	29.9	34.9	45.2	47.0	4.3	4.1	0.4

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MYELOID LEUKAEMIA (ICD9 205)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	4	5	3	5	6	2	8	3	6	7	7	5	6	8	7	7	5	0	95
1965	2	0	0	1	1	2	3	2	4	3	5	7	8	4	5	8	3	5	0	63
1966	2	1	2	2	2	0	2	1	4	3	4	10	4	7	7	12	4	2	0	69
1967	3	1	0	3	1	1	1	3	3	5	4	8	5	9	6	7	2	4	0	66
1968	6	1	4	0	3	1	4	2	3	5	5	4	9	7	8	3	3	0	0	68
1964-68	14	7	11	9	12	10	12	16	17	22	25	36	31	33	34	37	19	16	0	361
1969	2	4	3	3	2	3	4	7	10	7	5	5	10	6	3	8	6	4	0	92
1970	3	1	2	4	0	0	2	1	10	8	3	7	5	11	8	3	7	3	0	78
1971	2	0	3	2	2	3	2	3	3	3	4	6	6	8	10	5	10	5	0	77
1972	3	2	1	3	6	4	9	7	7	6	12	9	12	10	11	9	10	4	0	125
1973	2	1	5	5	6	6	2	3	10	12	12	12	13	11	13	15	6	7	0	141
1969-73	12	8	14	17	16	16	19	21	40	36	36	39	46	46	45	40	39	23	0	513
1974	3	2	1	4	5	4	7	3	6	6	7	12	11	13	6	6	13	9	0	118
1975	6	5	1	4	2	9	3	5	5	5	8	8	3	9	14	10	4	5	0	106
1976	3	2	6	5	7	6	7	4	6	8	8	5	4	5	12	13	10	7	0	118
1977	4	0	2	1	5	5	9	6	7	6	9	8	5	6	15	6	8	8	0	110
1978	2	2	3	1	8	6	5	13	6	11	7	14	12	12	10	20	12	12	0	156
1974-78	18	11	13	15	27	30	31	31	30	36	39	47	35	45	57	55	47	41	0	608
1979	2	0	1	3	6	4	3	10	7	10	7	17	12	18	18	15	14	12	1	160
1980	3	0	2	3	2	4	5	12	4	17	15	18	11	20	19	17	16	25	0	193
1981	3	1	3	3	2	8	7	7	7	9	17	13	12	16	23	18	16	13	0	178
1982	0	3	5	2	2	3	6	11	7	7	11	12	15	19	19	15	18	19	0	174
1983	2	2	2	8	5	12	7	7	8	10	9	14	19	21	24	19	22	10	1	202
1979-83	10	6	13	19	17	31	28	47	33	53	59	74	69	94	103	84	86	79	2	907

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MYELOID LEUKAEMIA (ICD9 205)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	1.1	1.6	1.1	2.3	2.9	0.9	3.4	1.4	3.2	4.3	5.1	4.2	5.9	9.6	11.9	20.2	22.7	2.9	2.6	0.2
1965	0.5	0.0	0.0	0.4	0.4	0.9	1.4	0.9	1.7	1.6	2.9	4.9	6.6	3.9	5.8	13.2	8.4	21.7	1.9	1.7	0.2
1966	0.6	0.3	0.6	0.7	0.8	0.0	0.9	0.4	1.7	1.5	2.3	6.8	3.2	6.6	7.9	19.4	10.8	8.4	2.0	1.7	0.2
1967	0.8	0.3	0.0	1.0	0.4	0.4	0.4	1.3	1.2	2.5	2.2	5.3	3.9	8.3	6.7	10.8	5.1	15.9	1.8	1.7	0.2
1968	1.8	0.3	1.1	0.0	1.1	0.4	1.8	0.9	1.2	2.4	2.8	2.5	6.8	6.3	8.8	4.5	7.4	0.0	1.9	1.8	0.2
1964-68	0.8	0.4	0.7	0.6	1.0	0.9	1.1	1.4	1.5	2.2	2.9	4.9	5.0	6.2	7.8	11.8	10.1	13.3	2.1	1.9	0.2
1969	0.6	1.0	0.8	0.9	0.7	1.2	1.8	3.1	4.2	3.2	2.7	3.1	7.4	5.2	3.2	11.6	14.2	14.1	2.4	2.2	0.2
1970	0.9	0.3	0.5	1.2	0.0	0.0	0.9	0.4	4.3	3.5	1.6	4.2	3.6	9.3	8.4	4.3	15.8	9.9	2.0	1.8	0.2
1971	0.6	0.0	0.8	0.6	0.6	1.1	0.8	1.3	1.3	1.3	2.1	3.5	4.2	6.6	10.3	6.9	21.6	15.4	1.9	1.7	0.2
1972	1.0	0.5	0.3	0.8	1.8	1.3	3.7	3.1	3.0	2.5	5.9	5.2	8.1	8.0	11.1	12.1	21.1	11.7	3.1	2.8	0.3
1973	0.7	0.3	1.3	1.4	1.8	1.9	0.8	1.3	4.3	5.1	5.6	6.9	8.4	8.6	12.8	19.7	12.2	19.5	3.4	3.1	0.3
1969-73	0.8	0.4	0.7	1.0	1.0	1.1	1.6	1.8	3.4	3.1	3.7	4.6	6.4	7.6	9.3	11.1	17.0	14.3	2.6	2.4	0.2
1974	1.0	0.6	0.3	1.1	1.4	1.2	2.6	1.3	2.6	2.5	3.1	6.8	6.9	9.8	5.7	7.7	25.9	23.7	2.8	2.5	0.2
1975	2.0	1.5	0.3	1.0	0.5	2.6	1.1	2.1	2.1	2.1	3.4	4.4	1.8	6.7	13.0	12.5	7.8	12.7	2.4	2.3	0.2
1976	1.0	0.6	1.5	1.3	1.9	1.7	2.4	1.6	2.6	3.4	3.4	2.6	2.4	3.6	10.8	15.7	18.8	16.9	2.7	2.3	0.2
1977	1.4	0.0	0.5	0.3	1.3	1.4	2.9	2.4	3.0	2.6	3.8	4.0	2.9	4.1	13.1	7.1	14.4	18.5	2.4	2.2	0.2
1978	0.7	0.6	0.8	0.2	2.1	1.7	1.5	5.0	2.6	4.7	3.0	6.5	7.0	8.0	8.4	22.9	21.0	26.7	3.3	2.9	0.3
1974-78	1.2	0.7	0.7	0.8	1.4	1.7	2.1	2.5	2.6	3.1	3.3	4.9	4.2	6.4	10.2	13.3	17.6	19.8	2.7	2.5	0.2
1979	0.7	0.0	0.3	0.7	1.5	1.1	0.9	3.7	3.0	4.3	3.0	7.6	6.9	11.6	14.6	16.6	23.5	25.5	3.3	2.9	0.3
1980	1.0	0.0	0.6	0.7	0.5	1.1	1.4	4.3	1.7	7.3	6.4	7.8	6.2	12.4	15.1	18.2	25.9	50.9	3.8	3.4	0.3
1981	1.0	0.3	0.9	0.8	0.5	2.1	1.9	2.4	2.9	3.9	7.3	5.6	6.4	9.7	17.7	18.7	25.0	25.2	3.5	3.1	0.3
1982	0.0	1.0	1.5	0.5	0.5	0.8	1.7	3.5	2.8	3.0	4.7	5.1	7.6	11.4	14.1	15.0	27.5	35.2	3.3	2.9	0.3
1983	0.7	0.7	0.6	2.2	1.2	3.1	1.9	2.1	3.0	4.3	3.9	6.0	9.0	12.6	17.2	18.3	32.3	17.6	3.8	3.4	0.3
1979-83	0.7	0.4	0.8	1.0	0.9	1.6	1.6	3.2	2.7	4.6	5.1	6.4	7.3	11.5	15.8	17.4	27.0	30.5	3.5	3.1	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MONOCYTIC LEUKAEMIA (ICD9 206)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	1	0	1	0	1	0	1	1	1	0	0	8
1965	0	0	0	1	0	0	0	0	0	0	1	0	0	1	1	1	1	1	0	7
1966	0	0	0	0	0	0	0	0	1	1	0	0	0	0	1	4	1	0	0	8
1967	0	0	0	0	0	0	1	0	0	1	0	0	0	1	0	3	2	1	0	9
1968	0	0	0	0	0	0	0	0	0	0	3	1	0	0	2	1	2	0	0	9
1964-68	0	0	0	1	0	0	1	0	2	2	5	1	1	2	7	10	7	2	0	41
1969	0	0	0	0	0	0	0	0	0	1	0	4	0	1	3	1	1	1	0	12
1970	0	0	0	0	1	2	0	1	0	0	2	2	0	1	1	2	1	0	0	13
1971	0	0	0	1	0	0	1	0	0	0	2	1	2	0	2	0	0	0	0	9
1972	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	3
1973	1	0	0	1	0	1	0	0	0	1	0	0	0	1	1	3	1	1	0	11
1969-73	1	0	0	2	1	3	1	1	0	3	4	7	2	3	8	6	4	2	0	48
1974	0	0	1	1	0	0	0	0	0	0	0	1	1	0	0	1	1	2	0	8
1975	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	0	3	1	0	7
1976	0	0	0	0	0	1	0	0	0	0	1	1	2	2	0	2	2	0	0	11
1977	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	2	1	2	0	8
1978	0	0	0	1	0	0	0	0	0	0	0	0	0	2	0	3	2	0	0	8
1974-78	0	0	1	2	0	3	0	0	0	0	2	2	5	4	1	8	9	5	0	42
1979	0	0	0	1	1	0	0	0	0	1	0	0	0	0	1	1	0	5	0	10
1980	1	0	0	0	0	0	0	0	0	0	0	0	1	2	2	1	2	1	0	10
1981	1	0	0	0	2	0	0	0	0	0	0	0	1	2	4	0	0	0	0	10
1982	0	0	0	0	0	0	0	0	1	0	0	1	1	0	4	2	1	2	0	12
1983	0	0	0	0	0	0	0	0	0	1	0	0	4	0	1	1	0	1	0	8
1979-83	2	0	0	1	3	0	0	0	1	2	0	1	7	4	12	5	3	9	0	50

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MONOCYTIC LEUKAEMIA (ICD9 206)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.6	0.0	0.9	0.0	4.4	2.2	3.9	0.0	0.2	0.2	0.0
1965	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	1.1	1.4	2.2	3.9	7.2	0.2	0.2	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.0	0.0	0.0	0.0	1.4	8.6	3.8	0.0	0.2	0.2	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	0.0	0.0	0.0	1.1	0.0	6.3	7.6	6.9	0.3	0.2	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	0.6	0.0	0.0	2.8	2.1	7.5	0.0	0.2	0.2	0.0
1964-68	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.2	0.2	0.6	0.1	0.2	0.4	2.0	4.3	5.4	2.8	0.2	0.2	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	2.5	0.0	1.0	4.2	2.1	3.7	6.4	0.3	0.3	0.0
1970	0.0	0.0	0.0	0.0	0.3	0.7	0.0	0.4	0.0	0.0	1.1	1.2	0.0	1.0	1.4	4.1	3.6	0.0	0.3	0.3	0.0
1971	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	0.0	0.0	1.1	0.6	1.5	0.0	2.7	0.0	0.0	0.0	0.2	0.2	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	1.3	0.0	3.5	0.0	0.1	0.1	0.0
1973	0.3	0.0	0.0	0.3	0.0	0.3	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.9	1.3	6.0	3.4	5.5	0.3	0.3	0.0
1969-73	0.1	0.0	0.0	0.1	0.1	0.2	0.1	0.1	0.0	0.3	0.4	0.8	0.3	0.6	2.1	2.5	2.8	2.3	0.3	0.2	0.0
1974	0.0	0.0	0.2	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	0.0	0.0	2.0	3.4	10.8	0.2	0.2	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.0	0.6	0.0	0.0	0.0	10.3	5.3	0.2	0.1	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.6	1.3	1.7	0.0	3.7	6.7	0.0	0.3	0.2	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	1.1	3.5	3.3	10.2	0.2	0.2	0.0
1978	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.5	0.0	5.1	6.5	0.0	0.2	0.2	0.0
1974-78	0.0	0.0	0.0	0.1	0.0	0.2	0.0	0.0	0.0	0.0	0.2	0.2	0.6	0.7	0.2	2.9	6.0	5.2	0.2	0.2	0.0
1979	0.0	0.0	0.0	0.2	0.3	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	1.1	1.6	0.0	24.4	0.2	0.2	0.0
1980	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	1.4	2.1	1.6	6.0	4.8	0.2	0.2	0.0
1981	0.3	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	1.4	4.0	0.0	0.0	0.0	0.2	0.2	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	0.6	0.0	3.8	3.0	2.8	9.3	0.2	0.2	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	2.1	0.0	0.9	1.5	0.0	4.5	0.2	0.2	0.0
1979-83	0.1	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.1	0.2	0.0	0.1	0.8	0.6	2.4	1.5	1.7	8.5	0.2	0.2	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MONOCYTIC LEUKAEMIA (ICD9 206)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	1	0	1	0	0	0	0	0	1	2	1	1	1	0	0	2	2	0	0	12
1965	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	1	0	1	0	4
1966	0	0	0	0	0	0	1	1	0	0	0	1	1	0	0	1	3	2	0	10
1967	0	0	0	0	0	0	0	0	1	1	1	0	2	0	3	1	2	0	0	11
1968	0	1	1	0	0	0	0	0	1	0	0	1	2	1	1	0	1	2	0	11
1964-68	1	1	2	0	0	0	1	1	3	4	2	4	6	1	4	5	8	5	0	48
1969	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	2	0	1	0	5
1970	0	0	0	0	0	0	0	0	0	1	0	0	1	0	2	0	0	0	0	4
1971	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1	1	1	1	0	6
1972	0	1	0	0	0	1	2	0	0	0	1	1	0	0	0	0	1	0	0	7
1973	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	3	0	0	5
1969-73	0	1	0	0	0	1	3	0	0	1	1	1	3	3	3	3	5	2	0	27
1974	0	0	0	0	0	0	1	0	2	0	0	1	0	0	0	0	0	3	0	7
1975	0	0	0	0	0	1	0	1	1	0	1	0	1	2	0	1	0	1	0	9
1976	0	0	0	0	0	0	1	0	1	0	0	1	0	2	0	0	3	1	0	9
1977	0	0	0	0	0	0	1	0	0	0	3	0	1	0	0	0	1	0	0	6
1978	0	0	0	1	0	0	1	0	1	0	1	0	0	1	1	0	0	1	0	7
1974-78	0	0	0	1	0	1	4	1	5	0	5	2	2	5	1	1	4	6	0	38
1979	0	0	0	1	0	0	0	2	0	0	1	0	0	0	1	1	2	0	0	8
1980	0	1	1	0	0	1	0	0	0	0	1	0	0	0	1	0	2	1	0	8
1981	0	0	0	0	0	0	0	0	0	0	2	0	1	0	0	0	0	0	0	3
1982	0	0	0	0	0	1	1	0	0	0	0	2	0	0	1	1	1	1	0	8
1983	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	1	0	1	0	5
1979-83	0	1	1	1	0	2	1	2	0	0	5	2	2	1	3	3	5	3	0	32

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MONOCYTIC LEUKAEMIA (ICD9 206)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.3	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.5	1.1	0.6	0.7	0.8	0.0	0.0	3.4	5.8	0.0	0.4	0.3	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.7	0.0	0.0	0.0	1.7	0.0	4.3	0.1	0.1	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.4	0.0	0.0	0.0	0.7	0.8	0.0	0.0	1.6	8.1	8.4	0.3	0.2	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.6	0.0	1.6	0.0	3.3	1.5	5.1	0.0	0.3	0.3	0.0
1968	0.0	0.3	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.6	1.5	0.9	1.1	0.0	2.5	7.5	0.3	0.3	0.0
1964-68	0.1	0.1	0.1	0.0	0.0	0.0	0.1	0.1	0.3	0.4	0.2	0.5	1.0	0.2	0.9	1.6	4.3	4.2	0.3	0.2	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.5	0.0	0.0	2.9	0.0	3.5	0.1	0.1	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.7	0.0	2.1	0.0	0.0	0.0	0.1	0.1	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	1.0	1.4	2.2	3.1	0.1	0.1	0.0
1972	0.0	0.3	0.0	0.0	0.0	0.3	0.8	0.0	0.0	0.0	0.5	0.6	0.0	0.0	0.0	0.0	2.1	0.0	0.2	0.2	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	6.1	0.0	0.1	0.1	0.0
1969-73	0.0	0.1	0.0	0.0	0.0	0.1	0.3	0.0	0.0	0.1	0.1	0.1	0.4	0.5	0.6	0.8	2.2	1.2	0.1	0.1	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	7.9	0.2	0.1	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.4	0.0	0.4	0.0	0.6	1.5	0.0	1.2	0.0	2.5	0.2	0.2	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.4	0.0	0.0	0.5	0.0	1.4	0.0	0.0	5.6	2.4	0.2	0.2	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	1.3	0.0	0.6	0.0	0.0	0.0	1.8	0.0	0.1	0.1	0.0
1978	0.0	0.0	0.0	0.2	0.0	0.0	0.3	0.0	0.4	0.0	0.4	0.0	0.0	0.7	0.8	0.0	0.0	2.2	0.1	0.1	0.0
1974-78	0.0	0.0	0.0	0.1	0.0	0.1	0.3	0.1	0.4	0.0	0.4	0.2	0.2	0.7	0.2	0.2	1.5	2.9	0.2	0.1	0.0
1979	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.7	0.0	0.0	0.4	0.0	0.0	0.0	0.8	1.1	3.4	0.0	0.2	0.1	0.0
1980	0.0	0.3	0.3	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.8	0.0	3.2	2.0	0.2	0.1	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.7	1.0	1.5	1.9	0.1	0.1	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	0.6	0.0	1.0	0.0	1.8	0.1	0.1	0.0
1979-83	0.0	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.4	0.2	0.2	0.1	0.5	0.6	1.6	1.2	0.1	0.1	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

OTHER SPECIFIED LEUKAEMIA (ICD9 207)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	2
1965	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1966	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	2
1967	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	3
1968	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1
1964-68	0	1	0	0	0	0	0	1	1	0	2	1	1	0	0	1	0	0	0	8
1969	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	2	1	0	0	6
1970	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	1	0	0	0	3
1971	1	0	0	0	1	0	1	0	1	0	0	0	0	1	0	1	1	0	0	7
1972	0	0	0	0	1	0	0	0	0	1	1	0	0	3	3	1	0	0	0	10
1973	0	0	0	0	0	0	0	0	1	0	0	0	0	4	1	1	2	1	0	10
1969-73	1	0	0	1	2	0	2	0	3	1	1	1	1	8	4	6	4	1	0	36
1974	0	0	0	0	0	0	0	0	1	1	0	3	0	1	0	1	0	0	0	7
1975	0	0	0	0	0	0	0	2	0	0	1	0	1	1	0	1	1	0	0	7
1976	0	0	0	0	0	0	0	1	0	0	0	3	2	1	1	0	0	1	0	9
1977	1	0	0	0	0	0	0	0	1	0	1	0	1	1	2	1	0	0	0	8
1978	0	0	0	0	0	0	0	0	0	0	0	1	2	2	0	0	2	1	0	8
1974-78	1	0	0	0	0	0	0	3	2	1	2	7	6	6	3	3	3	2	0	39
1979	0	0	0	0	0	0	0	0	0	0	0	0	3	0	3	1	1	1	0	9
1980	0	0	0	0	1	1	2	0	0	3	1	1	1	2	1	2	0	0	1	16
1981	1	0	0	0	0	0	0	0	0	1	1	0	0	2	1	1	0	1	0	8
1982	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	2	1	0	0	5
1983	0	0	0	0	0	0	0	0	1	0	0	1	1	0	2	1	0	0	0	6
1979-83	1	0	0	0	1	1	2	0	1	5	2	2	5	5	7	7	2	2	1	44

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

OTHER SPECIFIED LEUKAEMIA (ICD9 207)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.8	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1967	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	2.1	0.0	0.0	0.1	0.1	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1964-68	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.2	0.1	0.2	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0
1969	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.8	0.0	0.0	4.2	3.7	0.0	0.2	0.1	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	2.1	0.0	0.0	0.1	0.1	0.0
1971	0.3	0.0	0.0	0.0	0.3	0.0	0.4	0.0	0.4	0.0	0.0	0.0	0.0	0.9	0.0	2.0	3.5	0.0	0.2	0.2	0.0
1972	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.4	0.5	0.0	0.0	2.8	3.9	2.0	0.0	0.0	0.3	0.3	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	3.6	1.3	2.0	6.9	5.5	0.3	0.2	0.0
1969-73	0.1	0.0	0.0	0.1	0.1	0.0	0.2	0.0	0.2	0.1	0.1	0.1	0.1	1.5	1.1	2.5	2.8	1.2	0.2	0.2	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	1.8	0.0	0.9	0.0	2.0	0.0	0.0	0.2	0.2	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.4	0.0	0.6	0.9	0.0	1.9	3.4	0.0	0.2	0.2	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	1.7	1.3	0.8	1.2	0.0	0.0	5.2	0.2	0.2	0.0
1977	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.0	0.6	0.8	2.3	1.8	0.0	0.0	0.2	0.2	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	1.3	1.5	0.0	0.0	6.5	5.0	0.2	0.2	0.0
1974-78	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.1	0.2	0.8	0.8	1.0	0.7	1.1	2.0	2.1	0.2	0.2	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.9	0.0	3.2	1.6	3.1	4.9	0.2	0.2	0.0
1980	0.0	0.0	0.0	0.0	0.3	0.3	0.6	0.0	0.0	1.3	0.4	0.5	0.6	1.4	1.0	3.2	0.0	0.0	0.3	0.3	0.0
1981	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	0.0	1.4	1.0	1.5	0.0	4.7	0.2	0.2	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.7	0.0	3.0	2.8	0.0	0.1	0.1	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	0.5	0.0	1.8	1.5	0.0	0.0	0.1	0.1	0.0
1979-83	0.1	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.1	0.4	0.2	0.2	0.6	0.7	1.4	2.2	1.2	1.9	0.2	0.2	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

OTHER SPECIFIED LEUKAEMIA (ICD9 207)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1965	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
1966	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1
1967	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
1968	0	0	0	0	0	1	0	0	0	0	1	0	1	0	0	0	0	0	0	3
1964-68	1	1	0	0	0	1	0	0	0	0	1	0	2	0	0	0	0	0	0	6
1969	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1
1970	0	0	1	0	0	0	0	0	0	0	0	0	1	0	2	0	1	0	0	5
1971	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	3
1972	1	0	0	0	1	0	0	0	0	0	0	1	1	1	1	1	0	0	0	7
1973	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	2	0	0	0	3
1969-73	1	0	2	0	1	0	0	0	0	1	0	2	3	1	3	4	1	0	0	19
1974	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	2
1975	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	1	0	4
1976	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	2
1977	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	2
1978	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1
1974-78	0	0	0	0	0	0	0	0	1	0	0	1	2	3	0	1	1	2	0	11
1979	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	2	1	0	0	5
1980	0	0	0	0	0	0	0	0	0	1	0	1	1	0	1	1	0	1	0	6
1981	0	0	0	0	0	0	0	0	0	0	0	0	1	1	2	1	0	0	0	5
1982	0	0	0	0	0	0	0	1	0	0	0	2	0	0	1	0	0	0	0	4
1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	2
1979-83	0	0	0	0	0	0	0	1	0	1	0	5	2	1	5	5	1	1	0	22

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

OTHER SPECIFIED LEUKAEMIA (ICD9 207)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1965	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1967	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.6	0.0	0.8	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0
1964-68	0.1	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1970	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.0	2.1	0.0	2.3	0.0	0.1	0.1	0.0
1971	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0	1.4	0.0	0.0	0.1	0.1	0.0
1972	0.3	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	0.8	1.0	1.3	0.0	0.0	0.2	0.2	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	2.6	0.0	0.0	0.1	0.1	0.0
1969-73	0.1	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.4	0.2	0.6	1.1	0.4	0.0	0.1	0.1	0.0
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	1.3	0.0	0.0	0.0	0.0	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.6	0.0	0.0	0.0	0.0	1.9	2.5	0.1	0.1	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0	0.0	2.3	0.0	0.0	0.0
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1974-78	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.2	0.4	0.0	0.2	0.4	1.0	0.0	0.0	0.0
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.0	2.2	1.7	0.0	0.1	0.1	0.0
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	0.6	0.0	0.8	1.1	0.0	2.0	0.1	0.1	0.0
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.6	1.5	1.0	0.0	0.0	0.1	0.1	0.0
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.9	0.0	0.0	0.7	0.0	0.0	0.0	0.1	0.1	0.0
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	1.0	0.0	0.0	0.0	0.0	0.0
1979-83	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.4	0.2	0.1	0.8	1.0	0.3	0.4	0.1	0.1	0.0

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

LEUKAEMIA OF UNSPECIFIED CELL TYPE (ICD9 208)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	20	27	9	4	1	3	3	7	1	4	4	11	15	11	16	11	10	9	0	166
1965	27	8	6	4	2	4	3	3	7	5	8	13	9	10	13	12	6	6	0	146
1966	36	24	5	3	2	4	1	4	1	8	4	8	5	20	13	13	7	9	0	167
1967	13	6	3	5	2	3	4	3	2	5	7	11	9	18	9	17	8	9	0	134
1968	18	8	12	2	2	5	4	2	7	7	6	9	14	12	21	18	13	5	0	165
1964-68	114	73	35	18	9	19	15	19	18	29	29	52	52	71	72	71	44	38	0	778
1969	11	3	5	6	3	2	4	2	5	5	2	5	5	13	13	18	8	11	0	121
1970	15	5	3	3	1	2	0	2	3	3	4	7	13	11	18	6	7	6	0	109
1971	8	4	2	3	1	0	2	1	3	3	4	9	7	6	6	16	11	3	0	89
1972	5	4	1	1	3	1	0	2	6	2	5	5	10	9	14	10	6	6	0	90
1973	6	1	5	3	1	3	3	1	4	1	5	8	13	14	12	10	4	9	1	104
1969-73	45	17	16	16	9	8	9	8	21	14	20	34	48	53	63	60	36	35	1	513
1974	9	4	4	2	3	1	4	0	2	1	9	9	10	15	12	7	7	3	1	103
1975	4	0	3	4	3	1	0	2	1	1	7	4	11	12	12	10	9	7	0	91
1976	2	1	1	2	2	1	2	2	1	3	5	6	8	17	12	10	9	7	0	91
1977	2	1	1	0	3	0	2	1	2	5	3	6	7	8	7	11	10	13	0	82
1978	0	2	0	1	1	0	2	1	1	1	8	3	3	5	11	10	5	8	0	62
1974-78	17	8	9	9	12	3	10	6	7	11	32	28	39	57	54	48	40	38	1	429
1979	0	0	2	1	1	0	1	1	0	1	2	1	7	5	6	8	11	4	0	51
1980	0	0	0	0	1	0	0	1	1	4	0	2	5	6	13	7	7	12	0	59
1981	1	0	1	1	0	0	0	5	1	2	3	5	7	15	9	17	6	6	0	79
1982	0	0	1	0	0	1	3	1	0	1	5	3	1	5	13	14	7	3	0	58
1983	1	0	1	0	1	0	1	0	1	1	3	3	7	6	8	8	6	7	2	56
1979-83	2	0	5	2	3	1	5	8	3	9	13	14	27	37	49	54	37	32	2	303

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

LEUKAEMIA OF UNSPECIFIED CELL TYPE (ICD9 208)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	5.1	7.3	2.7	1.5	0.5	1.5	1.3	3.0	0.5	2.1	2.4	7.9	13.2	12.2	23.4	24.2	39.4	66.2	5.1	4.8	0.4
1965	6.9	2.1	1.8	1.4	0.9	1.9	1.3	1.3	3.1	2.6	4.6	9.1	7.7	10.9	18.7	26.1	23.2	43.2	4.3	4.2	0.4
1966	9.4	6.1	1.4	1.0	0.8	1.9	0.4	1.7	0.4	4.1	2.2	5.4	4.2	21.5	18.5	27.9	26.9	64.3	4.9	4.8	0.4
1967	3.5	1.5	0.8	1.6	0.8	1.3	1.8	1.2	0.8	2.5	3.8	7.2	7.3	18.9	12.7	35.8	30.3	62.1	3.8	3.7	0.3
1968	5.0	2.0	3.2	0.6	0.7	2.1	1.7	0.8	2.9	3.4	3.3	5.7	11.1	12.3	29.5	37.7	48.5	33.3	4.6	4.5	0.4
1964-68	6.0	3.7	2.0	1.2	0.7	1.7	1.3	1.6	1.6	2.9	3.3	7.0	8.7	15.2	20.6	30.4	33.7	53.5	4.5	4.4	0.4
1969	3.2	0.7	1.3	1.8	1.0	0.8	1.7	0.8	2.1	2.3	1.1	3.1	3.9	13.0	18.1	37.4	29.4	70.1	3.3	3.3	0.3
1970	4.5	1.2	0.8	0.9	0.3	0.7	0.0	0.8	1.2	1.3	2.2	4.2	9.8	10.7	24.7	12.4	25.2	36.6	2.9	3.0	0.3
1971	2.5	1.0	0.5	0.8	0.3	0.0	0.8	0.4	1.2	1.3	2.1	5.3	5.1	5.7	8.1	32.7	39.0	17.3	2.4	2.2	0.2
1972	1.5	1.0	0.2	0.3	0.9	0.3	0.0	0.8	2.5	0.9	2.5	2.9	7.1	8.3	18.3	20.2	20.9	33.5	2.3	2.3	0.2
1973	1.9	0.3	1.2	0.8	0.3	0.9	1.2	0.4	1.6	0.4	2.4	4.7	8.9	12.7	15.1	20.1	13.7	49.5	2.6	2.6	0.3
1969-73	2.7	0.9	0.8	0.9	0.6	0.6	0.7	0.7	1.7	1.2	2.1	4.0	7.0	10.1	16.8	24.5	25.5	40.9	2.7	2.7	0.3
1974	2.8	1.1	1.0	0.5	0.9	0.3	1.5	0.0	0.8	0.4	4.1	5.3	6.7	13.3	14.6	13.8	24.0	16.2	2.5	2.5	0.3
1975	1.2	0.0	0.7	1.0	0.8	0.3	0.0	0.8	0.4	0.4	3.1	2.3	7.1	10.3	14.2	19.1	30.8	36.8	2.2	2.1	0.2
1976	0.6	0.3	0.2	0.5	0.5	0.3	0.7	0.8	0.4	1.3	2.2	3.4	5.1	14.1	13.9	18.4	30.3	36.1	2.2	2.1	0.2
1977	0.6	0.3	0.3	0.0	0.8	0.0	0.6	0.4	0.8	2.1	1.3	3.2	4.4	6.4	7.9	19.5	33.1	66.0	1.9	1.8	0.1
1978	0.0	0.6	0.0	0.2	0.3	0.0	0.6	0.4	0.4	0.4	3.5	1.5	1.9	3.8	12.1	17.0	16.2	40.0	1.4	1.3	0.1
1974-78	1.1	0.5	0.4	0.4	0.7	0.2	0.7	0.5	0.6	0.9	2.8	3.1	5.0	9.4	12.5	17.6	26.8	39.3	2.0	2.0	0.2
1979	0.0	0.0	0.5	0.2	0.3	0.0	0.3	0.4	0.0	0.4	0.9	0.5	4.5	3.7	6.4	13.1	34.6	19.5	1.1	1.0	0.1
1980	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	1.7	0.0	0.9	3.1	4.3	13.4	11.1	21.1	58.0	1.3	1.2	0.1
1981	0.3	0.0	0.3	0.2	0.0	0.0	0.0	1.7	0.4	0.8	1.3	2.3	4.1	10.6	8.9	26.3	17.6	28.3	1.7	1.6	0.2
1982	0.0	0.0	0.3	0.0	0.0	0.3	0.9	0.3	0.0	0.4	2.1	1.4	0.6	3.5	12.4	21.1	19.6	13.9	1.2	1.1	0.1
1983	0.3	0.0	0.3	0.0	0.2	0.0	0.3	0.0	0.4	0.4	1.3	1.4	3.7	4.2	7.3	11.7	15.9	31.4	1.1	1.0	0.1
1979-83	0.1	0.0	0.3	0.1	0.2	0.1	0.3	0.5	0.2	0.8	1.1	1.3	3.2	5.3	9.7	16.7	21.4	30.1	1.3	1.2	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

LEUKAEMIA OF UNSPECIFIED CELL TYPE (ICD9 208)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	29	13	6	5	2	4	10	4	3	6	4	9	4	12	14	7	2	5	0	139
1965	20	14	2	1	2	5	2	1	2	4	4	3	7	4	9	9	2	1	0	92
1966	16	15	3	2	1	4	5	1	2	2	7	5	6	8	11	9	5	3	0	105
1967	17	8	1	1	4	2	1	4	5	8	9	4	9	8	23	16	10	6	0	136
1968	8	10	4	2	1	2	2	2	3	3	2	8	8	9	16	17	10	9	0	116
1964-68	90	60	16	11	10	17	20	12	15	23	26	29	34	41	73	58	29	24	0	588
1969	5	7	1	1	2	2	2	2	2	7	9	6	5	9	9	10	7	7	0	93
1970	9	1	3	2	0	0	0	0	3	5	1	9	3	5	14	9	6	2	0	72
1971	6	4	3	4	1	4	1	3	0	1	1	3	1	4	5	10	10	6	0	67
1972	4	3	4	2	2	0	0	2	2	1	6	5	6	9	12	7	12	13	1	91
1973	4	1	1	0	6	0	2	1	4	1	2	4	6	7	7	11	10	8	0	75
1969-73	28	16	12	9	11	6	5	8	11	15	19	27	21	34	47	47	45	36	1	398
1974	3	1	0	4	5	2	1	1	2	2	5	6	3	9	6	13	6	12	1	82
1975	0	0	0	0	1	1	2	2	3	1	3	1	3	3	8	10	11	3	0	52
1976	0	0	4	4	5	1	1	4	5	1	6	6	4	8	7	11	11	12	0	90
1977	2	0	3	1	5	2	1	1	2	3	3	4	2	8	5	8	0	9	0	59
1978	1	1	0	1	1	0	4	0	2	1	2	2	4	6	1	8	8	11	0	53
1974-78	6	2	7	10	17	6	9	8	14	8	19	19	16	34	27	50	36	47	1	336
1979	0	0	1	1	1	0	1	0	1	1	2	3	4	1	12	9	14	9	0	60
1980	1	0	0	0	1	0	0	2	1	2	2	1	3	4	6	8	5	5	2	43
1981	0	0	0	0	1	0	0	0	1	0	1	4	0	6	3	7	9	4	0	36
1982	3	0	0	0	2	1	0	1	0	2	1	1	5	4	6	1	6	12	0	45
1983	0	1	0	0	0	4	3	2	1	0	1	2	7	12	10	5	9	9	2	68
1979-83	4	1	1	1	5	5	4	5	4	5	7	11	19	27	37	30	43	39	4	252

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

LEUKAEMIA OF UNSPECIFIED CELL TYPE (ICD9 208)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	7.8	3.7	1.9	1.9	0.9	1.9	4.6	1.7	1.4	3.2	2.4	6.5	3.4	11.8	16.9	11.9	5.8	22.7	4.2	4.0	0.3
1965	5.4	3.8	0.6	0.4	0.9	2.4	0.9	0.4	0.9	2.1	2.4	2.1	5.8	3.9	10.5	14.9	5.6	4.3	2.7	2.6	0.2
1966	4.4	4.0	0.9	0.7	0.4	1.8	2.3	0.4	0.8	1.0	4.0	3.4	4.8	7.5	12.4	14.5	13.4	12.7	3.0	2.8	0.2
1967	4.8	2.1	0.3	0.3	1.5	0.9	0.4	1.7	2.1	3.9	5.0	2.6	7.0	7.4	25.5	24.7	25.6	23.9	3.8	3.4	0.3
1968	2.3	2.6	1.1	0.6	0.4	0.8	0.9	0.9	1.2	1.4	1.1	5.1	6.1	8.1	17.5	25.4	24.6	34.0	3.2	2.7	0.3
1964-68	5.0	3.2	0.9	0.7	0.8	1.5	1.8	1.0	1.3	2.3	3.0	3.9	5.4	7.7	16.6	18.6	15.5	20.0	3.4	3.1	.0.3
1969	1.5	1.8	0.3	0.3	0.7	0.8	0.9	0.9	0.8	3.2	4.9	3.7	3.7	7.9	9.7	14.5	16.5	24.7	2.5	2.2	0.2
1970	2.8	0.3	0.8	0.6	0.0	0.0	0.0	0.0	1.3	2.2	0.5	5.3	2.1	4.2	14.8	12.8	13.6	6.6	1.9	1.7	0.2
1971	1.9	1.0	0.8	1.1	0.3	1.4	0.4	1.3	0.0	0.4	0.5	1.7	0.7	3.3	5.1	13.9	21.6	18.5	1.7	1.4	0.1
1972	1.3	0.8	1.0	0.6	0.6	0.0	0.0	0.9	0.9	0.4	2.9	2.9	4.0	7.2	12.1	9.4	25.3	37.9	2.2	1.8	0.2
1973	1.3	0.3	0.3	0.0	1.8	0.0	0.8	0.4	1.7	0.4	0.9	2.3	3.9	5.4	6.9	14.5	20.4	22.3	1.8	1.5	0.1
1969-73	1.8	0.9	0.6	0.5	0.7	0.4	0.4	0.7	0.9	1.3	1.9	3.2	2.9	5.6	9.7	13.0	19.6	22.3	2.0	1.7	0.2
1974	1.0	0.3	0.0	1.1	1.4	0.6	0.4	0.4	0.9	0.8	2.2	3.4	1.9	6.8	5.7	16.7	12.0	31.7	1.9	1.6	0.1
1975	0.0	0.0	0.0	0.0	0.3	0.3	0.7	0.8	1.3	0.4	1.3	0.6	1.8	2.2	7.4	12.5	21.3	7.6	1.2	0.9	0.1
1976	0.0	0.0	1.0	1.0	1.3	0.3	0.3	1.6	2.1	0.4	2.5	3.2	2.4	5.7	6.3	13.3	20.6	29.0	2.0	1.6	0.1
1977	0.7	0.0	0.8	0.3	1.3	0.6	0.3	0.4	0.9	1.3	1.3	2.0	1.2	5.5	4.4	9.4	0.0	20.8	1.3	1.1	0.1
1978	0.3	0.3	0.0	0.2	0.3	0.0	1.2	0.0	0.9	0.4	0.8	0.9	2.3	4.0	0.8	9.2	14.0	24.5	1.1	0.9	0.1
1974-78	0.4	0.1	0.4	0.5	0.9	0.3	0.6	0.7	1.2	0.7	1.6	2.0	1.9	4.8	4.8	12.1	13.5	22.7	1.5	1.2	0.1
1979	0.0	0.0	0.3	0.2	0.3	0.0	0.3	0.0	0.4	0.4	0.9	1.3	2.3	0.6	9.8	10.0	23.5	19.1	1.2	0.9	0.1
1980	0.3	0.0	0.0	0.0	0.3	0.0	0.0	0.7	0.4	0.9	0.9	0.4	1.7	2.5	4.8	8.6	8.1	10.2	0.8	0.7	0.1
1981	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	0.0	0.4	1.7	0.0	3.6	2.3	7.3	14.1	7.8	0.7	0.5	0.0
1982	1.0	0.0	0.0	0.0	0.5	0.3	0.0	0.3	0.0	0.9	0.4	0.4	2.5	2.4	4.5	1.0	9.2	22.2	0.8	0.7	0.1
1983	0.0	0.3	0.0	0.0	0.0	1.0	0.8	0.6	0.4	0.0	0.4	0.9	3.3	7.2	7.2	4.8	13.2	15.8	1.1	1.0	0.1
1979-83	0.3	0.1	0.1	0.1	0.3	0.3	0.2	0.3	0.3	0.4	0.6	1.0	2.0	3.3	5.7	6.2	13.5	15.1	0.9	0.7	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES (ICD9 195)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	0	2	1	6	4	5	2	10	3	2	0	35
1965	0	0	0	0	0	0	0	0	1	0	3	3	6	4	2	5	5	0	0	29
1966	0	0	0	0	0	1	1	0	0	0	2	2	2	7	7	2	7	1	0	32
1967	0	1	1	0	0	0	0	1	0	1	1	3	8	4	4	10	6	4	0	44
1968	0	0	1	0	0	0	1	1	1	3	3	3	5	8	13	10	6	5	0	60
1964-68	0	1	2	0	0	1	2	2	2	6	10	17	25	28	28	37	27	12	0	200
1969	0	0	0	0	0	0	0	0	2	1	0	3	7	9	9	12	6	9	0	58
1970	0	0	0	0	0	0	1	0	2	3	1	4	3	5	9	16	11	11	0	66
1971	0	0	1	0	0	0	0	0	0	3	8	9	12	8	13	14	6	7	0	81
1972	0	0	1	1	0	0	1	1	4	3	4	5	6	8	8	13	9	7	0	71
1973	2	0	1	0	1	0	1	3	3	13	5	7	3	9	8	19	11	8	2	96
1969-73	2	0	3	1	1	0	3	4	11	23	18	28	31	39	47	74	43	42	2	372
1974	0	0	0	0	1	0	0	0	1	6	11	7	14	21	14	10	8	19	1	113
1975	0	0	0	1	2	0	2	0	5	3	6	11	13	9	11	18	12	4	3	100
1976	0	0	0	1	0	0	0	2	4	9	14	10	23	16	11	17	18	14	0	139
1977	0	0	0	0	1	0	2	2	1	5	9	11	10	20	10	13	10	10	0	104
1978	0	0	0	0	0	0	0	0	1	2	6	11	11	18	20	15	17	12	0	113
1974-78	0	0	0	2	4	0	4	4	12	25	46	50	71	84	66	73	65	59	4	569
1979	0	0	0	1	1	0	0	2	2	4	7	7	11	18	12	14	15	11	0	105
1980	0	0	1	1	0	0	1	1	3	2	7	11	11	18	13	19	9	6	0	103
1981	0	0	0	0	1	0	1	1	3	2	7	10	12	16	22	15	13	10	1	114
1982	0	0	1	3	0	0	0	2	0	4	3	6	13	17	21	10	7	14	1	102
1983	0	2	0	0	0	1	0	0	3	3	4	8	19	10	6	17	12	7	0	92
1979-83	0	2	2	5	2	1	2	6	11	15	28	42	66	79	74	75	56	48	2	516

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES (ICD9 195)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.6	4.3	3.5	5.6	2.9	22.0	11.8	14.7	1.1	1.0	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.7	2.1	5.1	4.4	2.9	10.9	19.3	0.0	0.9	0.8	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.5	0.4	0.0	0.0	0.0	1.1	1.4	1.7	7.5	9.9	4.3	26.9	7.1	0.9	0.9	0.1
1967	0.0	0.3	0.3	0.0	0.0	0.0	0.0	0.4	0.0	0.5	0.5	2.0	6.5	4.2	5.6	21.1	22.7	27.6	1.3	1.2	0.1
1968	0.0	0.0	0.3	0.0	0.0	0.0	0.4	0.4	0.4	1.4	1.6	1.9	4.0	8.2	18.3	20.9	22.4	33.3	1.7	1.6	0.2
1964-68	0.0	0.1	0.1	0.0	0.0	0.1	0.2	0.2	0.2	0.6	1.1	2.3	4.2	6.0	8.0	15.9	20.7	16.9	1.2	1.1	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.5	0.0	1.8	5.4	9.0	12.5	24.9	22.1	57.3	1.6	1.5	0.2
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.8	1.3	0.5	2.4	2.3	4.9	12.3	33.0	39.6	67.1	1.8	1.6	0.1
1971	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	1.3	4.2	5.3	8.8	7.5	17.5	28.6	21.3	40.5	2.1	2.0	0.2
1972	0.0	0.0	0.2	0.3	0.0	0.0	0.4	0.4	1.6	1.3	2.0	2.9	4.3	7.4	10.4	26.3	31.4	39.1	1.8	1.7	0.2
1973	0.6	0.0	0.2	0.0	0.3	0.0	0.4	1.3	1.2	5.5	2.4	4.1	2.1	8.2	10.1	38.2	37.8	44.0	2.4	2.2	0.2
1969-73	0.1	0.0	0.1	0.1	0.1	0.0	0.2	0.3	0.9	2.0	1.9	3.3	4.5	7.4	12.5	30.2	30.5	49.1	2.0	1.8	0.2
1974	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	2.5	5.0	4.2	9.3	16.6	17.0	19.8	27.4	102.7	2.8	2.7	0.3
1975	0.0	0.0	0.0	0.2	0.6	0.0	0.7	0.0	2.1	1.3	2.7	6.4	8.4	7.8	13.0	34.4	41.1	21.1	2.4	2.2	0.2
1976	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.8	1.7	3.8	6.2	5.6	14.6	13.3	12.7	31.3	60.6	72.2	3.3	3.1	0.3
1977	0.0	0.0	0.0	0.0	0.3	0.0	0.6	0.8	0.4	2.1	4.0	5.8	6.3	15.9	11.3	23.0	33.1	50.8	2.4	2.3	0.2
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	2.6	5.5	7.0	13.8	22.1	25.5	55.2	60.0	2.5	2.4	0.3
1974-78	0.0	0.0	0.0	0.1	0.2	0.0	0.3	0.3	1.0	2.1	4.1	5.5	9.1	13.9	15.3	26.8	43.6	61.1	2.7	2.5	0.3
1979	0.0	0.0	0.0	0.2	0.3	0.0	0.0	0.7	0.8	1.7	3.0	3.4	7.0	13.4	12.8	22.9	47.2	53.7	2.3	2.2	0.2
1980	0.0	0.0	0.3	0.2	0.0	0.0	0.3	0.4	1.2	0.8	3.0	5.2	6.8	13.0	13.4	30.2	27.2	29.0	2.2	2.1	0.2
1981	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.3	1.2	0.8	3.0	4.6	7.1	11.3	21.8	23.2	38.1	47.2	2.4	2.2	0.3
1982	0.0	0.0	0.3	0.7	0.0	0.0	0.0	0.6	0.0	1.7	1.3	2.7	7.3	11.9	20.1	15.0	19.6	64.8	2.1	2.0	0.2
1983	0.0	0.6	0.0	0.0	0.0	0.3	0.0	0.0	1.1	1.3	1.7	3.6	10.1	7.0	5.5	24.8	31.7	31.4	1.8	1.7	0.2
1979-83	0.0	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.9	1.3	2.4	3.9	7.7	11.3	14.7	23.2	32.5	45.2	2.2	2.1	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES (ICD9 195)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	1	1	0	3	3	3	5	3	5	1	3	0	28
1965	0	0	0	0	0	0	1	0	2	0	2	3	3	2	2	3	1	1	0	20
1966	0	0	0	0	1	0	0	3	1	1	2	1	1	1	1	3	2	0	0	18
1967	1	0	0	0	0	1	0	2	3	5	6	4	0	3	3	6	6	8	0	47
1968	0	0	0	0	1	2	0	1	3	5	3	3	5	5	3	7	3	2	0	43
1964-68	1	0	0	0	2	3	1	7	10	11	16	14	12	16	12	24	13	14	0	156
1969	0	0	0	0	1	0	0	0	1	5	2	4	4	1	7	13	8	10	0	56
1970	1	0	1	0	1	1	1	1	3	2	1	3	10	5	4	5	8	9	0	56
1971	0	0	0	0	1	0	2	2	1	1	7	1	3	4	6	6	8	6	0	48
1972	0	0	0	1	0	2	0	0	5	0	3	10	6	6	12	13	10	7	0	75
1973	4	0	0	0	2	2	3	0	3	2	6	5	4	14	10	9	7	15	0	86
1969-73	5	0	1	1	5	5	6	3	13	10	19	23	27	30	39	46	41	47	0	321
1974	2	0	0	1	1	1	1	2	4	0	5	5	9	19	7	8	8	15	3	91
1975	0	0	0	1	0	1	1	2	6	7	9	6	7	5	5	10	11	19	0	90
1976	1	0	0	2	2	1	1	2	5	7	5	5	8	9	12	19	11	9	0	99
1977	0	0	0	0	2	4	1	0	3	3	8	10	4	12	13	7	16	11	0	94
1978	0	0	0	1	1	1	1	1	2	3	5	6	12	6	9	2	5	12	1	68
1974-78	3	0	0	5	6	8	5	7	20	20	32	32	40	51	46	46	51	66	4	442
1979	1	0	0	0	1	1	2	3	1	3	2	5	8	9	5	14	17	22	0	94
1980	1	0	0	0	0	2	0	2	2	8	13	6	5	12	17	12	19	18	0	117
1981	1	0	0	2	0	2	3	2	5	2	4	4	12	8	15	12	9	22	0	103
1982	0	0	0	0	0	0	1	2	4	7	8	10	7	9	8	10	12	17	1	96
1983	0	0	0	0	2	0	3	2	0	2	2	5	7	9	14	8	10	21	1	86
1979-83	3	0	0	2	3	5	9	11	12	22	29	30	39	47	59	56	67	100	2	496

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM OF OTHER AND ILL-DEFINED SITES (ICD9 195)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.0	1.8	2.2	2.5	4.9	3.6	8.5	2.9	13.6	0.9	0.7	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.9	0.0	1.2	2.1	2.5	1.9	2.3	5.0	2.8	4.3	0.6	0.5	0.1
1966	0.3	0.0	0.0	0.0	0.4	0.0	0.0	1.3	0.4	0.5	1.1	0.7	0.8	0.9	1.1	4.8	5.4	0.0	0.5	0.4	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.9	1.2	2.5	3.4	2.6	0.0	2.8	3.3	9.3	15.4	31.9	1.3	1.1	0.1
1968	0.0	0.0	0.0	0.0	0.4	0.8	0.0	0.4	1.2	2.4	1.7	1.9	3.8	4.5	3.3	10.4	7.4	7.5	1.2	1.0	0.1
1964-68	0.1	0.0	0.0	0.0	0.2	0.3	0.1	0.6	0.9	1.1	1.8	1.9	1.9	3.0	2.7	7.7	6.9	11.6	0.9	0.8	0.1
1969	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	2.3	1.1	2.5	2.9	0.9	7.6	18.9	18.9	35.3	1.5	1.1	0.1
1970	0.3	0.0	0.3	0.0	0.3	0.4	0.4	0.4	1.3	0.9	0.5	1.8	7.2	4.2	4.2	7.1	18.1	29.7	1.4	1.2	0.1
1971	0.0	0.0	0.0	0.0	0.3	0.0	0.8	0.9	0.4	0.4	3.6	0.6	2.1	3.3	6.2	8.3	17.3	18.5	1.2	1.0	0.1
1972	0.0	0.0	0.0	0.3	0.0	0.7	0.0	0.0	2.2	0.0	1.5	5.7	4.0	4.8	12.1	17.5	21.1	20.4	1.8	1.4	0.2
1973	1.3	0.0	0.0	0.0	0.6	0.6	1.2	0.0	1.3	0.8	2.8	2.9	2.6	10.9	9.8	11.8	14.3	41.8	2.0	1.7	0.2
1969-73	0.3	0.0	0.1	0.1	0.3	0.4	0.5	0.3	1.1	0.9	1.9	2.7	3.7	4.9	8.0	12.7	17.9	29.1	1.6	1.3	0.1
1974	0.7	0.0	0.0	0.3	0.3	0.3	0.4	0.9	1.7	0.0	2.2	2.8	5.6	14.4	6.7	10.3	16.0	39.6	2.0	1.7	0.2
1975	0.0	0.0	0.0	0.3	0.0	0.3	0.4	0.3	2.6	3.0	3.8	3.3	4.2	3.7	4.6	12.5	21.3	48.2	2.0	1.6	0.1
1976	0.3	0.0	0.0	0.5	0.5	0.3	0.3	0.8	2.1	3.0	2.1	2.6	4.7	6.4	10.8	23.0	20.6	21.7	2.2	1.8	0.2
1977	0.0	0.0	0.0	0.0	0.5	1.1	0.3	0.0	1.3	1.3	3.4	4.9	2.3	8.2	11.3	8.3	28.9	25.4	2.0	1.6	0.2
1978	0.0	0.0	0.0	0.2	0.3	0.3	0.3	0.4	0.9	1.3	2.1	2.8	7.0	4.0	7.6	2.3	8.7	26.7	1.4	1.2	0.1
1974-78	0.2	0.0	0.0	0.3	0.3	0.5	0.3	0.6	1.7	1.7	2.7	3.3	4.8	7.2	8.3	11.1	19.1	31.9	1.9	1.6	0.2
1979	0.3	0.0	0.0	0.0	0.3	0.3	0.6	1.1	0.4	1.3	0.9	2.2	4.6	5.8	4.1	15.5	28.5	46.7	1.8	1.4	0.1
1980	0.3	0.0	0.0	0.0	0.0	0.5	0.0	0.7	0.8	3.4	5.6	2.6	2.3	7.4	13.5	12.9	30.7	36.7	2.2	1.8	0.2
1981	0.3	0.0	0.0	0.5	0.0	0.5	0.8	0.7	2.0	0.9	1.7	1.7	6.4	4.9	11.5	12.5	14.1	42.7	1.9	1.6	0.2
1982	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.6	1.6	3.0	3.4	4.3	3.5	5.4	6.0	10.0	18.3	31.5	1.7	1.4	0.1
1983	0.0	0.0	0.0	0.0	0.5	0.0	0.8	0.6	0.0	0.9	0.9	2.1	3.3	5.4	10.0	7.7	14.7	36.9	1.4	1.1	0.1
1979-83	0.2	0.0	0.0	0.1	0.2	0.3	0.5	0.7	1.0	1.9	2.5	2.6	4.1	5.8	9.0	11.6	21.0	38.7	1.8	1.5	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

SECONDARY AND UNSPECIFIED NEOPLASM OF LYMPH NODES (ICD9 196)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	1	0	0	1	1	0	4	4	0	3	0	2	2	0	18
1965	0	0	1	0	0	0	0	0	0	0	0	1	4	4	1	0	0	0	0	11
1966	0	0	0	0	1	0	0	0	0	1	2	0	3	1	1	0	1	0	0	10
1967	0	0	0	0	0	0	0	1	2	3	1	2	4	3	2	2	0	1	0	21
1968	0	0	0	0	0	0	0	0	0	0	1	1	1	1	0	2	1	1	0	8
1964-68	0	0	1	0	1	1	0	1	3	5	4	8	16	9	7	4	4	4	0	68
1969	0	0	0	0	0	0	0	2	1	0	2	2	3	2	3	3	2	1	0	21
1970	0	0	0	0	0	0	0	0	1	0	2	0	1	2	5	1	1	1	0	14
1971	0	0	0	0	1	0	0	0	1	2	2	0	6	0	0	2	2	1	0	17
1972	0	1	0	0	0	2	0	0	1	3	1	4	3	1	1	2	1	2	0	22
1973	0	0	0	0	0	0	1	0	0	3	5	1	4	4	3	0	2	1	1	25
1969-73	0	1	0	0	1	2	1	2	4	8	12	7	17	9	12	8	8	6	1	99
1974	0	0	1	0	1	1	0	1	1	1	7	2	0	5	3	3	0	1	1	26
1975	0	0	0	0	0	0	0	0	0	0	4	2	7	4	3	0	0	1	0	21
1976	0	0	0	0	0	1	2	0	1	1	2	2	3	5	1	1	2	1	1	23
1977	0	0	0	0	0	0	0	0	1	2	3	3	2	6	4	5	1	0	1	28
1978	0	0	0	0	0	0	1	0	0	0	5	1	4	6	3	1	2	1	1	25
1974-78	0	0	1	0	1	2	3	1	3	4	21	10	16	26	14	10	5	4	4	125
1979	0	0	0	0	0	0	1	1	0	3	3	3	7	5	4	0	1	0	0	28
1980	0	0	0	0	0	1	1	3	0	4	3	5	3	4	3	2	4	2	0	35
1981	0	0	0	0	0	0	1	1	1	3	4	9	5	4	9	2	0	0	1	40
1982	0	0	0	0	1	2	0	1	0	2	2	3	2	6	4	4	1	1	1	30
1983	0	0	0	0	0	0	1	1	2	4	8	5	5	7	3	2	1	2	1	42
1979-83	0	0	0	0	1	3	4	7	3	16	20	25	22	26	23	10	7	5	3	175

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

SECONDARY AND UNSPECIFIED NEOPLASM OF LYMPH NODES (ICD9 196)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.5	0.5	0.0	2.9	3.5	0.0	4.4	0.0	7.9	14.7	0.6	0.6	0.1
1965	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.7	3.4	4.4	1.4	0.0	0.0	0.0	0.3	0.4	0.1
1966	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.5	1.1	0.0	2.5	1.1	1.4	0.0	3.8	0.0	0.3	0.3	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.5	0.5	1.3	3.3	3.2	2.8	4.2	0.0	6.9	0.6	0.6	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.6	0.8	1.0	0.0	4.2	3.7	6.7	0.2	0.2	0.0
1964-68	0.0	0.0	0.1	0.0	0.1	0.1	0.0	0.1	0.3	0.5	0.5	1.1	2.7	1.9	2.0	1.7	3.1	5.6	0.4	0.4	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.4	0.0	1.1	1.2	2.3	2.0	4.2	6.2	7.4	6.4	0.6	0.5	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.1	0.0	0.8	1.9	6.9	2.1	3.6	6.1	0.4	0.4	0.1
1971	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	0.9	1.1	0.0	4.4	0.0	0.0	4.1	7.1	5.8	0.4	0.4	0.0
1972	0.0	0.3	0.0	0.0	0.0	0.7	0.0	0.0	0.4	1.3	0.5	2.3	2.1	0.9	1.3	4.0	3.5	11.2	0.6	0.6	0.0
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.3	2.4	0.6	2.7	3.6	3.8	0.0	6.9	5.5	0.6	0.6	0.1
1969-73	0.0	0.1	0.0	0.0	0.1	0.1	0.1	0.2	0.3	0.7	1.2	0.8	2.5	1.7	3.2	3.3	5.7	7.0	0.5	0.5	0.1
1974	0.0	0.0	0.2	0.0	0.3	0.3	0.0	0.4	0.4	0.4	3.2	1.2	0.0	4.4	3.6	5.9	0.0	5.4	0.7	0.6	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.8	1.2	4.5	3.4	3.6	0.0	0.0	5.3	0.5	0.5	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.3	0.7	0.0	0.4	0.4	0.9	1.1	1.9	4.1	1.2	1.8	6.7	5.2	0.5	0.5	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	1.3	1.6	1.3	4.8	4.5	8.8	3.3	0.0	0.6	0.6	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	2.2	0.5	2.5	4.6	3.3	1.7	6.5	5.0	0.5	0.5	0.1
1974-78	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.1	0.2	0.3	1.9	1.1	2.1	4.3	3.2	3.7	3.4	4.1	0.6	0.6	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.0	1.2	1.3	1.4	4.5	3.7	4.3	0.0	3.1	0.0	0.6	0.6	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.3	0.3	1.1	0.0	1.7	1.3	2.3	1.9	2.9	3.1	3.2	12.1	9.7	0.7	0.7	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	1.3	1.7	4.2	3.0	2.8	8.9	3.1	0.0	0.0	0.8	0.8	0.1
1982	0.0	0.0	0.0	0.0	0.2	0.5	0.0	0.3	0.0	0.8	0.8	1.4	1.1	4.2	3.8	6.0	2.8	4.6	0.6	0.6	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.8	1.7	3.4	2.3	2.7	4.9	2.7	2.9	2.6	9.0	0.8	0.8	0.1
1979-83	0.0	0.0	0.0	0.0	0.1	0.2	0.2	0.5	0.2	1.3	1.7	2.3	2.6	3.7	4.6	3.1	4.1	4.7	0.7	0.7	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

SECONDARY AND UNSPECIFIED NEOPLASM OF LYMPH NODES (ICD9 196)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	0	1	0	1	3	0	0	5	5	1	1	0	17
1965	0	0	0	0	1	0	1	1	1	0	0	2	2	0	1	2	0	0	0	11
1966	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	3
1967	0	0	0	0	0	0	0	1	0	0	1	1	0	0	1	1	1	1	0	7
1968	0	0	0	0	0	0	0	1	0	3	0	4	0	3	3	0	1	0	0	15
1964-68	0	0	0	0	1	0	1	3	2	3	3	10	2	3	10	9	3	3	0	53
1969	0	0	0	0	0	0	0	0	1	1	4	2	3	2	1	1	0	1	0	16
1970	0	0	0	1	0	0	0	0	0	1	0	1	2	0	0	0	1	1	0	7
1971	0	0	0	0	0	0	0	0	1	2	1	4	0	4	0	0	1	1	0	14
1972	0	0	0	0	0	2	0	0	3	1	2	1	2	5	2	3	2	1	0	24
1973	0	0	0	0	0	0	0	2	3	3	5	2	4	2	2	3	0	0	1	27
1969-73	0	0	0	1	0	2	0	2	8	8	12	10	11	13	5	7	4	4	1	88
1974	0	0	0	2	0	0	0	1	2	0	1	1	2	1	2	5	1	1	0	19
1975	0	0	0	0	0	1	0	0	1	2	1	1	1	4	3	0	3	1	0	18
1976	0	0	0	0	0	0	2	1	0	0	0	0	2	3	5	4	0	3	0	20
1977	0	0	0	1	0	2	0	0	2	0	3	3	2	2	5	2	5	2	0	29
1978	0	0	0	0	0	0	0	0	2	4	4	2	3	0	1	1	0	2	0	19
1974-78	0	0	0	3	0	3	2	2	7	6	9	7	10	10	16	12	9	9	0	105
1979	0	0	1	0	0	0	0	0	2	1	2	2	2	3	4	3	1	4	0	25
1980	0	0	0	1	0	0	0	0	1	2	2	2	5	1	3	3	2	1	0	23
1981	0	0	0	1	0	0	1	0	2	1	1	6	7	1	3	4	4	3	0	34
1982	0	0	0	0	0	1	1	0	2	2	4	5	1	6	1	3	0	0	0	26
1983	0	0	0	0	1	0	1	1	1	3	4	3	4	7	6	3	5	3	0	42
1979-83	0	0	1	2	1	1	3	1	8	9	13	18	19	18	17	16	12	11	0	150

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

SECONDARY AND UNSPECIFIED NEOPLASM OF LYMPH NODES (ICD9 196)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.6	2.2	0.0	0.0	6.0	8.5	2.9	4.5	0.5	0.4	0.0
1965	0.0	0.0	0.0	0.0	0.4	0.0	0.5	0.4	0.4	0.0	0.0	1.4	1.7	0.0	1.2	3.3	0.0	0.0	0.3	0.3	0.0
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	1.6	0.0	4.2	0.1	0.1	0.0
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.6	0.7	0.0	0.0	1.1	1.5	2.6	4.0	0.2	0.2	0.0
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.4	0.0	2.5	0.0	2.7	3.3	0.0	2.5	0.0	0.4	0.4	0.1
1964-68	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.3	0.2	0.3	0.3	1.4	0.3	0.6	2.3	2.9	1.6	2.5	0.3	0.3	0.0
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	2.2	1.2	2.2	1.7	1.1	1.5	0.0	3.5	0.4	0.4	0.0
1970	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.6	1.4	0.0	0.0	0.0	2.3	3.3	0.2	0.2	0.0
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	0.5	2.3	0.0	3.3	0.0	0.0	2.2	3.1	0.3	0.3	0.0
1972	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.0	1.3	0.4	1.0	0.6	1.3	4.0	2.0	4.0	4.2	2.9	0.6	0.5	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.3	1.3	2.3	1.1	2.6	1.6	2.0	3.9	0.0	0.0	0.6	0.6	0.1
1969-73	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.2	0.7	0.7	1.2	1.2	1.5	2.1	1.0	1.9	1.7	2.5	0.4	0.4	0.0
1974	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.4	0.9	0.0	0.4	0.6	1.3	0.8	1.9	6.4	2.0	2.6	0.4	0.4	0.0
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.9	0.4	0.6	0.6	3.0	2.8	0.0	5.8	2.5	0.4	0.4	0.0
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.7	0.4	0.0	0.0	0.0	0.0	1.2	2.1	4.5	4.8	0.0	7.2	0.4	0.4	0.0
1977	0.0	0.0	0.0	0.3	0.0	0.6	0.0	0.0	0.9	0.0	1.3	1.5	1.2	1.4	4.4	2.4	9.0	4.6	0.6	0.5	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	1.7	1.7	0.9	1.7	0.0	0.8	1.1	0.0	4.5	0.4	0.4	0.0
1974-78	0.0	0.0	0.0	0.2	0.0	0.2	0.1	0.2	0.6	0.5	0.8	0.7	1.2	1.4	2.9	2.9	3.4	4.3	0.5	0.4	0.0
1979	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.8	0.4	0.9	0.9	1.2	1.9	3.3	3.3	1.7	8.5	0.5	0.4	0.0
1980	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.4	0.9	0.9	0.9	2.8	0.6	2.4	3.2	3.2	2.0	0.5	0.4	0.0
1981	0.0	0.0	0.0	0.3	0.0	0.0	0.3	0.0	0.8	0.4	0.4	2.6	3.7	0.6	2.3	4.2	6.3	5.8	0.6	0.6	0.1
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.8	0.9	1.7	2.1	0.5	3.6	0.7	3.0	0.0	0.0	0.5	0.5	0.1
1983	0.0	0.0	0.0	0.0	0.2	0.0	0.3	0.3	0.4	1.3	1.7	1.3	1.9	4.2	4.3	2.9	7.3	5.3	0.8	0.7	0.1
1979-83	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.1	0.6	0.8	1.1	1.6	2.0	2.2	2.6	3.3	3.8	4.3	0.6	0.5	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

SECONDARY MALIGNANT NEOPLASM OF RESPIRATORY AND DIGESTIVE SYSTEMS (ICD9 197)

YEAR OF DIAGNOSIS	AGE (YEARS)																		ALL AGES
	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	
1964	0	0	0	0	0	0	2	0	1	1	1	0	8	1	3	2	1	1	0	21
1965	1	0	0	0	0	0	0	0	1	2	5	2	2	3	4	4	3	1	0	28
1966	0	0	0	0	0	0	0	0	1	1	4	8	4	2	7	3	1	2	0	33
1967	0	0	0	0	0	0	0	1	0	1	2	6	5	4	6	7	0	3	0	35
1968	0	0	0	0	0	0	1	1	1	1	2	3	4	9	8	4	2	3	0	39
1964-68	1	0	0	0	0	0	3	2	4	6	14	19	23	19	28	20	7	10	0	156
1969	0	0	0	0	0	0	1	0	0	3	5	3	6	1	6	8	2	4	0	39
1970	0	0	0	0	0	0	0	0	0	3	3	4	7	10	12	7	3	2	0	51
1971	0	0	0	0	0	0	0	1	1	2	6	6	6	11	9	6	3	4	0	55
1972	0	0	0	0	0	0	0	1	1	0	5	3	4	9	5	4	3	0	0	35
1973	0	0	0	0	0	0	0	1	3	1	7	2	6	8	9	9	5	3	1	55
1969-73	0	0	0	0	0	0	1	3	5	9	26	18	29	39	41	34	16	13	1	235
1974	0	0	0	1	0	0	0	1	3	4	3	2	6	3	4	5	5	2	0	39
1975	0	0	0	0	0	1	1	1	0	1	1	6	5	11	9	6	5	6	0	53
1976	0	0	0	0	0	0	0	0	4	2	2	6	7	10	8	9	10	2	0	60
1977	0	0	0	0	1	0	1	0	1	4	4	3	3	5	11	7	4	4	0	48
1978	0	0	0	0	1	0	0	0	0	2	10	5	7	14	17	12	7	3	0	78
1974-78	0	0	0	1	2	1	2	2	8	13	20	22	28	43	49	39	31	17	0	278
1979	0	0	0	0	0	0	0	0	0	4	5	9	18	12	16	7	7	7	0	85
1980	0	0	0	0	0	0	2	0	1	0	3	6	11	17	18	6	13	5	1	83
1981	0	0	0	0	0	1	0	0	0	1	3	8	13	16	9	11	11	0	1	74
1982	0	0	0	0	0	1	0	1	4	3	6	9	14	10	14	6	7	10	2	87
1983	1	0	0	0	0	0	1	0	3	3	6	12	16	18	14	8	6	9	3	100
1979-83	1	0	0	0	0	2	3	1	8	11	23	44	72	73	71	38	44	31	7	429

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

SECONDARY MALIGNANT NEOPLASM OF RESPIRATORY AND DIGESTIVE SYSTEMS (ICD9 197)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.5	0.5	0.6	0.0	7.0	1.1	4.4	4.4	3.9	7.4	0.6	0.6	0.1
1965	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.0	2.9	1.4	1.7	3.3	5.8	8.7	11.6	7.2	0.8	0.8	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	2.2	5.4	3.3	2.1	9.9	6.4	3.8	14.3	1.0	0.9	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.5	1.1	3.9	4.1	4.2	8.5	14.7	0.0	20.7	1.0	1.0	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.5	1.1	1.9	3.2	9.3	11.3	8.4	7.5	20.0	1.1	1.1	0.1
1964-68	0.1	0.0	0.0	0.0	0.0	0.0	0.3	0.2	0.3	0.6	1.6	2.6	3.8	4.1	8.0	8.6	5.4	14.1	0.9	0.9	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	1.4	2.7	1.8	4.7	1.0	8.4	16.6	7.4	25.5	1.1	1.0	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	1.6	2.4	5.3	9.7	16.5	14.4	10.8	12.2	1.4	1.3	0.2
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.9	3.2	3.5	4.4	10.4	12.1	12.3	10.6	23.1	1.4	1.4	0.2
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	2.5	1.8	2.8	8.3	6.5	8.1	10.5	0.0	0.9	0.9	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.2	0.4	3.4	1.2	4.1	7.3	11.3	18.1	17.2	16.5	1.4	1.3	0.1
1969-73	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	0.4	0.8	2.7	2.1	4.2	7.4	10.9	13.9	11.3	15.2	1.2	1.2	0.1
1974	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	1.2	1.7	1.4	1.2	4.0	2.7	4.9	9.9	17.1	10.8	1.0	0.9	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.0	0.4	0.4	3.5	3.2	9.5	10.7	11.5	17.1	31.6	1.3	1.2	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	0.8	0.9	3.4	4.4	8.3	9.3	16.5	33.7	10.3	1.4	1.3	0.1
1977	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.0	0.4	1.7	1.8	1.6	1.9	4.0	12.5	12.4	13.2	20.3	1.1	1.1	0.1
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.8	4.4	2.5	4.4	10.8	18.7	20.4	22.7	15.0	1.7	1.7	0.2
1974-78	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.2	0.7	1.1	1.8	2.4	3.6	7.1	11.3	14.3	20.8	17.6	1.3	1.2	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	2.2	4.3	11.5	8.9	17.1	11.4	22.0	34.1	1.8	1.8	0.2
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.4	0.0	1.3	2.8	6.8	12.2	18.6	9.5	39.3	24.2	1.7	1.7	0.2
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	1.3	3.7	7.7	11.3	8.9	17.0	32.3	0.0	1.5	1.4	0.2
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.3	1.6	1.3	2.5	4.1	7.9	7.0	13.4	9.0	19.6	46.3	1.7	1.7	0.2
1983	0.3	0.0	0.0	0.0	0.0	0.0	0.3	0.0	1.1	1.3	2.5	5.4	8.5	12.6	12.8	11.7	15.9	40.4	1.9	1.9	0.2
1979-83	0.1	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.6	0.9	2.0	4.1	8.4	10.4	14.1	11.7	25.5	29.2	1.7	1.7	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

SECONDARY MALIGNANT NEOPLASM OF RESPIRATORY AND DIGESTIVE SYSTEMS (ICD9 197)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	2	1	0	2	4	1	3	1	2	5	2	0	23
1965	0	0	0	0	0	0	0	0	0	0	1	1	4	0	6	3	1	1	0	17
1966	0	0	0	1	0	0	0	0	0	2	2	1	3	4	9	3	5	1	0	31
1967	0	0	0	0	0	1	0	1	3	3	2	3	6	2	6	5	7	6	0	45
1968	0	0	0	0	0	0	0	0	0	0	4	8	2	5	6	3	1	1	0	30
1964-68	0	0	0	1	0	1	0	3	4	5	11	17	16	14	28	16	19	11	0	146
1969	0	0	0	0	0	0	0	0	1	1	4	6	1	9	10	7	2	6	0	47
1970	0	0	0	0	0	1	0	3	1	1	2	7	5	2	6	5	6	13	0	52
1971	0	0	0	0	0	0	3	1	0	5	5	4	5	1	8	9	9	6	0	56
1972	0	0	0	0	0	0	0	0	1	5	2	8	9	9	8	10	10	4	0	66
1973	0	0	0	2	0	0	1	1	2	3	6	9	8	8	8	12	3	4	1	68
1969-73	0	0	0	2	0	1	4	5	5	15	19	34	28	29	40	43	30	33	1	289
1974	0	0	0	0	0	0	0	0	2	7	4	5	5	7	7	6	6	5	0	54
1975	0	0	0	0	0	0	0	1	3	4	6	9	12	7	11	9	11	4	1	78
1976	0	0	0	0	1	1	1	1	2	2	5	1	8	4	10	14	16	5	0	71
1977	0	0	0	0	0	0	0	1	1	0	6	7	5	6	11	9	12	8	0	66
1978	0	0	0	0	0	0	2	2	2	2	3	5	9	8	15	16	13	8	1	86
1974-78	0	0	0	0	1	1	3	5	10	15	24	27	39	32	54	54	58	30	2	355
1979	0	0	0	0	0	2	0	2	1	4	3	5	8	11	12	11	14	15	1	89
1980	0	0	0	0	0	1	1	1	1	3	7	7	7	17	14	17	9	9	0	94
1981	0	0	0	1	0	0	2	1	2	1	4	8	11	6	17	11	16	7	1	88
1982	0	0	0	0	1	1	0	2	2	3	5	8	5	11	19	21	20	17	1	116
1983	0	0	0	0	0	1	1	2	1	3	4	8	11	20	26	20	17	22	1	137
1979-83	0	0	0	1	1	5	4	8	7	14	23	36	42	65	88	80	76	70	4	524

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

SECONDARY MALIGNANT NEOPLASM OF RESPIRATORY AND DIGESTIVE SYSTEMS (ICD9 197)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.5	0.0	1.2	2.9	0.8	3.0	1.2	3.4	14.5	9.1	0.7	0.6	0.1
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.7	3.3	0.0	7.0	5.0	2.8	4.3	0.5	0.4	0.1
1966	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	1.0	1.1	0.7	2.4	3.8	10.2	4.8	13.4	4.2	0.9	0.7	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.4	1.2	1.5	1.1	2.0	4.7	1.8	6.7	7.7	17.9	23.9	1.3	1.0	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.2	5.1	1.5	4.5	6.6	4.5	2.5	3.8	0.8	0.7	0.1
1964-68	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.3	0.3	0.5	1.3	2.3	2.6	2.6	6.4	5.1	10.1	9.1	0.8	0.7	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	2.2	3.7	0.7	7.9	10.8	10.2	4.7	21.2	1.2	1.0	0.1
1970	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.3	0.4	0.4	1.1	4.2	3.6	1.7	6.3	7.1	13.6	42.9	1.3	1.1	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.4	0.0	2.1	2.6	2.3	3.5	0.8	8.2	12.5	19.5	18.5	1.4	1.1	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	2.1	1.0	4.6	6.0	7.2	8.1	13.5	21.1	11.7	1.6	1.3	0.1
1973	0.0	0.0	0.0	0.5	0.0	0.0	0.4	0.4	0.9	1.3	2.8	5.1	5.2	6.2	7.9	15.8	6.1	11.1	1.6	1.4	0.2
1969-73	0.0	0.0	0.0	0.1	0.0	0.1	0.3	0.4	0.4	1.3	1.9	4.0	3.9	4.8	8.2	11.9	13.1	20.5	1.4	1.2	0.1
1974	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	3.0	1.8	2.8	3.1	5.3	6.7	7.7	12.0	13.2	1.2	1.1	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.3	1.7	2.6	5.0	7.2	5.2	10.2	11.2	21.3	10.2	1.7	1.4	0.2
1976	0.0	0.0	0.0	0.0	0.3	0.3	0.3	0.4	0.9	0.9	2.1	0.5	4.7	2.8	9.0	16.9	30.0	12.1	1.5	1.2	0.1
1977	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.0	2.5	3.5	2.9	4.1	9.6	10.6	21.7	18.5	1.4	1.1	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.8	0.9	0.9	1.3	2.3	5.2	5.3	12.6	18.3	22.7	17.8	1.7	1.3	0.1
1974-78	0.0	0.0	0.0	0.0	0.1	0.1	0.2	0.4	0.9	1.3	2.0	2.8	4.7	4.5	9.7	13.1	21.7	14.5	1.5	1.2	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.7	0.4	1.7	1.3	2.2	4.6	7.1	9.8	12.2	23.5	31.8	1.7	1.4	0.1
1980	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.4	1.3	3.0	3.0	3.9	10.6	11.1	18.2	14.6	18.3	1.8	1.5	0.2
1981	0.0	0.0	0.0	0.3	0.0	0.0	0.5	0.3	0.8	0.4	1.7	3.4	5.9	3.6	13.0	11.4	25.0	13.6	1.6	1.3	0.2
1982	0.0	0.0	0.0	0.0	0.2	0.3	0.0	0.6	0.8	1.3	2.1	3.4	2.5	6.6	14.1	21.0	30.6	31.5	2.0	1.5	0.2
1983	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.6	0.4	1.3	1.7	3.4	5.2	12.0	18.6	19.3	25.0	38.7	2.3	1.8	0.2
1979-83	0.0	0.0	0.0	0.1	0.1	0.3	0.2	0.5	0.6	1.2	2.0	3.1	4.4	8.0	13.5	16.5	23.8	27.1	1.9	1.5	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

SECONDARY MALIGNANT NEOPLASM OF OTHER SPECIFIED SITES (ICD9 198)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	1	0	0	1	0	1	3	5	4	6	5	5	1	2	2	0	36
1965	0	0	0	0	1	1	0	0	2	2	4	3	3	7	3	3	1	1	0	31
1966	0	0	0	0	0	0	0	0	4	4	2	4	1	5	0	5	2	1	0	28
1967	1	0	0	0	0	0	0	0	0	1	1	4	1	3	6	2	0	0	0	19
1968	0	0	0	0	1	0	0	0	0	1	1	2	2	1	2	2	1	1	0	14
1964-68	1	0	0	1	2	1	1	0	7	11	13	17	13	21	16	13	6	5	0	128
1969	0	0	0	0	0	0	1	2	2	2	3	4	4	4	5	4	5	0	1	37
1970	0	0	0	0	1	0	0	0	1	3	1	1	6	6	5	5	6	0	0	35
1971	0	0	0	0	0	0	0	0	0	1	2	9	6	8	3	3	3	3	1	39
1972	0	0	0	0	0	0	0	0	1	1	4	0	8	9	8	1	3	7	0	42
1973	0	0	0	0	0	0	0	0	1	2	4	1	4	5	7	8	7	4	0	43
1969-73	0	0	0	0	1	0	1	2	5	9	14	15	28	32	28	21	24	14	2	196
1974	2	0	0	0	0	0	0	1	0	0	1	2	6	7	8	9	1	2	3	42
1975	0	0	0	0	0	1	3	0	3	3	5	8	8	7	5	3	1	1	1	49
1976	0	0	0	0	0	0	0	1	1	4	2	6	9	5	6	6	2	4	0	46
1977	0	0	0	1	0	0	0	1	1	1	2	3	5	4	3	6	1	2	0	30
1978	0	0	0	0	0	0	0	1	1	2	5	8	7	13	10	6	3	4	0	60
1974-78	2	0	0	1	0	1	3	4	6	10	15	27	35	36	32	30	8	13	4	227
1979	0	0	0	0	0	0	0	3	0	2	6	7	9	10	9	12	5	4	0	67
1980	0	0	0	0	0	0	0	3	1	1	4	3	10	12	9	9	5	1	0	58
1981	0	0	0	0	0	0	1	1	0	2	5	4	12	8	16	7	3	6	1	66
1982	1	0	0	0	0	0	0	0	1	4	5	8	9	9	13	13	1	3	1	68
1983	0	0	0	0	0	0	0	0	2	0	6	6	19	9	17	12	9	4	0	84
1979-83	1	0	0	0	0	0	1	7	4	9	26	28	59	48	64	53	23	18	2	343

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

SECONDARY MALIGNANT NEOPLASM OF OTHER SPECIFIED SITES (ICD9 198)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.4	0.0	0.0	0.4	0.0	0.5	1.6	2.9	2.9	5.3	5.6	7.3	2.2	7.9	14.7	1.1	1.1	0.1
1965	0.0	0.0	0.0	0.0	0.4	0.5	0.0	0.0	0.9	1.0	2.3	2.1	2.6	7.7	4.3	6.5	3.9	7.2	0.9	0.9	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.7	2.0	1.1	2.7	0.8	5.4	0.0	10.7	7.7	7.1	0.8	0.8	0.1
1967	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.5	2.6	0.8	3.2	8.5	4.2	0.0	0.0	0.5	0.5	0.1
1968	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.5	0.5	1.3	1.6	1.0	2.8	4.2	3.7	6.7	0.4	0.4	0.0
1964-68	0.1	0.0	0.0	0.1	0.2	0.1	0.1	0.0	0.6	1.1	1.5	2.3	2.2	4.5	4.6	5.6	4.6	7.0	0.7	0.7	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.8	0.8	0.9	1.6	2.5	3.1	4.0	7.0	8.3	18.4	0.0	1.0	0.9	0.1
1970	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.4	1.3	0.5	0.6	4.5	5.8	6.9	10.3	21.6	0.0	0.9	0.9	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.1	5.3	4.4	7.5	4.0	6.1	10.6	17.3	1.0	1.0	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	2.0	0.0	5.7	8.3	10.4	2.0	10.5	39.1	1.1	1.1	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.9	1.9	0.6	2.7	4.5	8.8	16.1	24.1	22.0	1.1	1.0	0.1
1969-73	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.2	0.4	0.8	1.5	1.8	4.1	6.1	7.5	8.6	17.0	16.4	1.0	1.0	0.1
1974	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.0	0.5	1.2	4.0	6.2	9.7	17.8	3.4	10.8	1.0	1.0	0.1
1975	0.0	0.0	0.0	0.0	0.0	0.3	1.0	0.0	1.2	1.3	2.2	4.7	5.2	6.0	5.9	5.7	3.4	5.3	1.1	1.1	0.1
1976	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	1.7	0.9	3.4	5.7	4.1	6.9	11.0	6.7	20.6	1.1	1.1	0.1
1977	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.4	0.4	0.4	0.9	1.6	3.2	3.2	3.4	10.6	3.3	10.2	0.7	0.7	0.1
1978	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.8	2.2	4.0	4.4	10.0	11.0	10.2	9.7	20.0	1.3	1.3	0.2
1974-78	0.1	0.0	0.0	0.0	0.0	0.1	0.2	0.3	0.5	0.8	1.3	3.0	4.5	5.9	7.4	11.0	5.4	13.5	1.0	1.0	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.8	2.6	3.4	5.7	7.4	9.6	19.6	15.7	19.5	1.5	1.4	0.2
1980	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.4	0.4	1.7	1.4	6.2	8.6	9.3	14.3	15.1	4.8	1.2	1.2	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.8	2.1	1.9	7.1	5.7	15.9	10.8	8.8	28.3	1.3	1.3	0.2
1982	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.7	2.1	3.7	5.1	6.3	12.4	19.5	2.8	13.9	1.4	1.3	0.2
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	2.5	2.7	10.1	6.3	15.6	17.5	23.8	17.9	1.6	1.6	0.2
1979-83	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.5	0.3	0.8	2.2	2.6	6.9	6.9	12.7	16.4	13.3	16.9	1.4	1.4	0.2

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

SECONDARY MALIGNANT NEOPLASM OF OTHER SPECIFIED SITES (ICD9 198)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	0	0	0	0	1	1	1	1	4	3	1	0	3	1	1	0	17
1965	0	0	0	0	0	0	0	0	0	3	2	2	3	2	3	3	2	0	0	20
1966	0	0	0	0	0	0	1	0	1	0	1	2	5	3	5	1	4	0	0	23
1967	0	0	0	0	0	0	0	0	2	0	1	2	3	4	4	0	0	2	0	18
1968	0	0	0	0	0	0	0	0	0	2	4	8	0	3	4	0	1	1	0	23
1964-68	0	0	0	0	0	0	1	1	4	6	9	18	14	13	16	7	8	4	0	101
1969	0	0	0	0	0	0	0	0	2	1	0	3	0	0	3	3	1	2	0	15
1970	0	0	0	0	0	0	0	2	2	2	1	4	0	2	3	4	4	1	0	25
1971	0	0	0	0	0	0	0	3	0	1	1	3	2	4	5	4	2	3	0	28
1972	0	0	0	0	0	1	0	0	1	1	1	2	5	6	1	4	9	4	0	35
1973	0	0	0	0	0	1	0	0	6	0	4	4	5	5	7	7	2	10	1	52
1969-73	0	0	0	0	0	2	0	5	11	5	7	16	12	17	19	22	18	20	1	155
1974	0	0	0	0	1	0	0	1	0	4	2	3	5	4	5	7	4	3	0	39
1975	1	0	0	0	1	0	1	0	1	1	4	5	3	10	11	6	3	4	1	52
1976	0	0	0	1	1	1	1	0	0	2	2	3	4	10	5	7	3	6	1	47
1977	0	0	0	0	0	0	1	2	2	4	3	0	4	5	6	5	1	2	0	35
1978	0	0	0	0	1	0	0	2	3	3	8	4	5	5	9	9	6	2	1	58
1974-78	1	0	0	1	4	1	3	5	6	14	19	15	21	34	36	34	17	17	3	231
1979	0	0	0	0	0	0	2	0	2	4	3	7	4	7	5	12	6	7	0	59
1980	0	0	0	1	0	0	0	1	3	2	3	2	5	4	7	8	7	8	0	51
1981	0	0	0	0	0	1	1	1	0	4	7	9	10	10	12	8	9	3	0	75
1982	0	0	0	0	0	1	0	3	1	2	4	6	13	3	18	11	11	8	2	83
1983	0	0	0	0	0	0	0	1	1	2	5	7	12	11	13	10	13	14	1	90
1979-83	0	0	0	1	0	2	3	6	7	14	22	31	44	35	55	49	46	40	3	358

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

SECONDARY MALIGNANT NEOPLASM OF OTHER SPECIFIED SITES (ICD9 198)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.5	0.5	0.6	2.9	2.5	1.0	0.0	5.1	2.9	4.5	0.5	0.5	0.0
1965	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	1.2	1.4	2.5	1.9	3.5	5.0	5.6	0.0	0.6	0.5	0.1
1966	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.4	0.0	0.6	1.4	4.0	2.8	5.6	1.6	10.8	0.0	0.7	0.6	0.1
1967	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.6	1.3	2.3	3.7	4.4	0.0	0.0	8.0	0.5	0.5	0.1
1968	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	2.2	5.1	0.0	2.7	4.4	0.0	2.5	3.8	0.6	0.6	0.1
1964-68	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.3	0.6	1.0	2.4	2.2	2.4	3.6	2.2	4.3	3.3	0.6	0.5	0.1
1969	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.5	0.0	1.8	0.0	0.0	3.2	4.4	2.4	7.1	0.4	0.3	0.0
1970	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.9	0.9	0.5	2.4	0.0	1.7	3.2	5.7	9.0	3.3	0.6	0.5	0.1
1971	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.3	0.0	0.4	0.5	1.7	1.4	3.3	5.1	5.5	4.3	9.2	0.7	0.6	0.1
1972	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.4	0.5	1.1	3.4	4.8	1.0	5.4	18.9	11.7	0.8	0.7	0.1
1973	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	2.6	0.0	1.9	2.3	3.2	3.9	6.9	9.2	4.1	27.9	1.2	1.0	0.1
1969-73	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.4	0.9	0.4	0.7	1.9	1.7	2.8	3.9	6.1	7.8	12.4	0.8	0.6	0.1
1974	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.4	0.0	1.7	0.9	1.7	3.1	3.0	4.8	9.0	8.0	7.9	0.9	0.7	0.1
1975	0.3	0.0	0.0	0.0	0.3	0.0	0.4	0.0	0.4	0.4	1.7	2.8	1.8	7.4	10.2	7.5	5.8	10.2	1.1	1.0	0.1
1976	0.0	0.0	0.0	0.3	0.3	0.3	0.3	0.0	0.0	0.9	0.8	1.6	2.4	7.1	4.5	8.5	5.6	14.5	1.0	0.8	0.1
1977	0.0	0.0	0.0	0.0	0.3	0.0	0.3	0.8	0.9	1.7	1.3	0.0	2.3	3.4	5.2	5.9	1.8	4.6	0.7	0.7	0.1
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.8	1.3	1.3	3.4	1.9	2.9	3.3	7.6	10.3	10.5	4.5	1.2	1.0	0.1
1974-78	0.1	0.0	0.0	0.1	0.2	0.1	0.2	0.4	0.5	1.2	1.6	1.6	2.5	4.8	6.5	8.2	6.4	8.2	1.0	0.8	0.1
1979	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.8	1.7	1.3	3.1	2.3	4.5	4.1	13.3	10.1	14.9	1.2	0.9	0.1
1980	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.4	1.3	0.9	1.3	0.9	2.8	2.5	5.5	8.6	11.3	16.3	1.0	0.8	0.1
1981	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.3	0.0	1.7	3.0	3.9	5.3	6.1	9.2	8.3	14.1	5.8	1.4	1.2	0.2
1982	0.0	0.0	0.0	0.0	0.0	0.3	0.0	1.0	0.4	0.9	1.7	2.6	6.6	1.8	13.4	11.0	16.8	14.8	1.5	1.2	0.1
1983	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.4	0.9	2.1	3.0	5.7	6.6	9.3	9.6	19.1	24.6	1.5	1.2	0.1
1979-83	0.0	0.0	0.0	0.1	0.0	0.1	0.2	0.4	0.6	1.2	1.9	2.7	4.6	4.3	8.4	10.1	14.4	15.5	1.3	1.1	0.1

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: MALES

MALIGNANT NEOPLASM WITHOUT SPECIFICATION OF SITE (ICD9 199)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	1	0	0	0	1	2	3	3	4	22	20	25	27	29	22	16	7	0	182
1965	0	1	0	0	1	3	2	4	3	7	20	20	31	25	46	17	27	10	0	217
1966	4	1	1	0	1	1	0	1	4	6	18	22	24	40	33	27	14	6	0	203
1967	3	0	0	0	0	2	1	1	2	10	13	19	20	27	26	25	12	16	0	177
1968	2	0	1	0	0	1	1	0	6	9	6	14	28	23	34	24	21	7	0	177
1964-68	9	3	2	0	2	8	6	9	18	36	79	95	128	142	168	115	90	46	0	956
1969	0	0	0	0	0	2	0	2	3	8	7	18	21	32	20	21	19	10	0	163
1970	0	0	0	1	0	0	2	0	0	4	8	17	20	27	19	23	17	10	0	148
1971	2	0	0	0	0	0	0	1	3	8	9	15	26	32	22	25	10	10	0	163
1972	3	0	0	1	0	0	0	1	4	6	11	19	31	25	45	27	15	12	0	200
1973	2	1	0	1	0	0	1	3	3	6	11	22	27	33	43	38	17	15	1	224
1969-73	7	1	0	3	0	2	3	7	13	32	46	91	125	149	149	134	78	57	1	898
1974	2	0	1	1	0	3	3	7	5	7	20	20	27	39	53	41	21	25	4	279
1975	0	0	0	1	2	1	2	3	5	12	19	21	31	29	44	27	27	19	5	248
1976	1	0	1	2	3	3	2	2	10	6	23	21	24	51	51	45	32	16	0	293
1977	1	0	1	2	1	5	4	3	8	12	27	20	34	46	39	43	25	21	0	292
1978	0	0	1	1	2	0	1	4	0	8	24	33	52	65	46	49	32	19	1	338
1974-78	4	0	4	7	8	12	12	19	28	45	113	115	168	230	233	205	137	100	10	1450
1979	2	0	1	3	1	0	0	0	3	7	24	22	28	46	62	42	31	23	0	295
1980	1	0	0	3	0	2	3	2	6	7	23	18	29	53	48	50	30	31	0	306
1981	2	1	1	1	1	4	1	3	2	6	14	28	37	45	50	41	39	23	0	299
1982	0	0	0	1	0	0	1	3	3	5	21	29	23	56	53	35	35	28	4	297
1983	1	2	1	0	2	2	1	0	4	10	16	36	19	44	43	41	36	21	1	280
1979-83	6	3	3	8	4	8	6	8	18	35	98	133	136	244	256	209	171	126	5	1477

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: MALES

MALIGNANT NEOPLASM WITHOUT SPECIFICATION OF SITE (ICD9 199)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.3	0.0	0.0	0.0	0.5	0.9	1.3	1.4	2.1	12.9	14.3	21.9	30.0	42.5	48.5	63.0	51.5	5.5	5.3	0.6
1965	0.0	0.3	0.0	0.0	0.4	1.4	0.9	1.7	1.3	3.7	11.4	14.0	26.6	27.3	66.3	37.0	104.2	71.9	6.4	6.2	0.8
1966	1.0	0.3	0.3	0.0	0.4	0.5	0.0	0.4	1.7	3.1	10.1	15.0	20.1	43.0	46.9	57.9	53.8	42.9	5.9	5.8	0.7
1967	0.8	0.0	0.0	0.0	0.0	0.9	0.4	0.4	0.8	5.0	7.1	12.5	16.3	28.4	36.7	52.6	45.5	110.3	5.1	5.0	0.5
1968	0.6	0.0	0.3	0.0	0.0	0.4	0.4	0.0	2.5	4.3	3.3	8.9	22.2	23.7	47.8	50.2	78.4	46.7	5.0	4.8	0.6
1964-68	0.5	0.2	0.1	0.0	0.2	0.7	0.5	0.8	1.6	3.7	8.9	12.9	21.4	30.4	48.0	49.3	69.0	64.8	5.6	5.4	0.6
1969	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.8	1.2	3.7	3.8	11.1	16.3	32.0	27.9	43.7	69.9	63.7	4.5	4.3	0.5
1970	0.0	0.0	0.0	0.3	0.0	0.0	0.8	0.0	0.0	1.8	4.3	10.2	15.1	26.2	26.1	47.4	61.2	61.0	4.0	3.8	0.4
1971	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.4	1.2	3.4	4.8	8.8	19.1	30.2	29.6	51.1	35.5	57.8	4.3	4.2	0.5
1972	0.9	0.0	0.0	0.3	0.0	0.0	0.0	0.4	1.6	2.6	5.5	11.1	22.0	23.1	58.7	54.7	52.3	67.0	5.2	5.0	0.6
1973	0.6	0.3	0.0	0.3	0.0	0.0	0.4	1.3	1.2	2.6	5.3	13.0	18.6	30.0	54.0	76.3	58.4	82.4	5.7	5.4	0.6
1969-73	0.4	0.1	0.0	0.2	0.0	0.1	0.2	0.6	1.1	2.8	4.8	10.8	18.3	28.3	39.7	54.8	55.3	66.7	4.7	4.6	0.5
1974	0.6	0.0	0.2	0.3	0.0	0.9	1.1	2.9	2.0	3.0	9.1	11.9	18.0	34.5	64.4	81.0	71.9	135.1	6.8	6.6	0.7
1975	0.0	0.0	0.0	0.2	0.6	0.3	0.7	1.2	2.1	5.0	8.5	12.2	20.0	25.0	52.1	51.5	92.5	100.0	5.9	5.6	0.6
1976	0.3	0.0	0.2	0.5	0.8	0.8	0.7	0.8	4.2	2.5	10.1	11.7	15.2	42.3	59.0	82.7	107.7	82.5	7.0	6.5	0.7
1977	0.3	0.0	0.3	0.5	0.3	1.4	1.3	1.2	3.3	5.0	11.9	10.6	21.5	36.5	44.2	76.1	82.8	106.6	6.7	6.5	0.7
1978	0.0	0.0	0.3	0.2	0.5	0.0	0.3	1.5	0.0	3.3	10.5	16.6	33.0	50.0	50.7	83.2	103.9	95.0	7.6	7.2	0.8
1974-78	0.3	0.0	0.2	0.3	0.4	0.7	0.8	1.5	2.3	3.8	10.0	12.7	21.6	38.0	53.9	75.1	91.9	103.5	6.8	6.5	0.7
1979	0.6	0.0	0.3	0.7	0.3	0.0	0.0	0.0	1.2	2.9	10.4	10.6	17.9	34.2	66.2	68.6	97.5	112.2	6.5	6.2	0.7
1980	0.3	0.0	0.0	0.7	0.0	0.6	0.9	0.7	2.5	2.9	9.9	8.5	18.0	38.2	49.5	79.5	90.6	149.8	6.5	6.2	0.7
1981	0.7	0.3	0.3	0.2	0.3	1.1	0.3	1.0	0.8	2.5	5.9	13.0	21.9	31.9	49.7	63.4	114.4	108.5	6.2	5.9	0.6
1982	0.0	0.0	0.0	0.2	0.0	0.0	0.3	1.0	1.2	2.1	8.9	13.3	12.9	39.2	50.7	52.6	98.0	129.6	5.9	5.6	0.6
1983	0.3	0.6	0.3	0.0	0.5	0.5	0.3	0.0	1.5	4.2	6.7	16.3	10.1	30.9	39.4	59.9	95.2	94.2	5.6	5.2	0.6
1979-83	0.4	0.2	0.2	0.4	0.2	0.4	0.3	0.5	1.4	2.9	8.4	12.4	16.0	34.9	50.7	64.5	99.1	118.5	6.1	5.8	0.6

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE FREQUENCIES: FEMALES

MALIGNANT NEOPLASM WITHOUT SPECIFICATION OF SITE (ICD9 199)

YEAR OF DIAGNOSIS	AGE (YEARS) 0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	UNK	ALL AGES
1964	0	0	0	1	0	0	1	6	11	4	11	16	20	19	11	14	15	2	0	131
1965	1	0	1	0	0	1	2	3	5	7	11	15	14	15	9	8	13	13	0	118
1966	0	1	0	0	2	2	2	1	5	6	13	21	14	21	17	17	21	9	0	152
1967	1	1	1	1	1	1	0	7	10	8	11	10	17	11	16	17	13	12	0	138
1968	1	0	0	0	1	1	2	2	5	6	12	16	16	13	12	16	15	13	0	131
1964-68	3	2	2	2	4	5	7	19	36	31	58	78	81	79	65	72	77	49	0	670
1969	0	0	0	0	1	2	0	2	2	9	5	7	13	10	18	8	15	16	0	108
1970	1	0	0	0	0	1	0	5	7	9	5	15	13	14	11	14	15	8	0	118
1971	0	0	0	0	0	2	0	2	6	2	13	9	10	4	15	23	12	21	0	119
1972	1	0	0	1	1	1	1	4	5	6	10	16	17	15	13	20	18	17	1	147
1973	1	0	0	1	1	3	2	4	6	7	7	21	18	17	20	16	23	9	4	160
1969-73	3	0	0	2	3	9	3	17	26	33	40	68	71	60	77	81	83	71	5	652
1974	1	0	1	0	2	3	3	4	6	14	11	14	19	28	21	29	23	22	3	204
1975	1	0	0	1	1	1	1	1	9	3	11	17	24	17	24	27	24	20	0	182
1976	1	0	2	2	1	3	4	4	4	8	14	21	28	27	20	32	34	26	0	231
1977	1	0	1	1	2	0	2	4	8	6	13	14	27	34	21	25	24	25	0	208
1978	0	0	0	0	1	0	2	3	3	11	12	27	32	24	24	32	37	28	0	236
1974-78	4	0	4	4	7	7	12	16	30	42	61	93	130	130	110	145	142	121	3	1061
1979	2	0	0	0	2	2	4	0	2	4	9	15	11	23	23	16	25	28	0	166
1980	0	0	1	1	2	1	1	2	5	8	10	17	24	26	23	38	31	37	0	227
1981	2	0	0	1	1	1	5	0	0	6	14	17	16	27	26	34	27	38	0	215
1982	2	0	0	2	1	0	3	5	2	7	8	10	22	23	24	32	19	38	0	198
1983	2	0	0	2	0	0	0	1	7	6	14	10	22	25	27	33	38	35	1	223
1979-83	8	0	1	6	6	4	13	8	16	31	55	69	95	124	123	153	140	176	1	1029

SEE TEXT SECTION 'INTERPRETATION OF DATA'

ONTARIO CANCER REGISTRY

INCIDENCE RATES PER 100,000: FEMALES

MALIGNANT NEOPLASM WITHOUT SPECIFICATION OF SITE (ICD9 199)

	AGE SPECIFIC RATES																	SUMMARY RATES		
YEAR	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85+	AAR CAN	AAR WORLD	CUM RISK % 0-74
1964	0.0	0.0	0.0	0.4	0.4	0.0	0.5	2.6	5.0	2.2	6.7	11.6	16.9	18.7	13.3	23.9	43.4	9.1	3.9	3.4	0.4
1965	0.3	0.0	0.3	0.0	0.0	0.5	0.9	1.3	2.2	3.7	6.5	10.6	11.6	14.5	10.5	13.2	36.2	56.5	3.5	3.0	0.3
1966	0.0	0.3	0.0	0.0	0.8	0.9	0.9	0.4	2.1	3.1	7.5	14.4	11.2	19.8	19.2	27.5	56.5	38.0	4.4	3.7	0.4
1967	0.3	0.3	0.3	0.3	0.4	0.4	0.0	3.0	4.2	3.9	6.2	6.6	13.2	10.1	17.8	26.3	33.3	47.8	3.8	3.3	0.3
1968	0.3	0.0	0.0	0.0	0.4	0.4	0.9	0.9	2.1	2.8	6.6	10.2	12.1	11.7	13.2	23.9	36.9	49.1	3.6	3.0	0.3
1964-68	0.2	0.1	0.1	0.1	0.3	0.5	0.6	1.6	3.1	3.1	6.7	10.6	13.0	14.9	14.8	23.0	41.1	40.7	3.8	3.3	0.3
1969	0.0	0.0	0.0	0.0	0.3	0.8	0.0	0.9	0.8	4.1	2.7	4.3	9.6	8.7	19.4	11.6	35.4	56.5	2.8	2.4	0.3
1970	0.3	0.0	0.0	0.0	0.0	0.4	0.0	2.2	3.0	3.9	2.7	8.9	9.3	11.9	11.6	19.9	33.9	26.4	3.0	2.6	0.3
1971	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.9	2.6	0.8	6.8	5.2	6.9	3.3	15.4	31.9	26.0	64.6	2.9	2.3	0.2
1972	0.3	0.0	0.0	0.3	0.3	0.3	0.4	1.7	2.2	2.5	4.9	9.2	11.4	12.0	13.1	26.9	37.9	49.6	3.5	2.9	0.3
1973	0.3	0.0	0.0	0.3	0.3	0.9	0.8	1.7	2.6	3.0	3.2	12.0	11.7	13.2	19.6	21.1	46.9	25.1	3.7	3.1	0.3
1969-73	0.2	0.0	0.0	0.1	0.2	0.6	0.3	1.5	2.2	2.8	4.1	8.0	9.8	9.9	15.9	22.4	36.2	44.0	3.2	2.7	0.3
1974	0.3	0.0	0.3	0.0	0.6	0.9	1.1	1.7	2.6	5.9	4.8	7.9	11.9	21.1	20.0	37.2	45.9	58.0	4.6	3.8	0.4
1975	0.3	0.0	0.0	0.3	0.3	0.3	0.4	0.4	3.9	1.3	4.7	9.4	14.5	12.6	22.2	33.7	46.5	50.8	4.0	3.3	0.4
1976	0.3	0.0	0.5	0.5	0.3	0.8	1.[illegible]	1.6	1.7	3.4	5.9	11.0	16.6	19.2	18.0	38.7	63.8	62.8	5.0	4.1	0.4
1977	0.3	0.0	0.3	0.3	0.5	0.0	0.6	1.6	3.4	2.6	5.5	6.9	15.8	23.4	18.3	29.5	43.3	57.7	4.3	3.7	0.4
1978	0.0	0.0	0.0	0.0	0.3	0.0	0.6	1.1	1.3	4.7	5.1	12.6	18.6	16.0	20.2	36.6	64.7	62.4	4.8	3.9	0.4
1974-78	0.3	0.0	0.2	0.2	0.4	0.4	0.8	1.3	2.6	3.6	5.2	9.6	15.5	18.5	19.7	35.1	53.1	58.5	4.5	3.8	0.4
1979	0.7	0.0	0.0	0.0	0.5	0.5	1.2	0.0	0.8	1.7	3.9	6.7	6.3	14.8	18.7	17.7	41.9	59.4	3.2	2.6	0.3
1980	0.0	0.0	0.3	0.2	0.5	0.3	0.3	0.7	2.1	3.4	4.3	7.4	13.4	16.1	18.2	40.7	50.2	75.4	4.3	3.4	0.3
1981	0.7	0.0	0.0	0.3	0.3	0.3	1.4	0.0	0.0	2.6	6.0	7.3	8.5	16.4	20.0	35.3	42.3	73.8	3.9	3.1	0.3
1982	0.7	0.0	0.0	0.5	0.2	0.0	0.8	1.6	0.8	3.0	3.4	4.3	11.1	13.8	17.9	32.0	29.1	70.4	3.5	2.9	0.3
1983	0.7	0.0	0.0	0.5	0.0	0.0	0.0	0.3	2.7	2.6	6.0	4.3	10.5	15.0	19.3	31.8	55.8	61.5	3.9	3.1	0.3
1979-83	0.5	0.0	0.1	0.3	0.3	0.2	0.7	0.5	1.3	2.7	4.7	6.0	10.0	15.2	18.8	31.6	43.9	68.1	3.8	3.0	0.3

SEE TEXT SECTION 'INTERPRETATION OF DATA'

Index

AAR (Age Adjusted Rate)
  Canada
    See site of interest
  Explanation and sample calculation, 12-13
  World
    By sites, 5 year summary, 20-21
    For individual years, see site of interest
Advantages and limitations of data, 7-8
Age adjusted rate
  See AAR
Age specific rate
  Calculation of, 12
  See site of interest or ASR
Age standardized rate
  See AAR
All sites (ICD9: 140-208, excl. 173), 24-27
Anus
  See Rectum
Articular cartilage
  See Bone
Artifacts in data, 8, 11
ASR (Age Specific Rate)
  Explanation and sample calculation of, 12
  For specific site, see site of interest

Bladder (ICD9: 188), 166-169
Body of uterus
  See Uterus
Bone (ICD9: 170), 132-135
Brain (ICD9: 191), 178-181
Breast (ICD9: 174/5), 144-147
Bronchus
  See Lung

Cancer Act, 1
Cancer by region in Ontario, 14
Case resolution of linked records
  Composite record, 7
  Logic used, 5-7
  of Metastatic cancers, 6-7
  of Multiple primary cancers, 7
  Quality control in, 7
  Reliability of sources, 6
  Resolution of site, 5-7
  Same histology codes, 6
  Use of histology in determining site codes, 6-7
Cervix uteri (ICD9: 180)
  High rates, 11
  Tables, 150-151
Characteristics of OCR data
  Death certificate only, 11
  Multiple primaries, 7-8
  Non-specific sites, high proportion of, 11
  Childhood cancer, 11
Coding
  Changes over time, 3-4
  Conversion of ICD codes, 4
  Quality control, 3-4, 8
Colon (ICD9: 153), 80-83
Completeness of data, 7-8, 11
Composite case records, 7
Composite identification records, 7
Connective tissue (ICD9: 171), 136-139
Cost of registry, 7-8
Creation of incidence data
  Linkage of cases, 4-5
  Preprocessing, 3-4
  Resolution of cancers, 5-7
CUM RISK% $_{0-74}$
  For specific site see site of interest
  Explanation and sample calculation, 12-13
Cumulative Risk%
  See CUM RISK% $_{0-74}$

Data
  Description summary (n, %DIST, AAR WORLD), 20-21
  Production
  Quality indicators (HV%, DCO%, M/I%), 22-23
Data sources
  Abstract cards, 2-3
  Additional sources, 8
  Death registration transcripts, 2-3
  Hospital separation forms (hospital discharge), 2-3
  Number of records by year and source type, 3
  OCF drug service programme, 2-3
  OCF biopsy service reports, 2-3
  Pathology reports, 2-3
  Quality checks, 3-4
  Radiotherapy outpatient reports, 2-3
Death Certificate Only
  See DCO%
Differences from other registries, 8, 11
Digestive organs
  Other and ill defined sites within the digestive organs (ICD9: 159), 104-107
  Secondary malignant neoplasms of respiratory and digestive organs (ICD9: 197), 238-241
Distribution of cancers by site
  See %Dist
%Dist (percent distribution of cancers)
  By site, 20-21
DCO% (percent Death Certificate Only)
  Definition of, 11
  By site, 22-23
Duodenum
  See Small intestine
Duplicate reporting of cancers, 8, 11

Endocrine glands (ICD9: 194), 190-193
Extrahepatic bile ducts
  See Gallbladder
Eye (ICD9: 190), 174-177

Fallopian tubes
  See Ovary
Floor of mouth (ICD9: 144), 44-47
  See also Mouth, Naso-, Oro-, Hypopharynx
Frequency
  See n

Gallbladder (ICD9: 156), 92-95
Generalized iterative record linkage system
See GIRLS
Genital organs
Other and unspecified female (ICD9: 184), 158-159
Penis, and other male (ICD9: 187), 164-165
GIRLS (Generalized Iterative Record Linkage System)
Use in linking records, 4-5
Gum (ICD9: 143), 40-43

Heart
See Other thoracic organs
Histiocytic tissue
See Lymphoid tissue
Histologically verified
See HV%
History of the Registry
Early cancer incidence surveys, 1-2
Sources for early surveys, 1-2
Quality of early data, 1-2
HMRI (Hospital Medical Records Institute)
Source of data, 2
Hodgkin's disease (ICD9: 201), 198-201
Hospital Medical Records Institute, 2
HV% (Percent Histologically Verified)
By site, 22-23
Definition, 11
Hypopharynx (ICD9: 148), 60-63

ICD (International Classification of Diseases)
Conversion of early revisions to ninth revision, 4
Fourth digit, 6, 14
Full or abbreviated title, 16
ICDO-M (International Classification of Diseases for Oncology - Morphology)
Change of OCR coding from MOTNAC to ICDO-M, 4
Ill-defined sites
See Other and ill-defined sites
Information, request for, 14
International Classification of Diseases
See ICD
Intrahepatic bile ducts
See Liver

Kidney (ICD9: 189), 170-173

Larynx (ICD9: 161), 112-115
Leukaemia
Lymphoid (ICD9: 204), 210-213
Myeloid (ICD9: 205), 214-217
Monocytic (ICD9: 206), 218-221
Other (ICD9: 207), 222-225
Unspecified cell type (ICD9: 208), 226-229
Linkage
GIRLS computer program, 4
Linkage category and grey area, 4-5
Non-NYSIIS linkage, 5
NYSIIS code, 4
Quality control of linkage, 5
Reduction in manual linkage workload, 4-5
Role in generating incidence data, 4
Sequence of linkages, 5
Weights used to link, 4-5

Lip (ICD9: 140) 28-31
Liver (ICD9: 155), 88-91
Lung (ICD9: 162), 116-119
Lymphoid tissue
Other neoplasm of lymphoid and histiocytic tissue (ICD9: 202), 202-205
Leukaemia
See Leukaemia, Lymphoid
Lymphosarcoma and reticulosarcoma (ICD9: 200), 194-197
Lymph glands, secondary
See Secondary neoplasms

M/I% (Mortality: Incidence Ratio as a percentage), 22-23
Definition of, 11
By site, 22-23
Manual of Tumour Nomenclature and Coding
See MOTNAC
Mediastinum
See Other thoracic organs
Melanoma of skin (ICD9: 172), 140-143
Middle ear
See Nasal cavities
Ministry of Health
See MOH
MOH (Ministry of Health)
Source of information for OCF, 1-2
Mortality/Incidence Ratio
See M/I%
MOTNAC (Manual Of Tumour Nomenclature And Coding)
Change of morphology coding from MOTNAC to ICDO-M, 4
Mouth
Floor of (ICD9: 144), 44-47
Other parts of (ICD9: 145), 48-51
Unspecified parts of (ICD9: 145), 48-51
Multiple myeloma (ICD9: 203), 206-209

n (frequency of cancers)
By sites, all ages, 20-21
Single site, by age group
See site of interest
Nasal cavities (ICD9: 160), 108-111
Nasopharynx (ICD9: 147), 56-59
Nervous system unspecified (ICD9: 192), 182-185
New York State Intelligence Identification System
See NYSIIS
Nose
See Nasal cavities
NYSIIS (New York State Individual Identification System)
For linking records, 4-5

OCF (Ontario Cancer Foundation)
See OCTRF
OCR (Ontario Cancer Registry)
See OCTRF
OCTRF (Ontario Cancer Treatment and Research Foundation)
Incorporation date, 1
Financial support, 1
History, 1-2
Regional centres, 1
Oesophagus (ICD9: 150), 68-71
OHIP (Ontario Health Insurance Plan)

Source of data for OCF, 2
Ontario
Descriptive data, 1
Major ethnic groups, 1
Ontario Cancer Foundation
See OCTRF
Ontario Cancer Institute, 1
Ontario Cancer Registry
See OCTRF
Ontario Health Insurance Plan
See OHIP
Oropharynx (ICD9: 146), 52-55
Other and ill-defined sites (ICD9: 195), 230-233
Other thoracic organs (ICD9: 164)
Thymus, Heart, and Mediastinum, 124-127
Ovary (ICD9: 183), 156-157

Pancreas (ICD9: 157), 96-99
Pathology reports
Artifact in, 8
Penis
See Genital organs
Percent Death Certificate Only
See DCO%
Percent Distribution of cancers
See %Dist
Percent Histologically Verified
See HV%
Peritoneum
Other and ill defined sites within the peritoneum (ICD9: 159), 104-107
Retroperitoneum and Peritoneum (ICD9: 158), 100-103
Pharynx unspecified
Other and ill-defined sites within the lip, oral cavity, and pharynx (ICD9: 149), 64-67
Placenta (ICD9: 181), 152-153
Planning
Use of OCF data in, 7-8
Pleura (ICD9: 163), 120-123
PMH (Princess Margaret Hospital)
Establishment and role of, 1
Populations
Canada, 1971, 17
Ontario, 18-19
World Standard, 17
Production of data, 8-9
Prostate (ICD9: 185), 160-161

Quality of data
Indices of, 22-23

Radiation Therapy, 1
Rates
Lag time in producing, 8
RCC (Regional Cancer Centres)
List of centres, 1
Rectosigmoid junction
See Rectum
Rectum, rectosigmoid junction (ICD9: 154), 84-87
Respiratory
Other and ill defined sites within the respiratory system and intrathoracic organs (ICD9: 165), 128-131
Secondary malignant neoplasm of respiratory and digestive systems (ICD9: 197), 238-241
Reticulosarcoma
See Lymphosarcoma
Retroperitoneum
See Peritoneum

Salivary Glands (ICD9: 142), 36-39
Secondary neoplasms
Lymph nodes (ICD9: 196), 234-237
Respiratory and digestive (ICD9: 197), 238-241
Other specified sites (ICD9: 198), 242-245
Sequence of site specific tables, 12
Sinuses
See Nasal cavities
Skin
See Melanoma of skin
Small intestine (ICD9: 152), 76-79
Specific Rates
See ASR
Standard populations
See Populations
Standardized Rates
See ASR or AAR
Statistics, method of calculation, 12-13
Stomach (ICD9: 151), 72-75

Testis (ICD9: 186), 162-163
Thymus
See Other thoracic organs
Thyroid Gland (ICD9: 193), 186-189
Trachea
See Lung
Tongue (ICD9: 141), 32-35

Unspecified
Digestive (ICD9: 159), 104-107
Female genital (ICD9: 184), 158-159
Leukaemia, unspecified cell type (ICD9: 208), 226-229
Lymph nodes (ICD9: 196), 234-237
Mouth (ICD9: 145), 48-51
Nervous system (ICD9: 192), 182-185
Pharynx (ICD9: 149), 64-67
Respiratory (ICD9: 165), 128-131
Site (ICD9: 199), 246-249
Urinary organs (ICD9: 189), 170-173
Uterus (ICD9: 179), 148-149
Urinary organs
See Kidney
Uterine adnexa
See Ovary
Uterus
Part unspecified (ICD9: 179), 148-149
Body of uterus (ICD9: 182), 154-155